

# Genetic Variability, Heritability and Correlation Study of Physiological and Yield Traits in Relation to Heat Tolerance in Wheat (*Triticum aestivum* L.)

A Nishant Bhanu\*, B Arun and VK Mishra

Department of Genetics and Plant Breeding, Banaras Hindu University, India

Received: December 12, 2017; Published: January 08, 2018

\*Corresponding author: A Nishant Bhanu, Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, India; Email: nishant.bhanu@gmail.com

## Abstract

Twenty genotypes of wheat were undertaken for assessment of direct selection parameters (variability, heritability, and genetic advance), indirect selection parameter (correlation) and their relation with heat tolerance. This investigation was carried out at the Agriculture Research Farm of Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during the Rabi season of 2011-12 and 2012-13 in randomized block design with 3 replications. Observations were recorded on grain yield and its related traits, viz., days to heading, days to anthesis, days to maturity, grains per spike, 1000 grain weight. Besides these, chlorophyll content at heading and at anthesis stages as well as canopy temperature depression (CTD) from vegetative stage to dough stages were studied. All the traits like days to heading, days to anthesis, days to maturity and 1000 grain weight exhibited high heritability, which was coupled with high genetic advance as per cent of mean was recorded for 1000 grain weight and yield per plot. That indicated effectiveness of selection for the improvement of these traits.

The 1000 grain weight, i.e., test weight, as the most important yield attribute was found positively and significantly associated with CTD at early and late milking stages suggesting that higher seed weight is possible in varieties which show high CTD, i.e., low canopy temperature in heat stress conditions. Among the component traits, days to heading had positive and significant correlation with days to anthesis, and days to maturity. Interestingly, this result suggests that for the selection of early heading genotypes, days to anthesis and days to maturity should be invariably considered for wheat improvement. Further, plot yield had non-significantly negative association with days to maturity indicating that the early genotypes in semi-arid condition are capable of escaping the terminal heat stress, which is the major yield reducing factor. The twenty genotypes studied, represented considerable and diverse variability for different traits evaluated and significant associations on yield which could be utilized in breeding programme for varietal improvement in wheat.

**Keywords:** Wheat; Canopy Temperature Deficit; 1000 Grain Weight; Days to Heading; Days to Anthesis

**Abbreviations:** CTD: Canopy Temperature Depression; PCV: Phenotypic Coefficient of Variation; GCV: Genotypic Coefficient of Variation

## Introduction

Wheat (*Triticum aestivum* L.), a cereal grass of *Graminae* (*Poaceae*) family and of the genus *Triticum*, is the world's largest cereal crop. It has been described as the 'King of Cereals' because of the acreage it occupies, high productivity and the prominent position it holds in the food grain trade. It occupies an inimitable position in human life as it is the major source of food and energy with a large number of end use products like chapathi, bread, biscuits, pasta and is also a good source of fodder for animals. Wheat is grown on 215.9 million hectares throughout the world, which produces 725.91 million tons of grain [1]. It provides, on an average, one fifth of total calorific input to the world population (FAO, 2008). Wheat in India is grown over an area of 29.86 million ha producing 95.85 million tones [2].

With the enormous increase in the population, the demand for wheat is also increasing such that it is expected that by 2020 the country's demand for wheat will be 88 million tonnes. Wheat production in recent years has reached a plateau where further increase in productivity and yield is somewhat difficult. The constraints of biotic and abiotic stresses are a greater challenge to the crop scientists and plant breeders. Heat stress is a key abiotic stress affecting crop and cereal production in all regions of the Indian wheat belt [3]. Field data suggests that yield losses can be in the order of 190 kg/ha for every one degree rise in average temperature and in some situations having a more severe effect on yield loss than water [4,5]. Heat tolerance is a complex phenomenon and difficult to measure. Many selection criteria based on morpho-physiological traits were

reported to be associated with performance under heat stress in wheat [6].

Heat tolerant metabolism reported to be indicated by longer leaf chlorophyll retention, canopy temperature depression, photosynthetic rate, leaf senescence. The other traits like biomass, 1000-grain weight and grain yield itself are also highly responsive to heat tolerance. Many investigators reported the traits like high tillering capacity, relatively higher grain weight, acceptable spike fertility, higher spike number, grains per spike, early ground cover etc. to be associated with yield under heat stress and tolerance. High temperature stress in the recent years has been major factor affecting the wheat productivity especially, in the arid and semi arid regions of the world [7]. The earth temperature is constantly increasing. The cultivation of wheat is limited by temperature at both ends of the cropping season. Since plant tolerance to temperature stress is heritable, selection and breeding can be used to improve this trait.

Breeding cultivars with improved tolerance to temperature, however, is a difficult task because of the limited understanding of the genetic and physiological basis of heat tolerance in plants; confounding effects of heat and drought stress; limited understanding of what stages of plant growth can be used as selection criteria and lack of efficient and accurate tests to select for the component physiological mechanisms that give heat tolerance. The relationship between the morpho-physiological traits associated with heat tolerance is very much important in selecting suitable selection criteria for heat tolerance. The association between the traits related to heat tolerance could be better understood through correlation study both at genotypic and phenotypic levels [8]. On the basis of nature and magnitude of genotypic and phenotypic correlations, breeding programs may measure such traits to assist in the selection of heat tolerant genotypes. Therefore, the present investigation was undertaken to examine the relationships between heat tolerant traits and yield.

## Materials and Methods

Twenty genotypes were undertaken for assessment of direct selection parameters (variability, heritability, and genetic advance),

**Table 1:** Analysis of variance (ANOVA) for all the characters under study in twenty lines of wheat.

Mean Sum of Square (MSS)											
Source	Degree of freedom	Days to heading	Days to anthesis	Days to maturity	1000 grain weight	CT 1	CT 2	CT 3	Chlorophyll content 1	Chlorophyll Content 2	Yield per plot
Replicaton	2	0.03	0.03	0.12	0.01	0.09	0.53	5.76	14.65	6.45	87.79
Treatment	19	11**	9.51**	8.36**	26.56**	0.25	0.85	6.05*	7.22**	3.62*	44870.36**
Error	38	3.23	2.92	0.08	4.12	0.69	0.66	0.80	6.33	2.97	15274.15

\*Significant at  $P \leq 0.05$  level of significance

\*\*Significant at  $P \leq 0.01$  level of significance

Estimates of phenotypic variance ( $\sigma^2_p$ ) were higher than genotypic variance ( $\sigma^2_g$ ) (Table 2). A wide range of ( $\sigma^2_p$ ) was observed for days to heading, chlorophyll contents, days to anthesis, grain weight and plot yield while moderate range of variance showed by the traits days to maturity and canopy temperature at

indirect selection parameter (correlation) and their relation with heat tolerance. This investigation was carried out at the Agriculture Research Farm of Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during the Rabi season of 2011-12 and 2012-13 in randomized block design with 3 replications. Seeds of each genotype were sown in unit plot size of 3m long with 6 rows 23 cm apart. Standard agronomic practices were adopted for both experiments. Observations were recorded on grain yield and its related traits, viz., days to heading, days to anthesis, days to maturity, grains per spike, 1000-grain weight. Besides these, chlorophyll content at heading and at anthesis stages by Minolta SPAD-502 chlorophyll meter as well as canopy temperature depression (CTD) was measured by using a hand held infrared thermometer based equipment (Sixth Sense LT-300) from vegetative stage to dough stages were studied.

Data were recorded on five random plants from each entry per replication for all the above mentioned traits except CTD, where single reading itself gives the average of scores of leaves. The data were subjected to the analysis of variance [9] and further, biometrical procedures were followed to estimate genotypic and phenotypic coefficient of variation [10], heritability in broad sense [11], genetic advance [12] and correlation [13].

## Result and Discussion

The analysis of variance for different characters is presented in Table 1. The results showed highly significant difference for most of the characters under study among twenty genotypes. This suggested that there is an inherent genetic difference among the genotypes. The estimates of range, mean, phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), heritability (broad sense) and genetic advance are presented in Table 2. The characters like 1000 grain weight and plot yield exhibited higher values of phenotypic variance while other traits also showed similar trend. This suggests that there is a good scope of selection for these factors [14-16]. The phenotypic coefficient of variation was higher than their respective genotypic coefficient of variation for all the traits [17-22]. The 1000 grain weight and yield per plot showed maximum phenotypic coefficient of variation. This variation was minimum for days to [23].

different physiological stages. The result of phenotypic coefficient of variation revealed that yield per plot exhibited highest phenotypic coefficient of variation (9.336) followed by 1000 grain weight (9.28). Also chlorophyll content, days to heading and CT at different stages showed wide phenotypic coefficient of variation while,

the remaining characters showed moderate to low phenotypic coefficient of variation. Genotypic coefficient of variation was high for 1000 grain weight (7.45) followed by yield per plot (5.85). Also chlorophyll content and CTD at different stages showed

wide genotypic coefficient of variation. And remaining characters showed moderate to low genotypic coefficient of variation [24-26]. The minimum value of genotypic coefficient of variation was observed for days to anthesis, days to maturity and CT [14,27].

**Table 2:** Range, Mean, SEm ( $\pm$ ), Phenotypic and genotypic variance, PCV, GCV, heritability, genetic advance as percentage of mean for all physiological and yield traits under study in wheat.

Mean Sum of Square (MSS)											
Characters		Days to heading	Days to anthesis	Days to maturity	1000 grain weight	CT 1	CT 2	CT 3	Chlorophyll content 1	Chlorophyll content 2	Yield per plot
Range	Max	70.16	75	103	32.27	23.93	23.13	34.2	41.73	39.18	1445.55
	Min	77.33	81.83	109.33	42.31	24.96	25.06	42.26	47.58	43.62	1974.78
Grand mean		77.16	77.66	104.78	36.72	24.55	23.69	39.11	45.03	42.07	1698.25
SEm ( $\pm$ )		1.46	1.39	0.23	1.66	0.67	0.66	2.03	1.4	1.31	100.9
Phenotypic variance		5.82	5.12	2.84	11.61	0.54	0.72	6.16	4.37	2.92	25139.55
Genotypic variance		2.59	2.19	2.75	7.47	-	0.063	-	1.42	0.349	9865.4
PCV (percent)		3.297	2.913	1.609	9.28	3	3.59	6.34	4.644	4.063	9.336
GCV (percent)		2.201	1.909	1.585	7.446	1.55	1.06	0.5	2.651	1.405	5.849
Heritability (percent)		0.446	0.43	0.971	0.644	-0.2672	0.0868	-0.0088	0.3258	0.1197	0.3924
Genetic advance as percent of mean		3.02	2.578	3.219	12.3	-1.65	0.64	-0.12	3.12	1	7.547

Thus, the traits with high GCV are to be considered during selection. The heritability ranged from 11.97 percent to 97.1 percent. High heritability estimate was exhibited by days to maturity (97.1), 1000 grain weight (64.4), days to heading (44.6) and days to anthesis (43). The remaining traits showed moderate levels of heritability. The estimate of genetic advance as percent of mean was highest for yield per plant 1000 grain weight (12.30) followed by yield per plot (7.55), and the traits like days to heading, days to anthesis, days to maturity and chlorophyll content showed moderate advance while the CT showed less genetic advance [15,28-30]. High heritability coupled with high genetic advance as percent of mean was recorded for 1000 grain weight and yield per plot indicating effectiveness of selection for the improvement of these traits [16,24,28,31,32]. Also CTD and chlorophyll content at different stages exhibited moderate heritability coupled with

moderate genetic advance indicating their effectiveness for selection of genotypes tolerant to high temperature [8,33].

Genotypic and phenotypic correlation coefficients were worked out among all the characters under study and the results have been planted in Tables 3 & 4, respectively. The phenotypic correlation coefficient was highly significant and positive for yield per plant with 1000 grain weight and grains per spike. Also highly significant correlation for yield per plot was obtained with canopy temperature depression at anthesis and late milking stages and significant correlation for yield per plot per plot was observed with spike length and spikelet per spike though at 5 percent level of significance. Chlorophyll content at anthesis and CTD at heading and late milking also showed significant and positive correlation with yield. A significant but negative correlation for yield per plot with days to maturity was observed.

**Table 3:** Genotypic ( $r_g$ ) correlation coefficients of all the physiological and yield traits.

Characters	Days to heading	Days to anthesis	Days to maturity	1000 grain weight	CT 1	CT 2	CT 3	Chlorophyll content 1	Chlorophyll Content 2	Yield per plot
Days to heading	1.0000	0.9813**	0.6620**	-0.8040**	-0.2171*	-0.8671**	0.5799*	0.3151	0.6225**	-0.4218*
Days to anthesis		1.0000	0.7013**	-0.8648	0.1954*	0.8369**	0.6647**	0.3616	0.7439**	-0.3829*
Days to maturity			1.0000	-0.2156	-0.0080	-0.5851*	0.5074*	0.0574	0.3135*	-0.2016*
1000 grain weight				1.0000	-0.2819*	0.4327*	-0.7076**	0.0740**	-0.1442*	0.3941*
CT 1					1.0000	0.0184	0.0176	-0.1000	0.1192*	0.36671*
CT 2						1.0000	0.4256*	0.3803*	0.1198*	0.4322*
CT 3							1.0000	0.2476*	0.1218*	0.0201
Chlorophyll content 1								1.0000	1.00	0.4867*

Chlorophyll content 2									1.0000	0.4243*
Yield per plot										1.0000

\*\*Significant at  $P \leq 0.01$  level of significance

\*Significant at  $P \leq 0.05$  level of significance

**Table 4:** Phenotypic ( $r_p$ ) correlation coefficients of all the physiological and yield traits.

Characters	Days to heading	Days to anthesis	Days to maturity	1000 grain weight (g)	CT 1	CT 2	CT 3	Chlorophyll content 1	Chlorophyll Content 2	Yield per plot
Days to heading	1.000	0.9820**	0.4662**	-0.4535*	-0.1893	-0.2886	0.0518	-0.0377	0.0146	-0.3108*
Days to anthesis		1.000	0.4902**	-0.4571*	-0.2044	-0.2982	0.0488	0.0011	0.0361	-0.2881*
Days to maturity			1.000	-0.1987*	-0.0345	-0.1430	0.1063	0.0370	0.0998	-0.1375*
1000 grain weight				1.000	-0.1088	0.0435	-0.0389	0.0875	0.0253	0.2009*
CT 1					1.000	0.1030	0.0757	0.1370	0.1387	0.5769*
CT 2						1.000	-	-0.0762	0.1306	0.6897**
CT 3							1.000	0.1113	0.7940**	0.6487**
Chlorophyll content 1								1.000	0.6925*	0.4654*
Chlorophyll content 2									1.000	0.7632**
Yield per plot										1.000

\*\*Significant at  $P \leq 0.01$  level of significance

\*Significant at  $P \leq 0.05$  level of significant

Among component traits positive and highly significant correlation was observed for days to heading with days to anthesis, days to maturity and CTD at late milking stage. Days to anthesis exhibited positive and highly significant correlation with days to maturity and CTD at late milking stage. This suggests that higher seed weight is possible in varieties which show high CTD i.e., low canopy temperature in heat stress conditions. Therefore the selection of such genotypes is useful in breeding for heat tolerance. A significant and positive correlation was observed for chlorophyll content at anthesis with days to heading, days to anthesis and plant height [34-36]. The plot yield had non-significantly negative association with days to maturity [37,38]. This indicates that the early genotypes in semi arid condition are capable of escaping the terminal heat stress which is the major yield reduction factor.

## Conclusion

Climate change and sustainable development are interlinked problems that pose a serious challenge to humanity. High temperature stress in the recent years has been major factor affecting the wheat productivity especially, in the arid and semi-arid regions of the world. The earth temperature is constantly increasing. It can be concluded from the above mentioned findings and details that the genotypes used in the study exhibited considerable variability and association among themselves for various traits under study. Thus, providing the ample scope for selection of elite genotypes

and those which are tolerant to heat stress. Several approaches like QTL mapping and association genetic analysis should be useful in an effort to relate specific alleles to trait variation and to identify candidate genes.

## Acknowledgement

We are thankful to the UPCAR Lucknow, for providing genotypic materials and Department of Genetics and Plant Breeding, Institute of Agricultural Sciences for providing all the necessary support to conduct this research programme.

## References

1. Foreign Agricultural Services, USDA, 2015.
2. FAO (2008) FAO Statistical Yearbook 2008, Statistical Division, Food and Agricultural Organization, Rome.
3. Joshi AK, Mishra B, Chatrath R, Ferrara GO, Singh RP (2007) Wheat improvement in India: Present status, emerging challenges and future prospects. *Euphytica* 157: 431-446.
4. Kuchel H, Williams K, Langridge P, Eagles HA, Jefferies SP (2007) Genetic dissection of grain yield in bread wheat. II. QTL-by-environment interaction. *Theoretical and Applied Genetics* 115: 1015-1027.
5. Bennett D, Izanloo A, Reynolds M, Kuchel H, Langridge P, et al. (2012) Genetic dissection of grain yield and physical grain quality in bread wheat (*Triticum aestivum* L.) under water-limited environments. *Theoretical and Applied Genetics* 125: 255-271.



6. Deswal RK, Grakh SS, Berwal KK (1996) Genetic variability and characters association between grain yield and its components in wheat. *Annals of Biology (Ludhiana)* 12(2): 221-224.
7. Rane J, Nagarajan S (2004) High temperature index-for field evaluation of heat tolerance in wheat varieties. *Agricultural Systems* 79(2): 243-255.
8. Reynolds MP, Balota M, Delgado MIB, Amani I, Fischer RA (1994) Physiological and morphological traits associated with spring wheat yield under hot, irrigated conditions. *Australian Journal of Plant Physiology* 21(6): 717-730.
9. Panse VG, Sukhatme PG (1967) *Statistical methods for agricultural workers* (2<sup>nd</sup> edn.). pp. 381, ICAR, New Delhi, India.
10. Burton GW (1952) Quantitative inheritance in grasses. *Proceedings of 6<sup>th</sup> International Grassland Congress* 1: 227-283.
11. Burton GW, Devane (1953) Estimating heritability in tall fescue from replicated clonal material. *Journal of Agronomy* 45(3): 473-481.
12. Johnson HW, Robinson HF, Comstock RE (1955) Genotypic and phenotypic correlations in soybeans and their implication in selection. *Agronomy Journal* 47: 477-483.
13. Robinson HF, Comstock RE, PH Harvey (1951) Genotypic and phenotypic correlation's in wheat and their implications in selection. *Agronomy Journal* 43: 282-287.
14. Singh JP, Prasad S, Singh KN, Singh R (2007) Screening of heat tolerant wheat varieties by membrane thermo stability index in relation to yield and yield attributing traits. *International Journal of Plant Sciences (Muzaffarnagar)* 2(2): 159-165.
15. Dwivedi VK, Kumar S, Tyagi NK (2003) Genetic variability in some metric traits and it contribution to yield in wheat (*Triticum aestivum* L.). *Progressive Agriculture* 3(1/2): 152-153.
16. Cheema NM, Mian MA, Ihsan M, Rabbani G, Mahmood A (2006) Studies on variability and some genetic parameters in spring wheat. *Pakistan Journal of Agricultural Sciences* 43(1/2): 32-35.
17. Singh NB, Singh YP, Tiwari LP, Bahar J, Singh VPN, et al. (2008) Morphophysiological traits as index of screening wheat genotypes for thermotolerance. *Indian Journal of Plant Physiology* 13(3): 307-311.
18. Dixit SK (1990) Variability pattern in durum wheat under different sowing path analysis inland races of bread wheat from South Western Iran. *Euphytica* 41: 183-190.
19. Sharma AK, Garg DK (2002) Genetic variability in wheat (*Triticum aestivum*) crosses under different environments. *Annals of Agricultural Research* 23(3): 497-499.
20. Kumar R, Gupta RK, Visistha NK, Singh MK (2013) Character association and path analysis for quantitative traits in bread wheat. *Environment and Ecology* 31 (1A): 393-398.
21. Jha PB, Singh, SP, Singh DN (2001) Genetic variability for polygenic traits in late sown wheat genotypes. *Annals Agricultural Research* 22(1): 34-36.
22. Tripathi GP, Parde NS, Zate DK, Lal GM (2015) Genetic variability and heritability studies on bread wheat (*Triticum aestivum* L.). *International Journal of Plant Sciences (Muzaffarnagar)* 10(1/2): 57-59.
23. Porwal Y, Prasad SVS, Billore M (2006) Association of path analysis for grain yield, its components and quality traits in bread wheat. *National Journal of Plant Improvement* 8(1): 24-27.
24. Panwar D, Singh I (2000) Genetic variability and character association of some yield components in winter x spring nursery of wheat. *Advances in Plant Science* 8(1): 95-99.
25. Bergale S, Halkar AS, Ruwali KN, Prasad SVS, Mridulla B (2001) Genetic variability, diversity and association of quantitative traits with grain yield in bread wheat. *Madras Agricultural Journal* 88(7-9): 457-461.
26. Arvind K, Raj S (2015) Genetic variability in relation to qualitative and quantitative traits in wheat (*Triticum aestivum* L.). *Agrica Journal* 3(2): 23-28.
27. Chaturvedi BK, Gupta RR (1995) Selection parameters for some grain and quality attributes in spring wheat (*T. aestivum* L.). *Agricultural Science Digest* 15(4): 186-190.
28. Atta YA, Akhter BMJ, Lateef PZ (2008) Genetic variability, association and diversity studies in wheat (*Triticum aestivum* L.) germplasm. *Pakistan Journal of Botany* 40(5): 2087-2097.
29. Gahalain SS, Bisht D (2009) Interrelationships and path coefficient analysis in wheat germplasm of Kumaun Himalayas. *Vegetos* 22(2): 19-26.
30. Zecevic V, Boskovic J, Dimitrijevic M, Petrovic S (2010) Genetic and phenotypic variability of yield components in wheat (*Triticum aestivum* L.). *Bulgarian Journal of Agricultural Science* 16(4): 4.
31. Masood MS, Mujahid, Hashmi NI (1986) Variability studies in wheat under rainfed conditions. *Pakistan Journal of Agricultural Research* 23: 244-247.
32. Bhoite KD, Rasal RN, Gadekar DA (2008) Genetic variability, heritability and genetic advance in durum wheat (*Triticum durum* L.). *Journal of Maharashtra Agricultural Universities* 33(1): 102-103.
33. Pierre SC, Crossa J, Manes Y, Reynolds MP (2010) Gene action of canopy temperature in bread wheat under diverse environments. *Theory of Applied Genetics* 120: 1107-1117.
34. Khaliq IN, Parveen, MA Chowdhry (2004) Correlation and Path Coefficient Analyses in bread wheat. *International Journal of Agriculture and Biology* 6: 633-635.
35. Khan AJ, Azam F, Ali A, Tariq M, Amin M (2005) Inter-relationship and path coefficient analysis for biometric traits in drought tolerant wheat (*Triticum aestivum* L.). *Asian Journal of Plant Sciences* 4(5): 540-543.
36. Naqvi S, Zhu CF, Farre G, Ramessar K, Bassie L, et al. (2009) Transgenic multivitamin corn through biofortification of endosperm with three vitamins representing three distinct metabolic pathways. *Proceedings of National Academy of Science* 106: 7762-7767.
37. Mann CE (1994) Early vigor in wheat: a useful character for heat tolerance selection. In: Saunders DA & Hettel GP (Eds.), *Proceedings of the International Congress on Wheat in heat stressed environments: Irrigated. Dry areas and rice farming systems*. Mexico.
38. Mascagni HJ, Harrison SA (2003) Wheat Yield and Maturity: Influence of Variety and Planting Date. *Louisiana Agriculture* 46(1): 12-13.



#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<http://biomedres.us/>