

Dysplastic Lumbar Spondylolysis Presenting As Hyperventilation Syndrome in a Young Man



Shin-Tsu Chang*^{1,2}

¹Department of Physical Medicine and Rehabilitation, Taichung Veterans General Hospital, Taichung City, Taiwan

²Department of Physical Medicine and Rehabilitation, School of Medicine, National Defense Medical Center, Taipei City, Taiwan

Received: May 09, 2018; Published: May 15, 2018

***Corresponding author:** Shin-Tsu Chang, Department of Physical Medicine and Rehabilitation, School of Medicine, National Defense Medical Center, Taichung Veterans General Hospital, 1650 Taiwan Boulevard Section 4, Taichung, No.161, Sec. 6, Minquan East Road, Neihu District, Taipei, Taiwan, Tel: 886-4-23741350; Fax: 886-945-605523; Email: ccdivlaser1959@gmail.com

Abstract

Many causes contribute to hyperventilation; however, the pathophysiology involved is still unknown, and the skeletal origins of the condition have scarcely been examined. We present a case of a 20-year-old man with a history of hyperventilation and subtle lower back pain due to spondylolysis of the fifth lumbar vertebra with grade I spondylolisthesis. The duration and frequency of hyperventilation increased after performing physical modality and abated after discontinuing physical modality. We discuss the relationship and associated pathophysiology between hyperventilation and spondylolysis, which is the first report in the literature. We suggest that physical therapy might be harmful for some patients with hyperventilation, because the hyperventilation might be a clinical manifestation of dysplastic lumbar spondylolysis.

Keywords: Hyperventilation; Spondylolysis; Pars; Pain; Mechanoreceptor

Introduction

Hyperventilation, defined as breathing in excess of metabolic requirements, is associated with a reduction in arterial PaCO₂ and respiratory alkalosis, as well as a wide range of symptoms. Many causes contribute to hyperventilation syndrome; however, the pathophysiology involved is still unknown, and the skeletal origins of the condition have scarcely been examined [1]. We present the case of a 20-year-old man with a history of hyperventilation and subtle lower back pain due to spondylolysis of the fifth lumbar vertebra with grade I spondylolisthesis. The duration and frequency of hyperventilation increased after performing physical modality (short wave diathermy and electric stimulation therapy). After discontinuing physical therapy, the hyperventilation abated. We discuss the relationship and associated pathophysiology between hyperventilation and spondylolysis.

Case Presentation

A 20-year-old soldier without history of major trauma experienced subtle lower back pain for 2 years since 2008. Half a year prior, he suffered from shortness of breath with near-syncope for several minutes after standing guard in 2010. A series of examinations did not show any specific findings, except a low PaCO₂; hyperventilation syndrome was considered. Due to intermittent subtle lower back pain, he was referred to our rehabilitation clinic

for help on 2010/10/12. Physical examination revealed pain during trunk extension and poor balance when standing on a single leg. Plain X-ray films of the lumbar spine showed bilateral spondylolysis of the fifth lumbar vertebra, together with grade I spondylolisthesis, and the ensuing computer tomography confirmed the lumbar pathology (Figure 1).




Figure 1: Images of computer tomography of the lumbar spine. Left Reconstructural Sagittal view shows lesion of pars interarticularis of fifth lumbar with grade I spondylolisthesis. Right Axial view shows bilateral lesion of pars interarticularis of fifth lumbar in the level of the corresponding line in left.

Physical therapy, including modality intervention (short wave diathermy and interferential current therapy), was prescribed for

his back pain. During rehabilitation, the back pain worsened and shortness of breath with cold sweat developed. Thereafter, the duration and frequency of hyperventilation increased several times during rehabilitation therapy, but laboratory data of blood and electrolytes were normal at every examination. Considering the risk of hyperventilation and the likelihood of falling down and acquiring an injury, we discontinued rehabilitation and instead prescribed analgesic medication and local injections. Thereafter, the lower back pain improved progressively, and no further hyperventilation attacks were reported.

Discussion

Many factors contribute to the hyperventilation syndrome, for example, organic, respiratory, central nervous system, physiologic/psychologic, and psychogenic factors. Of these factors, physiologic dysfunction derived from pain, exercise, anxiety, and excitement can induce hyperventilation, particularly when combined with other factors, and reduce PaCO₂ below the average level of 20 mmHg [2]. The precise pathophysiology of hyperventilation is unknown. It has been documented that hyperventilation induces increase in central excitability because hyperventilation does not alter peripheral conduction velocity [3]. Research in heart rate variability also confirmed that hyperventilation is strongly associated with neuronal excitability in human cortex [4]. On the other hand, involvement of peripheral nervous system might a role in pathophysiology because the increase in the excitability of cutaneous nerves and motor axons of median nerve matches the increase in compound sensory and muscle potentials of thenar muscles in patients with hyperventilation [5].

Spondylolysis is frequently observed in radiographic films as a defect in the pars interarticularis of the vertebral arch and may either occur asymptotically or be associated with significant lower back pain [6]. The clinical manifestations of symptomatic spondylolysis include focal back pain that may or may not radiate into the buttock or proximal lower limbs [7]. The onset of pain can be gradual or start after an acute injury, and mild symptoms can present for some time following an acute exacerbation [8]. Painful pars lesions are more common in young athletes than other populations [9]. The observation that the duration and frequency of hyperventilation increased during rehabilitation therapy in the patient presented here is of interest. The patient experienced worse back pain and subsequent hyperventilation during rehabilitation, which may be attributed to physiological dysfunction induced by modality therapy. With respect to modality therapy, an increase in body temperature due to thermal therapy or a change in mechanoreceptors due to electric stimulation may have contributed to the development of hyperventilation. However, it is difficult to confirm this hypothesis, because no similar case has been reported in the literature.

Various explanations have been proposed for the anatomical origin of lower back pain and lower limb pain associated with lumbar spondylolysis with/without spondylolisthesis. Detail study regarding lumbar fibrocartilage masses of scar tissue at the site of pars fracture, presence of nociceptive free nerve endings and mechanoreceptors has been confirmed, and those are assumed to sense position and movement [10]. Additional features in the nerve

endings of lumbar facet joint capsules have been demonstrated by using morphological and quantitative methods, and these features are quite different from those found in fibrocartilage masses at the site of pars lesions [11]. The complex of facet joint, pars and adjacent soft tissues is richly innervated by the medial branches of the dorsal rami. The facet capsule contains low-threshold mechanoreceptors and nociceptors, which can be hypersensitized by an inflammatory process [7,12], that alert the brain to undue stress in joint-related tissues.

We believe that the nociceptors surrounding the pars could be hypersensitized by stress or chronic derangement, like our case. Figure 2 demonstrates a possible mechanism for the relationship between hyperventilation and spondylolysis. Therapeutic interventions for spondylolysis vary, but complete bony healing of the lesion site is the final goal. Many authors have recommended activity restriction for spondylolytic patients [13,14] and some suggest anti-lordotic bracing to maintain lumbar lordosis [15]. Improving the dynamic stability and fine control of the lumbar spine, which may be achieved by the strengthening of core muscles, hamstring stretching, pelvic tilts, and specific stabilizing exercises, has been advocated by others [16].




Figure 2: Schematic model of the interaction between hyperventilation and spondylolysis.

Conclusion

This is the first report describing a relationship between hyperventilation and spondylolysis. The hyperventilation observed in our patient may initially have been caused by the subtle back pain associated with spondylolysis, and subsequent worsening factors might have included the additional stress or pain induced by modality therapy or inappropriate exercise intensity. We suggest that physical therapy may be harmful for some patients with hyperventilation, because the hyperventilation might be a clinical manifestation of dysplastic lumbar spondylolysis.

References

1. Beeckmans N, Vermeersch A, Lysens R, Van Wambeke P, Goossens N, et al. (2016) The presence of respiratory disorders in individuals with low back pain: A systematic review. *Man Ther* 26: 77-86.
2. Gardner WN (1996) The pathophysiology of hyperventilation disorders. *Chest* 109: 516-534.
3. Seyal M, Mull B, Gage B (1998) Increased excitability of human corticospinal system with hyperventilation. *Electroencephalogr ClinNeurophysiol* 3: 263-267.

4. Neginhal R, Herur A, Ambi U, Ankad R, Badami S, et al. (2017) Effect of hyperventilation on excitability of the human cortex and cardiac autonomic variability. *Indian J Physiol Pharmacol* 61: 7-13.
5. Macefield G, Burke D (1991) Paraesthesiae and tetany induced by voluntary hyperventilation. Increased excitability of human cutaneous and motor axons. *Brain* 114: 527-540.
6. Standaert CJ, Herring SA (2003) Spondylolysis: a critical review. *Br J Sports Med* 34: 415-422.
7. Van Kleef M, Vanelderden P, Cohen SP, Lataster A, Van Zundert J, et al. (2010) Pain originating from the lumbar facet joints. *Pain Pract* 10: 459-469.
8. Micheli LJ, Wood R (1995) Back pain in young athletes: significant differences from adults in causes and patterns. *Arch Pediatr Adolesc Med* 149: 15-18.
9. Soler T, Calderón C (2000) The prevalence of spondylolysis in the Spanish elite athlete. *Am J Sports Med* 28: 57-62.
10. Hasegawa S, Yamamoto H, Morisawa Y, Michinaka Y (1999) A study of mechanoreceptors in fibrocartilage masses in the defect of pars interarticularis. *J Orthop Sci* 4: 413-420.
11. Schneiderman GA, McLain RF, Hambly MF, Nielsen SL (1995) The pars defect as a pain source. A histologic study. *Spine (Phila Pa 1976)* 20: 1761-1764.
12. Cavanaugh JM, Lu Y, Chen C, Kallakuri S (2006) Pain generation in lumbar and cervical facet joints. *J Bone Joint Surg (Am)* 88: 63-67.
13. Standaert CJ, Herring SA, Halpern B, King O (2000) Spondylolysis. *Phys Med Rehabil Clin N Am* 11: 785-803.
14. Garet M, Reiman MP, Mathers J, Sylvain J (2013) Nonoperative treatment in lumbar spondylolysis and spondylolisthesis: a systematic review. *Sports Health* 5: 225-232.
15. Iwamoto J, Takeda T, Wakano K (2004) Returning athletes with severe low back pain and spondylolysis to original sporting activities with conservative treatment. *Scand J Med Sci Sports* 14: 346-351.
16. O'Sullivan PB, Phytty GD, Twomey LT, Allison GT (1997) Evaluation of specific stabilizing exercise in the treatment of chronic low back pain with radiologic diagnosis of spondylolysis or spondylolisthesis. *Spine (Phila Pa 1976)* 22: 2959-2967.



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>