

In Vitro Antibacterial Properties of Ethanol Extract from *Salvia Officinalis* (L.) Plant Growing Wild in Kosovo



Arben Haziri¹, Fatmir Faiku¹, Arben Mehmeti², Kemajl Kurteshi¹, Imer Haziri², Ibrahim Rudhani^{3*}

¹Mathematical and Natural Science Faculty, University of Prishtina, Kosovo

²Agricultural and Veterinary Faculty, University of Prishtina, Kosovo

³Medicinal Faculty, university of Prishtina, Kosovo

Received: January 27, 2018; Published: February 09, 2018

*Corresponding author: Ibrahim Rudhani, Medicinal Faculty, university of Prishtina, 10000 Prishtina, Kosovo; Email: ibrahim.rudhani@uni-pr.edu

Abstract

In this study the antibacterial efficiency of ethanol extract from *Salvia officinalis* (L.) growing wild in Kosovo were examined. Ethanol extract was tested against two gram positive bacteria *Staphylococcus aureus* (clinical isolate), *Listeria monocytogenes* (clinical isolate) and one gram negative bacteria *Escherichia coli* (clinical isolate). The antibacterial activity was determined by using agar disc diffusion method. The inhibition zone of extract was compared to that of penicillin G. The ethanol extract showed activity in all of the concentrations 1, 3 and 5 mg/mL towards *E. coli*, *S. aureus* and *L. Monocytogenes*. Ethanol extract of the plant with concentration 5 mg/mL showed a stronger antibacterial activity towards bacteria *S. aureus* with inhibition zone of 9 mm. The antibacterial activity of the *S. officinalis* (L.) was due to the presence of various secondary metabolites such as phenols and flavonoids. Hence, this plant can be used to discover bioactive natural products that may serve as leads in the development of new pharmaceuticals.

Keywords: *Salvia officinalis* (L.); Antibacterial activity; Agar disk diffusion method; Ethanol extract

Introduction

The heterogeneous relief of Kosovo with the diversity of soil types, are responsible for the heterogeneousness of the Kosovo flora [1]. The secondary metabolism of the plants which are responsible for the biological activities are terpenes, phenolics and nitrogen-containing compounds. The antibiotics drugs obtained from plants are of greater interest in comparison to synthetic ones. The use of natural antibiotics from plants does not induce the secondary health effects (abdominal pain, diarrhea, cramping, headache, nausea and vomiting etc). Plant products may be used as antibiotic alternatives and do not cause resistance in bacteria [2].

The genus *Salvia*, commonly called sage, is the largest member of Lamiaceae or mint family containing over 900 species throughout the world [3]. *Salvia officinalis* L., probably the most known species of the *Salvia* genus and commonly known as garden or red sage, is a perennial hardy sub-shrub native to Mediterranean regions and is one of the most popular medicinal and culinary herbs used in the world [4,5]. Sage water extracts has been traditionally used for the treatment of digestive and circulation disturbances, bronchitis, cough, asthma, angina, mouth and throat

inflammations, depression, excessive sweating, skin diseases, and many other diseases. In addition, they possess a number of biological activities including antiseptic, antibacterial, antioxidant, antiviral and antimycobacterial activities [6-19]. The antibacterial effect of the ethanol extracts, obtained from the medicinal plants, has been studied by a number of researchers in different parts of the world. Our research group was interested to analyze the biological activities of different medicinal plants extracts growing in different region of Kosovo. The aim of this study was to investigate the antibacterial activity of ethanol extracts of *Salvia officinalis* L. growing wild in Kosovo.

Experimental

Plant Material

The *S. officinalis* L. plant was collected in 2015 in Kosovo. Voucher specimens were deposited in the herbarium of the Department of Plant Protection, University of Prishtina. The aerial part of this medicinal plant were air dried at room temperature within three weeks.

Preparation of Plant Organic Extracts

The aerial part of *S. officinalis* (L.) was air-dried and then milled with a mixer. A piece of finely powdered material (200 g) was extracted with 70% ethanol ($\text{CH}_3\text{CH}_2\text{OH}$, 4 L) during a 24 h period (three times). After removing the solvent by vacuum rotary evaporator (EYELA N-1000, Japan) the crud product was dissolved in DMSO to prepare the solution with the concentration 1, 3 and 5 mg/mL. The solutions are used in subsequent experiments for testing their antibacterial. Etanol (analytical grade) for extraction were obtained from Sigma-Aldrich.

Antibacterial Activity

The antibacterial activity of the *S. officinalis* (L.) extracts obtained by using the ethanol organic solvents are determined applying the Kirby-Bayer [19] method or disk method (d = 5.5 mm, maximum capacity 10 mg). *S. officinalis* (L.) extracts were tested in an *in vitro* experiment against bacterial strains; *S. aureus* (clinical isolate with code 3319), *L. monocytogenes* (clinical isolate with code 2653) and *E. coli* (clinical isolate with code 2813). For the research we used three different concentration of ethanol extract, 1, 3 and 5 mg/mL in DMSO as solvent and then placed in a Petri dishes (d=15 cm). The disks were incubated at 37 °C for 48 h; the control was also maintained with penicillin G dissolved in DMSO in a similar manner.

Results and Discussion

During the antibacterial activity, we have tested the extracts from the plant *S. officinalis* (L.) against bacteria: *S. aureus* (positive Gram bacteria), *L. monocytogenes* (positive Gram bacteria) and *E. coli* (negative Gram bacteria-clinical isolate). Antibacterial activity of the ethanol extracts were compared with Penicillin G- a substance, which can be found in the market and have been used with concentration 1, 3 and 5 mg/ mL as a standard. The comparison is done in accordance to the measurements of the diameter of the inhibition zone that is formed. These results are shown in Table 1.

Extracts of ethanol (1, 3 and 5 mg/ mL) shows antibacterial activities against *S. aureus*, *L. monocytogenes* and *E. coli* (Table 1 and Figure 1). The extracts of ethanol resulted in a lower activity than the penicillin G of the same concentration against *E. coli* (Table 1 and Figure 1). The extract of ethanol with concentration of 3 mg/mL and 5 mg/mL has higher inhibition zone (7 mm) as the standard (6 mm) against *L. monocytogenese*. The extract with concentration 1 mg/mL resulted in lower activity than the penicillin G with the same concentration against *L. monocytogenese* (Table 1 and Figure 1). The extracts of ethanol with concentration of 5 mg/mL have the same inhibition zone as the standard (inhibition zone 6 mm) against *S. aureus* (Table 1 and Figure 1).




Figure 1: Antibacterial activity of ethanol extracts from *S. officinalis* (L.).

Table 1: Inhibition zones diameters of *S. officinalis* (L.) ethanol extracts.

Extracts	Concentration (mg/ml)	Inhibition zones diameters (mm)		
		<i>E. coli</i> (c. i.)	<i>L. monocytogenes</i> (c. i.)	<i>S. aureus</i> (c. i.)
Ethanol	1	5	2	4
	3	6	7	6
	5	6	7	9
Penicillin	1	7	3	6
	3	7	5	7
	5	8	6	9

Conclusion

S. officinalis (L.) is a medicinal plant used in folk medicine for different health illnesses. The aim of this research was to analyse the antibacterial activities of ethanol extracts from *S. officinalis* (L.) plant growing wild in Kosovo. Based on the results, the three concentrations of the extract showed a very good activity by inhibiting the growth of all the tested bacteria. Ethanol extract of the plant with concentration 5 mg/mL showed a stronger antibacterial activity towards bacteria *S. aureus*, *E. coli* and *L.monocytogenes* (clinical isolate). Extract with concentration 5 mg/mL showed a higher antibacterial activity against bacteria *L.monocytogenes*. Results obtained from ethanol extracts of *S. officinalis*(L.) for antibacterial activity are logical, based on numerous studies where these extracts were analyzed in the content of flavonoids, phenols, terpenes, alkaloids, etc., and these components are responsible for the biological activity.


References

- Mehmeti A, Sherifi E, Demaj A (2007) Medical Herbs (European Agency for Reconstruction), 1st edition, Prishtine, Kosovo 46.
- Gibbons S (2005) Plants as a source of bacterial resistance modulators and anti-infective agents *Phytochemistry* 4(1): 63-78.
- Hamidpour S, Hamidpour M, Shahlari M (2013) Sage: The functional novel natural medicine for preventing and curing chronic illnesses. *International Journal of Case Reports and Images* 4: 671-677.
- Grzegorzcy KI, Króllicka A, Wysokińska H (2006) Establishment of *Salvia officinalis* L. hairy root cultures for the production of rosmarinic acid. *Z Naturforsch C* 61(5-6): 351-356.
- Qnais EY, Abu-Dieyeh M, Abdulla FA, Abdalla SS (2010) The antinociceptive and antiinflammatory effects of *Salvia officinalis* leaf aqueous and butanol extracts. *Pharm Biol* 48(10): 1149-1156.
- Hamidpour R, Hamidpour S, Shahlari M (2014) Chemistry, Pharmacology, and Medicinal Property of Sage (*Salvia*) to Prevent and Cure Illnesses such as Obesity, Diabetes, Depression, Dementia, Lupus, Autism, Heart Disease, and Cancer. *J Tradit Complement Med* 4(2): 82-88.
- Yang Z, Kitano Y, Chiba K, Shibata N, Kurokawa H et al. (2001) Synthesis of variously oxidized abietanedieterpenes and their antibacterial activities against MRSA and VRE. *Bioorg Med Chem* 9(2): 347-356.
- Abd-Elmageed MM, Hussein BA (2008) Cytotoxicity and antimicrobial activity of *Salvia officinalis* L. Flowers. *Sudan JMS* 3(2): 127-130.
- Lima CF, Andrade PB, Seabra RM, Fernandes-Ferreira M, Pereira-Wilson C (2005) The drinking of a *Salvia officinalis* infusion improves liver antioxidant status in mice and rats *J Ethnopharmacol* 97(2): 383-389.
- Baricevic D, Sosa S, Della LR, Tubaro A, Simonovska B, et al. (2001) Topical antiinflammatory activity of *Salvia officinalis* L. leaves: The relevance of ursolic acid. *J Ethnopharmacol* 75(2-3): 125-132.
- Akkol EK, Goger F, Kosar M, Baser KHC (2008) Phenolic composition and biological activities of *Salvia halophila* and *Salvia virgata* from Turkey. *Food Chem* 108(3): 942-949.
- Çadirci E, Süleyman H, Gürbüz P, Kuruüzüm UA, Güvenalp Z et al. (2012) Antiinflammatory effects of different extracts from three *Salvia* species. *Turk J Biol* 36: 59-64.
- Tada M, Okuna K, Chiba K, Ohnishi E, Yoshiia T (1994) Antiviral diterpenes from *Salvia officinalis* *Phytochemistry* 35(2): 539-541.
- Smidling D, Mitic-Culafic D, Vukovic-Gacic B, Simic D, Knezevic-Vukcevic J (2008) Evaluation of antiviral activity of fractionated extracts of Sage *Salvia officinalis* L (Lamiaceae). *Arch BiolSci Belgrade* 60(3): 421-429.
- Fiore G, Nencini C, Cavallo F, Capasso A, Bader A, et al. (2006) *In vitro* antiproliferative effect of six *Salvia* species on human tumor cell lines. *Phytother Res* 20(8): 701-703.
- Ryu SY, Lee CO, Choi SU (1997) *In vitro* cytotoxicity of tanshinones from *Salvia miltiorrhiza* *Planta Med* 63(4): 339-342.
- ZareShahneh F, Valiyari S, Baradaran B, Abdolalizadeh J, Bandeghagh A, et al. (2013) Inhibitory and cytotoxic activities of *salvia officinalis* L. Extract on human lymphoma and leukemia cells by induction of apoptosis. *Adv Pharm Bull* 3(1): 51-55.
- Aşkun T, Başer KH., Tümen G, Kürkçüoğlu M (2010) Characterization of essential oils of some *Salvia* species and their antimycobacterial activities). *Turk J Biol* 34(1): 89-95.
- Barry L (1991) Procedure and Theoretical Consideration for Testing Antimicrobial Agents in Agar Media. 5th edition, William Wilkins, Baltimore, Maryland 8-10.



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <http://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<http://biomedres.us/>