

**Appendix 2:** Prevalence of Anaphylaxis in Children in Kuwait Questionnaire. Title: The Prevalence of Anaphylaxis in Children in Kuwait.

Instructions: Kindly answer this survey with the help of a parent or caregiver. Please tick  the box that best correspond to your answer and provide specific details where required. All information will be kept confidential. Thank you for your willingness to help us in this study.

IDENTIFICATION NUMBER:

DATE:

AGE:

GENDER:  MALE  FEMALENATIONALITY:  Kuwaiti  Indian  Pakistani  Lebanese  Syrian Egyptian  OthersI. History of acute onset of illness (minutes to several hours- up to 12 hours):  Yes  No

If yes, please tick the corresponding signs/symptoms below:

 Skin (hives, pruritus, flush)  Mucosal tissue (swollen lips, tongue, uvula) Both

II. Respiratory compromise.

 Dyspnea  Wheeze/Bronchospasm Hypoxemia  Stridor (difficult inspiration with sound)

III. Reduced blood pressure documented in the hospital.

 Hypotonia (collapse)  Syncope Incontinence

IV. Persistent gastrointestinal symptoms.

 Crampy abdominal pain  Vomiting

V. Age during the incident?

 0-1 year  2-4 years 5-11 years  11-17 years

VI. Number of times the incident happened?

 1  2 More than 2

VII. Have you ever used Adrenaline auto injector?

 Yes  NoVIII. What was the trigger of the allergy attack?  Known  Unknown

If known, kindly list down below.

 Insect Bite  Medicine Food:  Others Milk Eggs Nuts

Fish

Others

Wards count: 6015 excluding title, abstract, tables, figures