

Current Knowledge of TB Patients on Initiation of Treatment in Mettu Town



Dame Tesfaye¹ and Sileshi Dubale*²

¹Department of Nursing, Mettu University, Mettu Ethiopia

²Department of Pharmacy, Mettu University, Mettu Ethiopia

Received:  August 01, 2018; **Published:**  September 12, 2018

***Corresponding author:** Sileshi Dubale, Department of Nursing, Mettu University, Mettu Ethiopia, Ethiopia, Email: sileshi.dubale@gmail.com

Abstract

Background: Knowledge on causes & initiation time of TB treatment of the community is critical and improve or worsen the transmission of the disease and treatment outcomes. Lack of awareness about the disease contribute to how compliance to chemotherapy and preventive measures of the disease.

The objectives of this study were: To assess timeline of initiation of the treatment and current knowledge of patients in Mettu Town. A facility based descriptive cross-sectional study was conducted from April 28/2016 –May 29/2016 and all patients attending Mettu Karl hospital and Mettu health center were included. Data was compiled and analyzed by using Statistical package for Social Sciences software (SPSS) version 20.0. Statistical significance was defined at a level of 0.05 and data was described with a confidence interval of 95%.

Results: Forty-six (52.87%) of respondents were female and 48.5% of the patients belong to age group of 15-29. Fifty-four (62.07%) of respondents residing in Urban setting and 54.02% were Oromo and 43.68% of respondents can read and write. Thirty-three (37.93%) of respondents recognized the symptoms one month before they presented to health facility for the first time.

Conclusion and Recommendation: Significant numbers of the respondents were not knowledgeable and delayed in the initiation of the treatment. Appropriate intervention should be taken to reverse this problem.

Keywords: Treatment Initiation; Current Knowledge; Tuberculosis

Background

TB is a common and deadly infectious disease caused usually by mycobacterium tuberculosis in human. About 10% of latent infection eventually progress to active TB disease which, if left untreated kills about half of those infected. Once person develop active tuberculosis diseases, the symptoms of this disease may be mild for months and this leads to delays in seeking care and results in the transmission of bacteria from infected person to the others. Lack of understanding about tuberculosis, stigma toward the disease, inaccessibility of treatment is the main reason for delayed presentation; while lack of information and dissatisfaction of the treatment are the main reason of non-completion of the treatment. Perceived knowledge on the causes of tuberculosis and health seeking behavior among community members may reduce or increase the disease transmission. The spread of the disease increases due to failure to recognize the symptoms early, hence theses may delay the diagnosis of tuberculosis and treatment [1-5]. TB is re-emerging as a public health concern because of the low practices of peoples to prevent the transmission of TB, inadequate

health coverage, impact of HIV epidemic and negative attitude of patients on its transmission and treatment.

Countries with high particularly sub Saharan Africa have witnessed profound increase in the number of tuberculosis cases [6-9]. In Ethiopia TB is major public health problem with increase rate of TB cases of 2.6% each year. Among these 97.1% are new cases of all forms TB. Smear positive cases account 32% and smear negative and extra pulmonary TB are 34% and 32% respectively. Lack of adequate knowledge about TB and noncompliance to treatment result in treatment failure. The likelihood of successful treatment of tuberculosis depends on the extent to which patient complete the pressured treatment regimen (adherence and noncompliance) [10-12]. Better understanding of level of the knowledge and time of initiation of the treatment about TB patient help for care provider and administrate to identify the gap and to take relevant action. Therefore, the purpose of this study was to assess timely initiation of the treatment and their current knowledge toward TB for patient on treatment in Mettu Town.

Objectives

General objectives

To assess the initiation time of the treatment and current knowledge of patients in Mettu town

Specific Objective

- To assess the time line of initiation of the treatment of tuberculosis patient
- To determine current knowledge of tuberculosis patient
- To explore factors affecting time of initiation of TB treatment

Method and Participants

Study Area

The study was conducted in health facilities of Mettu town, from April 2016 to May 2016. Mettu town is found in south west Ethiopia, in Illu Aba Bor zone 600km away from Addis Ababa, the capital of the country.

Sample Size Determination

All TB patient attending Mettu Karl Hospital and Mettu health center during the study period was included as convenient sample size.

Data Collection Procedure

Standard self-administered questionnaire and interview was developed in English and then changed to Afaan Oromo (Oromo Language) and then to English language). Questionnaires were further modified from guidelines and previous study before actual data collection. Data collection was face to face interview by principal investigator as it gets direct information from respondents.

Data Quality Assurance Measures

The appropriately designed data collection instrument was used. Every day the collected data was reviewed and checked for completeness and consistency of the respondents. Pre- test was done on 5% of total population out of the study and necessary correction was made on the language clarity sequencing and

workability of questionnaires.

Data Analysis

The collected data was analyzed manually after it was edited and checks for completeness and consistency, for further analysis, it was analyzed using SPSS. Descriptive analysis was used of describe the percentage of knowledge of TB patient and treatment initiation of TB patient to the area. Frequency distribution was using for qualitative data.

Ethical Consideration

Ethical clearance was obtained from ethical review board of Mattu university, college of public health and medical science department of nursing. Letter of permission present to Mattu Karl hospital and health center TB follow up clinic and verbal informed consent was obtained from each study is explain to respondent. Confidentiality of information was assured, and privacy of the respondent was maintained.

Limitation of the Study

Since the patients on follow up attend health center by appointment some TB patients were not involved in the study during the study period. Shortage of time of data collection and lake of similar literature in the study area to compare and contrast the result

Results

Time Lines of Treatment

About timelines of treatment initiation, the time when patients recognized the first symptoms, frequency of health facility visit before diagnosis, delays for diagnosis, and delays for treatment initiation and treatment phase were interviewed. Concerning of the time when they recognized first symptoms, thirty-three (37.93%) of respondents recognized the symptoms one month before they presented to health facility for the first time. Fourth eight (55.17%) of respondents visit health facility only once before diagnosis of TB, 37(42.53%) were diagnosed for TB within a week after the first visit and 62.07% were start treatment immediately after diagnosis of TB. Of total of respondents, 62.07% were in the continuation phase (Table 1).

Table 1: Time lines of treatment initiation of respondents in Mettu town, 2016.

Variable		Frequency	%
When recognised 1 st Symptom	1 week back	5	5.75
	2 weeks back	9	10.34
	3 weeks back	12	13.79
	1 month back	33	37.93
	>1 month	28	32.18
Frequency of health facility visit	1 time	48	55.17
	2 times	29	33.33
	3 times	9	10.34
	4 times	1	1.15
Delays for diagnosis	<1 week	37	42.53
	1-2 weeks	35	40.23

	2-4 weeks	10	11.49
	<4 weeks	5	5.57
Delays for treatment initiation	Start immediately	54	62.07
	<1 week	28	32.18
	1-2 week back	5	5.75
Treatment phase	Intensive phase	33	37.93
	Continuation phase	54	62.07

Knowledge about TB

Concerning knowledge of respondent’s questions about causes, signs and symptoms, transmission, curability, side effects of anti TB drugs, prevention and importance of completion of the treatment were asked. About cause of TB 47 (54.02%) of respondents were respond that TB caused by bacteria (Figure 1). Concerning perceived knowledge about sign and symptoms, transmission, curability, anti TB drugs treatment, prevention of TB was also asked and analyzed. Eighty-one 93.1% of respondents were respond as they know TB can transmitted and how it can be transmitted. Concerning anti TB drugs treatment 85.06% of respondents were know as they may die if they don’t treat and 91.95% of them know anti TB treatment was taken for 6 months. All the respondents had been taking the drugs once per day. About preventability of TB 86.21% know as TB

is preventable disease. To sum up 13 knowledge-based question were asked and 82.76% were answered the question above mean (knowledgeable) and 17.24% were not knowledgeable (Table 2).




Figure 1: Knowledge about causes of TB of Mettu.

Table 2: Knowledge of respondents about TB and its treatment in Mettu Town, 2016.

Variables		Frequency	%	
Signs & symptoms	Cough	83	95.4	
	Night sweating	37	42.53	
	Loss of appetite	20	23	
	Bloody sputum	15	17.24	
	Lymph node enlargement	17	19.54	
	Fever	43	49.45	
	Weight loss	28	32.18	
	Easily fatigability	55	63.22	
Perceived knowledge on transmission of TB	Coughing	79	97.53	
	Sneezing	44	54.32	
	Using the same utensil	47	58.02	
	Yes (93.1%)			
	Sleeping together	11	13.52	
	Drinking raw milk	5	6.17	
	Drinking contaminated water	2	2.47	
	Other	2	2.47	
Perceived knowledge on Anti-TB treatment	No	6	6.9	
	If not treated	I can lead normal life	1	1.15
		I will be cured by help of god	3	3.45
		I may die	74	85.06
		I can transmit the disease	11	12.64
		I will be disabled	1	1.15
	Total duration of the treatment	4 months	2	2.3
		6 months	80	91.95

		8 months	5	5.75
Perceived knowledge on prevention	Yes	Avoid coughing in front of people	73	97.33
		Not using the same utensil	37	49.33
		Drinking boiled milk	4	5.33
	No		12	13.79
Importance of completion of the treatment	To cure completely		85	97.7
	To avoid recurrence		14	16.9
	To decrease transmission		5	5.75TB

Table 3: Factor Affecting Time of Initiation of TB Treatment in Mettu Town, 2016.

Variables		Frequency	%	AOR (95%CI)	P-Values	Intercept
Sex	Male	41	47.13	1		0.029
	Female	46	52.87	2.4(1.9-2.10)	0.003	
Age	15-29 years	42	48.5	0.329(0.215-0.409)	0.806	
	30-49 years	29	33.34	1		
	50+ years	16	18.4	3.73(2.14-3.502)	0.067	2.74
Residence	Urban	54	62.07	1		1.60
	Rural	33	37.93	4.00(2.57-6.45)	0.001	
Marital status	Married	55	63.22	1		0.61
	Single	21	24.14	2	0.78	
	Divorced	6	6.9	0.97(0.09-0.23)	0.56	
	Widowed	5	5.75	2.30(1.9-4.56)	0.047	
Occupation	Farmer	36	41.38		0.001	0.34
	Merchant	20	23		0.67	
	Government employed	7	8.05	1		
	Private employed	15	17.25		0.08	
Educational status	Student	9	10.34		0.46	
	Illiterate	31	35.63	10.76(5.019-13.09)	0.001	3.56
	Read and write	18	20.7	4.06(2.34-6.04)	0.023	
	Grade 1-8	15	17.24	1.62(0.87-2.03)	0.09	
	Grade 9-10	19	21.8	0.2(0.12-0.97)	0.06	
	10+	4	4.6	1		

Factor Affecting Time of Initiation of TB Drug

The adjusted odd ratio (AOR) was calculated to identify associated factors and their significant. The analyzed risk factors are sex, age, patient residence, marital status, occupation and educational status of the Patients. Age of the patients, marital status and occupation had no such significant effect on knowledge and TB treatment dalliance. Sex, patient living, and educational status are significantly affecting patient TB treatment knowledge and they take lion's share for dalliance of TB diagnosis and treatment. The odds of Females delayed from seeking TB treatment is 2.4(1.9-2.10) times than male in Mettu town with P-value of 0.003. Patients residing rural areas are delayed from seeking TB diagnosis and treatment 4.00(2.57-6.45) times than urban resident patients with p-value of 0.001. Illiterate patients were 10.76(5.019-13.09) times

delayed from TB diagnosis and treatment than educated patient up to grade ten and above, with significant values (P<0.001) (Tables 2 & 3).

Discussion

This study provided information regarding TB patients, timelines of treatment initiation, knowledge about tuberculosis and its treatment and factor affecting time of initiation of TB drug. In this study 88.5% of TB patients belong to economically productive age group (15-54years) which is slightly higher than the finding documented in WHO report which was 82% of cases. This difference might show that the finding of this study is recent and it was also done in specific area and my total population or respondents also not comparable with that study since WHO report is global study [13] and 54.02% of respondents responded that

bacteria as the cause of tuberculosis whereas study done in south west Ethiopia in 2010 showed that 33.7% of Tb suspected patients had knowledge on the cause the disease [14]. This study shows that 95.4% respondents mentioned that cough is one sign and symptom of TB which is slightly higher than the study done in Tanzania which is 72% of them know cough as one sign and symptom of TB [15].

Knowledge about route of the transmission of the disease is another important factor in TB prevention and control problem. In our study 93.1% of respondents responded as they know as TB can be transmitted, which is also slightly higher than study done in southwest Ethiopia which indicate 83.8% of patients have knowledge about route of transmission [16]. This difference is may be due to our study was conducted in the place where majority of study population were from urban area and they may had access of health information more. About curability of the disease 97.7% of respondents were know that as TB is curable disease and 91.95% of them know anti TB treatment was taken for 6 months. This finding was the same with study conducted in Tanzania in which all respondents knew that TB was curable disease [17]. In this study 86.21% of respondents knew that as Tb is preventable disease with preventive method. This is slightly higher than study conducted in south west Ethiopia in which 82.3% of respondents know as Tb can be prevented and 69.9% know that not coughing /sneezing in front of other people as the main preventive measure [18]. Sex, Patient Residence and Educational status are significantly affecting Patient TB treatment knowledge and are significant factors for dalliance of TB diagnosis and treatment

Conclusion

More than half of patient visited health facility only once before diagnosis. Majority of them started the treatment immediately after diagnosis and were in continuation phase [19]. Most of them know as Tb can be transmitted and coughing in front of people and using the same utensil were common made of transmission known by respondents. Almost all respondents know as TB is curable disease and requires 6 months treatment more than half know that as anti TB treatment has no side effect. Majority of respondents know as TB is preventable disease and almost all of respondents respond avoidant direct coughing in front of people as the common preventive methods [20]. Patients living rural areas and illiterate patients were delayed from seeking TB diagnosis and treatment.

Recommendation

Based on the finding of the study the following recommendation was given. The Mettu health center and Mettu Karl hospital responsible bodies should make an effort to implement planned and well-organized health information. As early diagnosis and treatment initiation is important in TB control, attention should be given in all health facility. The focus of health education for public should mainly on cause, route of transmission, side effects of anti TB treatment, sign and symptoms and preventive methods. In service training for health care provider on TB prevention and control should be strengthened to update their knowledge and skills. Further study should be done across the country on related

topics. Focus should be given more for rural residence and un educated patients during health education

Acknowledgement

The investigators would like to acknowledge Mettu University for providing necessary resources.

Conflicts of Interest

No any conflicts of interest since the source for fund is from Mettu University for academic purpose.

References

1. WHO (2011) Tuberculosis fact sheet No. 104.
2. Cassie Thomas (2002) A literature review of problem of delayed presentation for the treatment and non-completion of treatment for tuberculosis in less developed countries and ways of addressing these problems using particular implementation of DOTs strategy. *Journal of management in medicine* 16(4-5): 371-400.
3. Auer C, Sarol J Jr, Tanner M, Weiss M (2000) Health seeking and perceived cause of tuberculosis among patients in manila. *Journal of Tropical medicine and international health* 5(9): 648-655.
4. Dye C, Scheele S, Dolin P, Pathania V, Ravigione MC (1999) Global burden of tuberculosis, estimated incidence, prevalence and mortality by country. *Journal of American Medical Association* 282(7): 677-686.
5. WHO (2013) global tuberculosis control. Geneva, USA, 08: 324-326.
6. Espinal MA, Laszlo A, Simonsen L, Boulahbal F, Kim SJ, et al. (2001) Global trends in resistance to anti tuberculosis drugs. *New England Journal of Medicine* 344(17): 1294-1303.
7. Michael KW, Belachew T, Jira C (2004) Tuberculosis defaulters from the "dots" regiemen in Jimma zone, south west Ethiopia. *Ethiop Med J* 42(4): 247-253.
8. WHO (2013) WHO global tuberculosis control geneva, USA.
9. World health organization (2003) Treatment of tuberculosis; guide lines for national programmers. Geneva, Switzerland, world health organization.
10. Global tuberculosis control. WHO report (2000) Geneva, World Helath Organization.
11. Booulue chimbartil, Fungladda W, Kaewkungwal J, Silachamroom U (2008) Treatment seeking behaviors and improvement in adherence to treatment regimen of tuberculosis patients using intensive triad-model program. *South East Asian J Trop Med Public Health* 39(3): 526-541.
12. Jane Mc, Harry JM Fnkenflige, Valerie M, Anna P Nieboer (2010) TB treatment initiation and adherence in South African community influenced move by perception than by knowledge of tuberculosis. *BMC public health* 10: 72.
13. AM Kilalel, Mushi AK, Lema LA, Kunda J, Makasi CE et al. (2008) Perception of tuberculosis and treatment seeking behavior in Ilala and Kinondoni Municipalities in Tanzania *Journal of Health research* 10(2): 89-94.
14. (2005) Central statistical Agency, Ethiopia Demographic and health survey 2005, Addis Ababa Ethiopia.
15. Mulenga C (2010) pulmonary TB patient's attitude to seek health.
16. Abebe G, Amare Deribew, Ludwig Apers, Kifle Woldemichael, Jaffer Shiffa (2010) Knowledge, health seeking behavior and perceived stigma.
17. Mohammed A, yousif (2007) knowledge of tuberculosis. A survey among tuberculosis patients in Omdurman, Sudan, *Sudanese Journal of public health* January 2(1).

18. World health organization (2011) global tuberculosis' control. Geneva, Switzerland.
19. Hang L Rider (2011) Epidemiology disease of Tb control in Paris and Berne by international union against TB and lung disease.
20. PE Mangeshol, E Sharo, WH Makunde, Keto GB, Mandara CI (2007) Community knowledge attitude and practice towards tuberculosis and its treatment in MP wapwa District. central Tanzania health research Bulletin 9(1): 38-43.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2018.09.001733](https://doi.org/10.26717/BJSTR.2018.09.001733)

Sileshi Dubale. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>