

Urethral Structure Length ≥ 2 cm is Significantly Associated with Lower Urethroplasty Success Rate, Results of our Large Case Series





Cristina Falavolti*¹, Roberto Giulianelli², Ervin Shehu¹, Pouria Alijani¹, Riccardo Giovannone¹, Tommasangelo Petitti³, Stuart F Boyd⁴ and Maurizio Buscarini³

¹Villa Betania Hospital, Italy

²Villa Claudia Hospital, Italy

³University Campus Biomedico, Italy

⁴University of Southern California, USA

Received:  September 29, 2018; Published:  October 09, 2018

*Corresponding author: Cristina Falavolti, Villa Betania Hospital, Via Pio IV 42 00165 Rome, Italy

Abstract

Urethral stricture is a relatively common disease in men and remains a reconstructive difficulty for urologists. It is associated with an unequivocal negative impact on the quality of life. The etiology is variable and the incidence is increasing in recent years with more frequent application of endoscopic instruments and indwelling catheters. Some of the most common etiologies of stricture disease include lichen sclerosis (LS), trauma, iatrogenesis, and infection [1-2]. Urethroplasty is considered the gold standard treatment for definitive correction of the disease reaching a success rate of 90%. Most of these strictures are short enough to make end-to-end urethral opposition. Normally, grafts are used when the strictures are too long or when patient already underwent a previously failed reconstruction attempt. The buccal mucosa is the most commonly used graft source [3]. The aim of this study was to investigate the correlation between urethral stricture etiology and length with the success rate in a large double-center series using the same surgical technique.

Materials and Methods

We retrospectively collected the data of 448 consecutive patients affected by urethral stricture who received surgery at University of Southern California of Los Angeles (USC) and University of Rome (UCBM) from 1998 to 2014. Patients were categorized based on the length of stricture in three different groups: group A 1-1.9 cm; group B 2-3cm and group C >3.1 cm. Urethral stricture etiologies and features (site and length) are showed in Table 1. Patients underwent urethroplasty according to the same technique. We performed end-to-end anastomosis for stricture <2cm (group A) and ventral buccal mucosa graft urethroplasty for patients with stricture >2cm (groups B and C). Comparative outcomes between the groups (by length) were assessed using analysis of a chi-square test. A logistic regression was used to calculate the odd ratio of success compared to the length. Statistical analysis was performed using STATA version 14.0 with 0.05 set as the level of significance. Database was analyzed to understand the impact of different urethral stricture lengths and etiologies on success rates. We collected data about one-year follow-up. All patients underwent urethral-cystoscopy and flowmetry at 1-6-12 month. Success was set as absence of urinary flow obstruction (no sign of endoscopic stricture and Q max > 10 ml/sec).

Results

Urethral strictures characteristics are showed in Table 1. Statistical results indicated that iatrogenic urethral strictures showed better urethroplasty results. Success rate significantly decreases for patients affected by urethral stricture ≥ 2 cm ($p < 0.05$) (Table 2). Success rate odd ratio of group 2 (stricture 2-3 cm) compared to group 1 (stricture 1-1.9 cm) is 0.26 ($p < 0.001$). Furthermore, success rate odd ratio of group 3 (stricture >2 cm) compared to group 1 is 0.17 ($p < 0.01$). There is no difference in the probability of success rate after surgery between group 2 and group 3 (Table 3).

Conclusion and Discussion

Urethral stricture is a relatively common condition and the prevalence in the US is estimated between approximately 200/100,000 in younger men to >600/100,000 in men older than 65 [4]. Traditionally, the most common treatment for urethral stricture is urethral dilatation and/or direct vision internal urethrotomy (DIVU) or Sachse. These treatments are preferred because they are simple with low morbidity [5]. After a first short period of symptoms improvements, urethral dilatation or urethrotomy show

a high number of recurrences. Often the efficacy of endoscopic procedures is very low. Pansadoro et al. [6] evaluated 224 patients with anterior urethral stricture and showed a high recurrence rate that improved according to the number of urethrotomy performed. They found a recurrence rate after first, second and third procedure of 58%, 82%, and 100% respectively. Also, Heyns et al found that the recurrence rate was 100% after the second endoscopic procedure [7]. The major part of patients of our series underwent previous

direct visual internal urethrotomy (DVIU) with high number of recurrences. That is why dr. Hamilton Russel introduced End-to-end (EE) urethroplasty in 1919. This procedure is still ideally suited for bulbar stricture with ≤ 3 cm long [8]. In 1993, a buccal mucosa graft (BMG) was introduced for the treatment of longer urethral stricture (> 3 cm) [9]. Urethroplasty success rate reaches 90% in many series for both techniques (excision and augmentation).

Table 1: STRICTURE LENGTH – groups A. 1.0-1.9 cm, B. 2.0-3.0 cm, C. >3.1 cm.

Etiology	N. of pts	<45 yrs old	>45 yrs old	Bulbar	Penile	Panurethral	Posterior	Previous direct visual internal urethrotomy (DVIU)
				(n. patients)	(n. patients)	(n. patients)	(n. patients)	(n. patients)
Perineal trauma	24	15	9	24 (A:6;B:9;C:9)	0	0	0	10
Urethral catheterization	51	14	37	25 (A:10;B:6;C:9)	10 (A:8;B:2;C:0)	16 (A:0;B:0;C:)	0	25
Idiopathic	79	30	49	51 (A:33;B:11;C:7)	12 (A:10;B:2;C:0)	16 (A:0; B:0; C:16)	0	39
Post prostatectomy	51	6	45	42 (A:21;B:12;C:9)	4 (A:4; B:0; C:0)	0	5 (A:0;B:3)	36
TUR	67	3	64	46 (A:29;B:11;C:6)	19 (A:15;B:4;C:0)	2 (A:0;B:0;C:)	0	43
Urethritis	14	11	3	12 (A:11; B:1; C:0)	2 (A:2; B:0; C:0)	0	0	3
Cystoscopy	7	1	6	7 (A:6; B:1)	0	0	0	0
Hypospadias	40	28	12	6 (A:1; B:2; C:3)	28 (A:21;B:6;C:1)	6 (A:0;B:0;C:)	0	22
Pelvic fracture	41	33	8	1 (A:1; B:0; C:0)	0	0	40 (A:15;B:16)	3
BXO	32	10	22	0	21 (A:0;B:10;C:11)	11 (A:0;B:0;C:)	0	28
Lichen								
Brachy therapy	28	0	28	21 (A:11;B:10;C:0)	7 (A:0; B:4; C:3)	0	0	21
Penile fracture	14	1	2	0	14 (A:14; B:0; C:0)	0	0	7

Note: BXO: Balanitis xerotica obliterans

Table 2: Success rate according to urethral stricture length.

Length	Success Rate		Total
	Yes	No	
1	24	194	218
	11.01	88.99	
2	35	75	110
	31.82	68.18	
3	49	71	120
	40.83	59.17	

Total	108	340	448
	24.11	75.89	100.00

Table 3: ODD Ratio of surgery success according to stricture length.

Success	Odds Ratio	Std. Err.	z	P> z	[95% Conf. Interval]	
Length						
1	.2650957	.0789639	-4.46	0.000	.1478618	.4752799
2	.1792552	.0511157	-6.03	0.000	.1025053	.3134708
Cons	8.083333	1.74909	9.66	0.000	5.289389	12.35309

For excisional urethroplasty, the diseased segment is excised, and the healthy ends anastomosed. For augmentation urethroplasty, the diseased segment is incised and the lumen widened by augmenting the incised urethra with a graft or flap of substitute epithelium [10]. Generally, management for short urethral stricture use end-to-end urethroplasty and for long urethral stricture use graft [11] but the surgical approach is determined by the etiology, localization, extension and previous treatment of stricture as well as the availability of local skin. Urethral strictures are a multifactorial issue. With increased application of endoscopic instruments and indwelling catheters, the incidence of iatrogenic urethral injury has increased obviously in recent years. In the literature, idiopathic and iatrogenic causes for urethral strictures are more common in the developed world, and each account for 33% of patients, respectively [12]. Inflammatory and traumatic causes account for 15% and 19% of strictures, respectively. In our series iatrogenic urethral strictures showed better urethroplasty results. According to the literature, we chose an end-to-end anastomosis for stricture <2 cm and BMG urethroplasty for patients with stricture > 2cm. We demonstrated that success rate significantly decreases for patients affected by urethral stricture ≥ 2 cm even though we used a BMG procedure.

Buccal mucosa is superior because it is easier to harvest, more readily available, resistant to infection with favorable tissue characteristics (thick epithelium, high content of elastic fibers and thin lamina propria) [13]. Several studies showed that BMG had superior success rate than end-to-end urethroplasty also in the management of short bulbar stricture [14]. In conclusion, urethral stricture is a significant issue and the incidence is increasing with advancing age (decreased tissue blood supply and ischemia have been proposed as a possible mechanism) and with urethral instrumentation. The most important feature in urethral stricture's treatment and success rate is the length. For stricture longer than 1 cm, endoscopic treatment should be avoided because prior urethrotomy is found to be a risk factor for failure and can make the urethroplasty more difficult [15]. Urethroplasty, in general, is an excellent option for a urethral stricture. A calculated approach is that initial urethrotomy followed by urethroplasty is the most cost-effective approach if there is recurrence of the stricture, unless the success rate of urethrotomy was likely to be inferior to 35% [16]. In our series, we reached very good result in the treatment of stricture shorter than 2 cm using an EE urethroplasty. BMG urethroplasty is a safe and effective procedure but if the urethral stricture is longer than 2 cm the success rate inevitably reduces.

References

- Stein DM, Thum DJ, Barbagli G (2013) A geographic analysis of male urethral stricture aetiology and location. *BJU Int* 112: 830-834.
- Wood DN, Andrich DE, Greenwell TJ, Mundy AR (2006) Standing the test of time: the long-term results of urethroplasty. *World J Urol* 24(3): 250-254.
- Dubey D, Kumar A, Mandhani A, Srivasta A, Kapoor R, et al. (2005) Buccal mucosa urethroplasty: a versatile technique for all urethral segments. *BJU Int* 95(4): 625-629.
- Santucci RA, Joyce GF, Wise M (2007) Male urethral stricture disease. *J Urol* 177(5): 1667-1674.
- Santucci RA, Mario LA, Mc Aninch JW (2002) Anastomotic urethroplasty for bulbar urethral stricture: analysis of 168 patients. *J Urol* 167(4): 1715-1719.
- Pansadoro V, Emiliozzi P (1996) Internal urethrotomy in the management of anterior urethral strictures: long-term followup. *J Urol* 156(1): 73-75.
- Heyns CF, Steenkamp JW, De Kock ML (1998) Treatment of male urethral strictures: is repeated dilation or internal urethrotomy useful? *J Urol* 160(2): 356-358.
- Gomez RG (2008) Stricture excision and primary anastomosis for anterior urethral stricture. *Urethral reconstructive surgery*. United States: Humana Press p. 107-118.
- Barbagli G (2008) Buccal mucosal graft urethroplasty. *Urethral reconstructive surgery*. United States: Humana Press pp. 119-135.
- Hudak SJ (2015) Use of overlapping buccal mucosa graft urethroplasty for complex anterior urethral strictures. *Transl Androl Urol* 4(1): 16-21.
- Mac Donald MF, Santucci RA (2005) Review and treatment algorithm of open surgical techniques for management of urethral stricture. *J Urol* 165(1): 9-15.
- Smith TG (2016) Current management of urethral stricture disease. *Indian J Urol* 32(1): 27-33.
- Bullock TL, Brandes SB (2007) Adult anterior urethral stricture: a national practice patterns surveys of board certified urologists in the united states. *J Urol* 177(2): 685-690.
- Angermeier KW (2008) Anterior urethral reconstruction: ventral grafts. *Textbook of reconstructive urologic surgery*. United Kingdom: Informa pp. 459-465.
- Breyer BN, Mc Aninch JW, Whitson JM (2010) Multivariate analysis of risk factors for long-term urethroplasty outcome. *J Urol* 183(2): 613-617.
- Wright JL, Wessells H, Nathens AB, Hollingworth W (2006) What is the most cost-effective treatment for 1 to 2-cm bulbar urethral strictures: societal approach using decision analysis. *Urology* 67(5): 889-893.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2018.09.001854](https://doi.org/10.26717/BJSTR.2018.09.001854)

Cristina Falavolti. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>