



Symptom Severity of Major Depressive Disorder: Associations with Functional Disability, And Prevalence of Co-Morbid Illnesses



Morteza Alibakhshikenari*

Shahid Beheshti University, Tehran, Iran

Received:  November 06, 2018; Published:  November 15, 2018

*Corresponding author: Morteza Alibakhshikenari, Shahid Beheshti University, Iran

Abstract

Objective: The objective of this study was to investigate the associations of symptom severity of major depressive disorder with functional disability and medical co-morbidities among Singapore residents.

Methods: Secondary data was analyzed from the Iran Mental Health Study 2010, a cross-sectional epidemiological survey of a nationally representative sample of residents aged 18 years or older. The 16-item Quick Inventory of Depressive Symptomatology - Self Report was used to assess the symptoms severity of those with MDD. Functional disability was assessed with Sheehan Disability Scale. Information on existing medical co-morbidities was also collected.

Results: Amongst 162 Singapore residents with 12-month MDD, 43.2% had mild to moderate symptoms and 56.8% had severe symptoms. Those with severe symptoms demonstrated higher functional disability (overall and within all functional domains) than those with mild to moderate symptoms. Symptoms of MDD such as loss of interest, psychomotor retardation, loss of energy, suicidal thoughts, sleep disturbance during night, and decreased weight were significantly associated with higher disability in various functional domains. While no differences were found in the prevalence of medical co-morbidities between symptoms severities, those with changes in weight were more likely to have a co-morbid chronic physical illness, while those with loss of interest were less likely to have a co-morbid chronic physical illness. Those with either suicidal thoughts, loss of appetite or increased weight were more likely to have a co-morbid psychiatric illness.

Conclusion: Our study found that symptom severity is associated with functional disability among Iran residents with MDD, highlighting the importance of managing MDD symptoms to minimize the impact on functioning. Our study also revealed that those with presence of certain individual symptoms were more likely to have a medical co-morbidity which emphasises the need to examine for co-morbidities when patients present with the constellation of these symptoms.

Keywords: Major Depressive Disorder; Depression; Symptom Severity; Disability; Co-Morbidity

Abbreviations: MDD: Major Depressive Disorder; GAD: Generalized Anxiety Disorder; SMHS: Singapore Mental Health Study; QIDS-SR16: Quick Inventory of Depressive Symptomatology - Self Report; CIDI: Composite International Diagnostic Interview; DSM-IV: Diagnostic and Statistical Manual of Mental Disorder, Fourth Edition; HAMD: Hamilton Rating Scale of Depression; SDS: Sheehan Disability Scale; SAS: Statistical Analysis Software

Introduction

Major depressive disorder (MDD) is a mood disorder accompanied by a multitude of symptoms including prolonged periods of sadness, lack of attention, fatigue, loss of pleasure and interest in everyday activities, withdrawal from social interaction, irregular sleep patterns, feeling of hopelessness and suicidal thoughts [1]. A WHO [2] report estimated that 322 million people, i.e., 4.4% of the global population, suffered from a depressive disorder in 2015 making it the leading cause of disability worldwide.

Prevalence of depressive disorders across United States of America, United Kingdom, China and Japan were 5.9%, 4.5%, 4.2% and 4.2% respectively [2]. MDD is associated with various adverse outcomes such as functional disability, impaired quality of life and suicidal behaviors [3-5]. Severity of MDD is an important albeit complex construct comprising of symptom severity, impairment and criteria count (Lux et al.). Studies have found that MDD patients who exhibit more severe symptoms were associated with greater functional

disability [4,6]. Fried and Nesse [7] further explored the association between individual depressive symptoms and the various domains of functional disability.

They found that both sadness and lack of attention were detrimental to all domains of functioning, while sleeping difficulties, loss of interest in everyday activities and self-blame were strongly associated with disability in work, social activities and relationship, respectively [7]. MDD is well-known to be highly co-morbid with other psychiatric disorders such as generalized anxiety disorder (GAD) and substance use disorder [8-9]. MDD patients who displayed more severe symptoms were more likely to have at least one co-morbid psychiatric disorder [8]. A cross-sectional study in Thailand found that MDD patients who displayed high suicidal risk may be more vulnerable to co-morbid anxiety and psychotic disorder [9]. Furthermore, MDD patients have a high prevalence of co-morbid chronic physical illnesses such as heart diseases, diabetes and hypertension [1,10]. It was suggested that MDD patients were more likely to engage in unhealthy lifestyle and diets that may lead to the development of these chronic physical illnesses [11]. Pereira et al. [12] study found that individuals with fewer numbers of depressive symptoms participated more frequently in physical activity.

In addition, Appelhans et al. [13] found that MDD patients who demonstrated more severe symptoms were associated with poorer diet quality which was characterized by unhealthy dietary pattern such as reduced intake of vegetables and fruits, and excessive intake of unhealthy fats, sodium and alcohol. Singapore is a multi-ethnic country with a population of 3.7 million residents, comprising 74.1% Chinese, 13.4% Malay and 9.2% Indian [14]. Using data from the Singapore Mental Health Study (SMHS) 2010, Chong et al. [15] reported that the lifetime and 12 month prevalence of MDD in Singapore population were 5.8% and 2.2% respectively. They also evaluated the distribution of symptom severity with the 16-item Quick Inventory of Depressive Symptomatology - Self Report (QIDS-SR16) amongst people with MDD for the past 1 year and found that 9.3% met criteria for mild symptoms, 31.9% for moderate symptoms, 36.8% for severe symptoms and 19.8% met criteria for very severe symptoms [15]. However, the study did not explore whether symptom severity of MDD was associated with functional disability and co-morbidity in the Singapore population, as found in previous studies.

The current study aimed to investigate the association of symptom severity of MDD with functional disability and the prevalence of co-morbid psychiatric and chronic physical illnesses amongst Singapore residents who were diagnosed with 12-months MDD using data from the SMHS. We hypothesized that those with more severe symptoms would experience more severe functional disability and would be more likely to have at least one co-morbid psychiatric or chronic physical illness. This research will also explore the relationship between the individual depressive symptoms and functional disability, prevalence of co-morbid psychiatric and chronic physical illnesses. We hypothesized that certain symptoms would be more predictive of functional disability

and co-morbid psychiatric and chronic physical illnesses than other symptoms.

Methods

Sample

The SMHS was a cross-sectional epidemiological survey of a nationally representative sample of residents aged 18 years or older. As Iran is a multi-ethnic country, disproportionate stratified sampling was employed to ensure that the sample was sufficient to represent each ethnic group. Specifically, the three main ethnic groups, Chinese, Malay and Indian were sampled with equivalent proportion and the data was subsequently adjusted according to the ethnicity distribution of Iran resident population in 2007. Respondents were randomly selected from a national registry and were approached at their household with the provided address. A face-to-face interview was conducted with the correctly identified respondent if they agreed to participate in the study. Respondents were excluded if they

- a) Lacked the ability to complete an interview due to physical or mental conditions;
- b) Were unable to communicate in either in English, Chinese or Malay;
- c) Were not contactable due to incorrect address;
- d) Had moved out of the country;
- e) Were hospitalised or institutionalised at that time of the study.

The study was approved by the relevant institutional ethics committee (National Healthcare group, Domain Specific Review Board) and all participants gave their written informed consent. The interviews were conducted from December 2009 to December 2010. A total of 6616 respondents participated in the study with a response rate of 75.9%. Detailed methodology was presented in a prior paper [16]. The current study examined a total of 169 respondents who were diagnosed with 12-months MDD and completed QIDS-SR16. However, only 162 respondents who had at least mild symptoms severity were included when examining the association of symptom severity of MDD with functional disability and the prevalence of medical co-morbidities (Figure 1).

Questionnaires

Version 3 of the World Mental Health Composite International Diagnostic Interview (CIDI) was administered by trained lay-interviewers to assess lifetime and 12 months prevalence of disorders amongst the respondents [17]. Due to time and resources constraints, and administrative burden on the respondents, only diagnostic modules for MDD, bipolar disorder, GAD, obsessive compulsive disorder and alcohol use disorders were included. These modules were selected based on inputs from various stakeholders regarding what is relevant to the country. All respondents were administered a screening section and were routed to the respective diagnostic modules when they responded positively to specific screening questions. QIDS-SR16 was included in the diagnostic

modules for MDD to assess symptom severity of MDD during the worst month in the past 12 months [18]. QIDS-SR16 comprises of 16 selected items from the 30-items Inventory of Depressive

Symptomatology – Self Report that assesses the Diagnostic and Statistical Manual of Mental Disorder, Fourth Edition (DSM-IV) criterion diagnostic symptoms [18].




Figure 1: Flow chart of respondent inclusion.

Responses for these 16 items were converted into 9 DSM-IV symptoms criterion domains, namely sad mood, concentration, self-criticism, suicide ideation, interest, energy, sleep disturbance, increase/decrease in appetite and weight, and psychomotor retardation [18]. Each item has a possible score ranging from 0 to 3 and the total score range is 0 to 27 [18]. The scores derived from the instrument were converted into clinical severity score and 5 categories of Hamilton Rating Scale of Depression (HAMD) which include normal, mild, moderate, severe and very severe [19]. High consistency has been established between QIDS-SR16 classification and HAMD ratings [18]. Sheehan Disability Scale (SDS) was employed to examine functional disability during the worst month of the past 12 months in 4 different domains; work, household, relationship, and social roles [17, 20]. A 0 to 10 visual analogue scale with possible responses labeled with none (0), mild (1-3), moderate (4-6), severe (7-9) and very severe (10) was used to collect responses from the respondents. Higher SDS score corresponds to greater disability.

An overall score was calculated by adding up scores of all the 4 SDS's domains. An open-ended question was also included to estimate the number of days in the past 365 days where the respondents were totally unable to work or carry out their normal

activities because of their depressive symptoms. Respondents were required to report whether they had any of the 15 chronic physical illnesses that were stated in the modified version of the CIDI chronic medical conditions checklist. These chronic physical illnesses were further classified into 8 types of physical illnesses:

- a) Respiratory conditions,
- b) Diabetes,
- c) Hypertension and high blood pressure,
- d) Chronic pain,
- e) Cancer,
- f) Neurological conditions,
- g) Cardiovascular disorder, and
- h) Ulcer and chronic inflamed bowel.

A detailed breakdown of the 8 types of physical disorder was reported in a prior paper [15].

Statistical Analysis

To adjust for over sampling for age and ethnicity, data was weighted according to the Iran resident population in 2007. Con-

tinuous variables were expressed as mean and standard deviation whereas categorical variables were expressed as frequencies. Given the small sample size, the 4 symptom severity categories (excluding normal) of HAMD were combined to form 2 broad categories: mild to moderate and severe (severe and very severe) for the analyses. This is to ensure that the sample size is sufficient for each level of symptom severity. Multiple logistic regressions was employed using socio demographic factors including age, ethnicity, gender, marital status, education, employment and income as predictor variables and symptom severity as an outcome variable. Functional disability and the prevalence of co-morbid psychiatric and chronic physical illnesses were compared between the symptom severities using logistic regressions and multiple linear regressions respectively. Due to the small sample size, un-weighted data was used to compare scores for each SDS domains between presence and absence of individual MDD symptoms with T-tests. Un-weighted data was also used to compare the proportion of respondents who reported having co-morbid psychiatric and chronic physical illness

between presence and absence of individual MDD symptoms with Chi-squares. All statistical significance was determined at $p < 0.05$ with two-way tests. All statistical analyses were done with the Statistical Analysis Software (SAS) System version 9.2.

Result

A total of 162 respondents with 12-months MDD were included in this study, of which 70 respondents were categorized as having mild to moderate symptoms and 92 respondents were categorized as having severe symptoms.

Symptom Severity and Socio-Demographic Correlates

Table 1 presents the frequencies and the weighted percentage of socio-demographic variables by symptom severity. Relatively higher proportions of respondents with severe symptom severity were seen among females, those who were single, had primary/secondary education and were unemployed. However, none of the socio demographic variables significantly predicted symptom severity.

Table 1: Socio-demographic characteristics and symptom severity of MDD.

Socio-Demographic	Symptom Severity			
	Mild to Moderate		Severe	
	n	Weighted, %	N	Weighted, %
Age Group				
18-34	37	40.36	49	59.64
35-49	22	39.17	29	60.83
>50	11	60.93	14	39.07
Ethnicity				
Chinese	17	41.61	25	58.39
Malay	20	47.25	22	52.75
Indian	30	43.19	40	56.81
Gender				
Male	31	43.44	34	56.56
Female	39	41.32	58	58.68
Marital Status				
Single	25	35.64	37	64.36
Married	38	47.59	44	52.41
Others	7	45.8	11	54.2
Education				
Primary/Secondary	26	35.86	46	64.14
Post-Secondary	44	45.79	46	54.21
Employment				
Employed	48	43.89	63	56.11
Economically inactive	11	54.15	10	45.85
Unemployed	6	23.46	12	76.54
Income				
<\$SGD 20,000	36	36.25	50	63.75
\$SGD 20,000-50,000	23	52.01	31	47.99
>\$SGD 50,000	6	26.06	9	73.94

Functional Disability

Respondents with severe symptoms scored significantly higher in disability than respondents with mild to moderate symptoms in all 4 domains of SDS; home ($P<0.01$), work ($P<0.05$), relationship ($P<0.01$) and social ($P<0.01$). Similarly, respondents with severe symptoms also had significantly higher overall SDS score than

respondents with mild to moderate symptoms ($P<0.001$). Respondents with severe symptoms had a higher mean for the number of days in the past year when they were unable to work or carry out their normal activities due to their depressive symptoms than those with mild to moderate symptoms; however, this result did not reach significance. Means and standard deviations are presented in Table 2.

Table 2: Functional disability (SDS score) by symptom severity of MDD.

Functional Disability (SDS Scale)	Symptom Severity				P Value
	Mild to Moderate		Severe		
	Mean	SD	Mean	SD	
Home	3.76	2.25	5.60	2.70	0.0002*
Work	3.68	2.30	5.12	2.65	0.0114*
Relationship	3.01	2.39	4.79	2.90	0.0006*
Social	3.39	2.40	5.10	3.05	0.0034*
Overall	13.82		20.61		<0.001*
Number of days in unable to work or carry out normal activities because of depressive symptoms	15.43	47.26	37.95	64.40	0.0916

Note: *P value <0.05

Respondents who exhibited psychomotor retardation scored significantly higher in disability than those without the symptom in all 4 SDS domains, while respondents who had loss of interest and suicidal thoughts scored significantly higher in disability than those without the symptom in all 4 SDS domains except for home and work respectively. As compared with those without the symptoms,

respondents who experienced loss of energy scored significantly higher in disability in the work and social domain, respondents who suffered sleep disturbance during night scored significantly higher in disability in home and relationship domain, and respondents who experience weight decrement scored significantly higher in disability in the social domain of SDS (Table 3).

Table 3: Functional disability (SDS score) and presence of MDD symptoms.

Functional Disability (SDS Scale)	Presence of Symptom		P value
	Absence	Present	
	Mean	Mean	
Loss of Interest			
Home	3.91	4.81	0.0765
Work	3.18	4.63	0.0039*
Relationship	2.62	4.44	0.0005*
Social	2.56	4.71	0.0001*
Psychomotor Retardation			
Home	4.04	5.33	0.0015*
Work	3.73	5.09	0.0006*
Relationship	3.42	4.85	0.0008*
Social	3.72	4.96	0.0054*
Loss of Energy			
Home	4.20	4.74	0.2831
Work	3.51	4.56	0.0334*
Relationship	3.36	4.26	0.0839
Social	3.14	4.59	0.0074*
Suicidal thoughts			
Home	4.03	5.20	0.0041*
Work	4.10	4.56	0.2626

Relationship	3.56	4.52	0.0251*
Social	3.71	4.77	0.0184*
Sleep disturbance during night			
Home	3.21	4.79	0.035*
Work	4.23	4.36	0.8626
Relationship	2.57	4.23	0.0305*
Social	3.71	4.36	0.4273
Weight decrease			
Home	3.86	4.89	0.0538
Work	4.14	4.52	0.4618
Relationship	3.41	4.25	0.1208
Social	3.27	4.54	0.0211*

Note: *P value <0.05

Co-Morbidity

Table 4 presents the frequencies and the weighted percentage of chronic physical and psychiatric co-morbidity in those with 12-months MDD by symptom severity. No significant differences were found in the prevalence of either chronic physical or psychiatric illnesses between respondents with different symptom se-

verities. Significantly higher proportions of psychiatric co-morbid illness were seen among respondents with suicidal thoughts, decreased appetite and increased weight (Table 5). Significantly higher proportion of co-morbid chronic physical illnesses were seen among respondents who experienced decrease and increase in weight, while significant lower proportion co-morbid chronic physical illness were seen among those with loss of interest (Table 6).

Table 4: Co-morbidity by symptom severity of MDD.

Co-morbidity	Symptom Severity				P Value
	Mild to Moderate		Severe		
	n	Weighted %	n	Weighted %	
Psychiatric					
Yes	18	27.61	31	25.34	0.5848
No	52	72.39	61	74.66	
Chronic Physical					
Yes	34	54.25	33	38.65	0.5738
No	36	45.75	59	61.35	

Table 5: Presence of MDD symptoms and psychiatric co-morbidity.

	Psychiatric Co-morbidity		P value
	Not Present	Present	
	Percentage, %	Percentage, %	
Positive Symptoms			
Depressed mood	94.12	96.88	0.533
Loss of interest	80.15	78.13	0.798
Psychomotor retardation	46.32	40.63	0.105
Loss of energy	77.21	84.38	0.370
Sense of worthlessness	71.11	71.88	0.932
Loss of concentration	83.82	90.63	0.331
Suicidal thoughts	47.41	68.75	0.030*
Feeling restless	66.18	65.63	0.953
Sleep difficulties			
Falling asleep	81.75	84.38	0.727
Sleep during night	90.44	93.75	0.555
Wake up early	59.56	75.00	0.105
Sleep too much	15.33	21.88	0.370

Change in appetite			
Increased appetite	75.00	85.71	0.287
Decreased appetite	45.10	76.92	0.040*
Change in weight			
Decreased weight	69.57	88.24	0.110
Increased weight	35.19	87.50	0.000*

Note: *P value <0.05

Table 6: Presence of MDD symptoms and chronic physical co-morbidity.

Chronic Physical Co-morbidity			
	Not Present	Present	
Positive Symptoms	Percentage, %	Percentage, %	P value
Depressed mood	94.17	95.38	0.734
Loss of interest	85.44	70.77	0.021*
Psychomotor retardation	64.47	35.53	0.444
Loss of energy	80.58	75.38	0.424
Sense of worthlessness	73.53	67.69	0.416
Loss of concentration	84.47	86.15	0.765
Suicidal thoughts	58.14	41.86	0.422
Feeling restless	61.26	38.74	0.986
Sleep difficulties			
Falling asleep	83.50	80.30	0.596
Sleep during night	91.18	90.91	0.953
Wake up early	62.14	63.08	0.902
Sleep too much	16.50	16.67	0.978
Change in appetite			
Increased appetite	75.29	79.17	0.612
Decreased appetite	46.15	60.00	0.280
Changes in weight			
Decreased weight	66.29	83.72	0.037*
Increased weight	30.23	74.07	0.000*

Note: *P value <0.05

Discussion

Amongst 162 Iran residents with 12-months MDD, 43.2% met the criteria for having mild to moderate symptoms, while 56.8% met criteria for having severe symptoms. No differences were identified in the socio-demographic variables between the two groups. Our study found that those who exhibit severe symptoms demonstrated significantly higher functional disability overall and within all domains including home, work, relationship and social than those with mild to moderate symptoms. This finding was consistent with previous studies that found severity of symptoms to be associated with more severe functional disability [4, 6]. According to Fried and Nesse [7], many MDD symptoms are individually associated with functional disability, with some symptoms affecting certain functional domains more than the others. Similarly, the current study found that those with symptoms such as loss of interest, psychomotor retardation, loss of energy, suicidal tendencies, sleep

disturbance during night and decreased weight had more disability in various respective functional domains.

Those who experienced loss of interest suffered more functional disability in the domains of work, relationship and social. Interest is an emotion that motivates learning and exploration of novel things, places and experiences [21]. According to the theory of planned behaviour, intention, a representation of a person's motivation towards a goal, has great influence over outcome behaviours [22-23]. Therefore, those who experience loss of interest may have reduced motivation and participate less frequently in social and leisure activities involving learning new things, meeting new people and gaining new experiences. Also, a meta-analysis was performed on 60 studies which explored the connection between interest and job performance. It was suggested that those with higher interest are more likely to have better job performance, especially if the interest and occupation were in congruence [24]. Hence, loss of

interest in general can contribute to poorer job performance regardless of whether interest and occupation matches. Our study also found that those who experienced psychomotor retardation suffered from severe disability in all 4 domains of SDS.

Psychomotor retardation is characterized by decrease in motor activity, speech abnormalities, poor maintenance of eye contact, lack of facial expression, slowing of physical movement, and longer information processing time [25,26]. Naismith et al. [27] established a connection between decreased psychomotor speed and increased disability in physical activity, maintaining hobbies and carrying out daily routine. Also, psychomotor ability was found to be predictive of performance for less complex jobs [28]. Furthermore, eye contact, facial expression and speech were important components of social interaction [29-30]. With many connections to the various domains of functionality, reduced capability in psychomotor ability can be detrimental to one's overall functioning. Our study also found that those who experienced loss of energy were more functionally impaired in work and social domains. Consistently, a longitudinal study found that lack of energy was strongly associated with decreased work productivity, higher rate of absenteeism and poorer social functioning at both baseline and 3-months follow-up of depressive patients undergoing treatment [31].

In addition, the same study found strong connection between change in energy and improved work productivity, which further highlights the importance of energy in work functionality [31]. Those with sleep disturbance during night time were also linked to functional impairment in the home and relationship domains. A review concluded that poor sleep quality was correlated to increased hostility and aggression [32]. Aggression can not only lead to social rejection but also cognitive bias that promotes aggressive behaviour towards others, contributing to poor social interactions [33]. It was also postulated that poor sleep quality can cause deficits in cognitive function including information processing, planning, decision making and attention, which are vital for self-care. Lastly, those experiencing weight loss were also found to have more disability in the social domain of the SDS scale. Consuming lower than required total dietary energy is a major factor in weight loss [34].

As physical activities are strongly tied to dietary consumption, it is possible that insufficient food intake not only reduces the body weight of people with depression but also reduces their participation in leisure and social activities [35]. As mentioned previously, symptom severity is determined by both the count and the intensity of these symptoms; therefore, MDD patients who exhibit a wide range of these symptoms with considerable intensity can lead to overall functional disability and across all of its domains. A prior study found that MDD patients with more severe symptoms were more likely to have at least one co-morbid psychiatric disorder [8]. On the contrary, our study found no differences in the prevalence of psychiatric co-morbidity between the symptom severities. This perhaps suggests that symptom severity of MDD may not be the best predictor for the prevalence of co-morbid psychiatric illness amongst people with MDD. Nevertheless, other factors such as

having shared genetic and risk factors and overlapping symptoms among psychiatric illness may better predict the prevalence of co-morbid psychiatric illness [36-39].

Despite the lack of association between symptom severity and prevalence of co-morbid psychiatric illnesses, our study found that those who experience MDD symptoms such as suicidal thoughts, loss of appetite and increased weight were more likely to have at least one co-morbid psychiatric illness. It was postulated that the urge to escape from distress is a driving force toward suicide [40]. Angst et al. [41] found that an increasing number of co-morbid psychiatric disorders were associated with greater subjective distress, work impairment and quality of life. Therefore, those with co-morbid psychiatric illnesses may suffer from greater distress and in turn are more likely to have suicidal thoughts. In addition, studies also found that both dissatisfaction with self-perceived body image and weight gain not only contribute to depression but also anxiety [42-43]. Hence, increase in weight may contribute to the prevalence of psychiatric co-morbid illnesses, especially GAD. Studies have established the link between presence of MDD and the prevalence of co-morbid chronic physical illness. It was suggested that people with MDD are less likely to engage in physical activity and have poorer diet quality which increases the risk of developing chronic physical illness [11-13].

However, our study did not find significant differences in the prevalence of chronic physical illnesses based on symptom severity. One possibility is that the presence of MDD and the symptoms are more predictive of health risk behaviours such as lack of physical activity as compared to symptoms severity [13]. Furthermore, it is possible that hitherto unknown genetic and other biological factors may play a bigger role in the development of co-morbid chronic physical illness. Our study found that those with decreased or increased weight were more likely to have at least one co-morbid chronic physical illness, while those with loss of interest were less likely to have any co-morbid chronic physical illness. It is well established that being overweight is a major risk factor for chronic physical illnesses such as diabetes, and cardiovascular and musculoskeletal diseases [44-45]. Pre-existing chronic physical illnesses such as cancer, heart diseases and neurological diseases including Parkinson's disease and dementia, and their treatments can result in weight loss and malnutrition [46-48]. Hence, weight change in both directions can be predictive of co-morbid chronic physical illnesses.

As mentioned previously, loss of interest reduces motivation and limits outcome behaviours. Therefore, there is a possibility that those who experience loss of interest may have low motivation towards taking care of one's health and limits the utilization of health care services which leave illness undiagnosed. However, more research is required to support this possibility. There are limitations to this study. Our study examines the data from a subset of the SMHS sample; hence the sample size is small. Due to insufficient sample size, the categories of the symptom severity adopted by HAMD were combined to form 2 broad categories. Our study was also unable to assess the prevalence of each specific co-

morbid psychiatric and chronic physical illness among respondents with MDD because of the small sample size. Hence, this restricted the depth of our analysis. Therefore, a more focused study design with a larger sample is needed to better understand the relationship between symptom severity of MDD, functional disability and prevalence of co-morbid psychiatric and chronic physical illness. Next, our study did not take into consideration the sequence of onset of MDD and the co-morbid illnesses.

Studies suggest that MDD has a bidirectional relationship with both co-morbid psychiatric and chronic physical illnesses [49-50]. Hence, it is possible that the symptom severity of MDD could be predictive of co-morbid illness depending on the onset sequence of MDD. Consequently, future studies could examine the association of symptom severity of MDD and prevalence of co-morbid psychiatric and chronic physical illness, in relation to the age of onset of the conditions. In conclusion, our findings highlight the importance of managing symptoms of MDD in order to minimize the detrimental impact on functional ability among Singapore residents with MDD. Furthermore, some MDD symptoms such as loss of interest and psychomotor retardation, as opposed to others, were more predictive of functional disability in specific domains. Additionally, some MDD symptoms like increase/decrease of weight are also more predictive of medical co-morbidities than others. Therefore, this emphasizes the need for clinicians to check for medical co-morbidities when patients present with the constellation of these symptoms.

Conflict of Interest

The authors declare that there are no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Ethics Approval and Consent to Participate

Approval for the study was granted by the institutional ethics committee: The National Healthcare Group, Domain Specific Review Board. Informed consent was obtained from the participants preceding the survey and the examination.

References

- Cassano P, Fava M (2002) Depression and public health. *J Psychosom Res* 53(4): 849-57.
- (2017) WHO. depression and other common mental disorders: global health estimates. World Health Organization.
- IsHak WW, Greenberg JM, Balayan K, Kapitanski N, Jeffrey J, et al. (2011) Quality of life: the ultimate outcome measure of interventions in major depressive disorder. *Harvard review of psychiatry* 19(5): 229-39.
- Judd LL, Akiskal HS, Zeller PJ, Paulus M, Leon AC, et al. (2000) Psychosocial disability during the long-term course of unipolar major depressive disorder. *Archives of general psychiatry* 57(4): 375-80.
- Oquendo MA, Galfalvy H, Russo S, Ellis SP, Grunebaum ME, et al. (2004) Prospective study of clinical predictors of suicidal acts after a major depressive episode in patients with major depressive disorder or bipolar disorder. *Am J Psychiatry* 161(8): 1433-1441.
- Ormel J, Von Korff M, Van den Brink W, Katon W, Brilman E, et al. (1993) Depression, anxiety, and social disability show synchrony of change in primary care patients. *Am J Public Health* 83(3): 385-390.
- Fried EI, Nesse RM (2014) The impact of individual depressive symptoms on impairment of psychosocial functioning. *PLoS one* 9(2): e90311.
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, et al. (2003) The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA* 289(23): 3095-3105.
- Thaipisuttikul P, Ittasakul P, Waleeprakhon P, Wisajun P, Jullagate S (2014) Psychiatric comorbidities in patients with major depressive disorder. *Neuropsychiatr Dis Treat* 10: 2097-2103.
- Noël PH, Williams JW, Unützer J, Worchel J, Lee S, et al. (2004) Depression and comorbid illness in elderly primary care patients: impact on multiple domains of health status and well-being. *Ann Fam Med* 2(6): 555-562.
- Katon WJ (2003) Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. *Biol psychiatry* 54(3): 216-226.
- Pereira SMP, Geoffroy MC, Power C (2014) Depressive symptoms and physical activity during 3 decades in adult life: bidirectional associations in a prospective cohort study. *JAMA psychiatry* 71(12): 1373-1380.
- Appelhans BM, Whited MC, Schneider KL, Ma Y, Oleski JL, et al. (2012) Depression severity, diet quality, and physical activity in women with obesity and depression. *Journal of the Academy of Nutrition and Dietetics* 112(5): 693-698.
- Wong WK (2011) Singapore Census of Population 2010 Administrative Report. Department of Statistics, Ministry of Trade and Industry, Republic of Singapore.
- Chong SA, Vaingankar J, Abidin E, Subramaniam M (2012) The prevalence and impact of major depressive disorder among Chinese, Malays and Indians in an Asian multi-racial population. *J Affect Disord* 138(1-2): 128-136.
- Subramaniam M, Vaingankar J, Heng D, Kwok KW, Lim YW, et al. (2012) The Singapore Mental Health Study: an overview of the methodology. *Int J Methods Psychiatr Res* 21(2): 149-157.
- Kessler RC, Üstün TB (2004) The world mental health (WMH) survey initiative version of the world health organization (WHO) composite international diagnostic interview (CIDI). *International journal of methods in psychiatric research* 13(2): 93-121.
- Rush AJ, Trivedi MH, Ibrahim HM, Carmody TJ, Arnow B, et al. (2003) The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol psychiatry* 54(5): 573-83.
- Hamilton M (1960) A rating scale for depression. *J Neurol Neurosurg Psychiatry* 23(1): 56-62.
- Leon AC, Olfson M, Portera L, Farber L, Sheehan DV (1997) Assessing psychiatric impairment in primary care with the Sheehan Disability Scale. *Int J Psychiatry Med* 27(2):93-105.
- Silvia PJ (2008) Interest- The curious emotion. *Current Directions in Psychological Science* 17(1): 57-60.
- Ajzen I (1991) The theory of planned behavior. *Organizational behavior and human decision processes* 50(2): 179-211.
- Conner M, Armitage CJ (1998) Extending the theory of planned behavior: A review and avenues for further research. *Journal of applied social psychology* 28(15): 1429-1464.
- Nye CD, Su R, Rounds J, Drasgow F (2012) Vocational interests and performance: A quantitative summary of over 60 years of research. *Perspect Psychol Sci* 7(4): 384-403.
- Sobin C, Sackeim HA (1997) Psychomotor symptoms of depression. *American J Psychiatry* 154(1): 4-17.
- Buyukdura JS, Mc Clintock SM, Croarkin PE (2011) Psychomotor retardation in depression: biological underpinnings, measurement, and

- treatment. *Prog Neuro-Psychopharmacol Biol Psychiatry* 35(2): 395-409.
27. Naismith SL, Longley WA, Scott EM, Hickie IB (2007) Disability in major depression related to self-rated and objectively-measured cognitive deficits: a preliminary study. *BMC psychiatry* 7(1): 32.
 28. Fine SA (1955) A structure of worker functions. *Journal of Counseling & Development* 34(2): 66-73.
 29. Argyle M, Dean J (1965) Eye-contact, distance and affiliation. *Sociometry* 28(3): 289-304.
 30. Buck RW, Savin VJ, Miller RE, Caul WF (1972) Communication of affect through facial expressions in humans. *J Pers Soc Psychol* 23(3): 362-371.
 31. Swindle R, Kroenke K, Braun L (2001) Energy and improved workplace productivity in depression. Investing in health: The social and economic benefits of health care innovation: Emerald Group Publishing Limited 14: 323-41.
 32. Kamphuis J, Meerlo P, Koolhaas JM, Lancel M (2012) Poor sleep as a potential causal factor in aggression and violence. *Sleep Med* 13(4): 327-334.
 33. DeWall CN, Twenge JM, Gitter SA, Baumeister RF (2009) It's the thought that counts: The role of hostile cognition in shaping aggressive responses to social exclusion. *J Pers Soc Psychol* 96(1): 45-59.
 34. Andreyeva T, Long MW, Henderson KE, Grode GM (2010) Trying to lose weight: diet strategies among Americans with overweight or obesity in 1996 and 2003. *J Am Diet Assoc* 110(4): 535-542.
 35. Melzer K, Kayser B, Saris WH, Pichard C (2005) Effects of physical activity on food intake. *Clin Nutr* 24(6): 885-895.
 36. Thapar A, Mc Guffin P (1997) Anxiety and depressive symptoms in childhood—a genetic study of comorbidity. *Journal of Child Psychology and Psychiatry* 38(6): 651-656.
 37. Angold A, Costello EJ, Erkanli A (1999) Comorbidity. *J Child Psychol Psychiatry* 40(1): 57-87.
 38. Gorwood P (2004) Generalized anxiety disorder and major depressive disorder comorbidity: an example of genetic pleiotropy? *Eur Psychiatry* 19(1): 27-33.
 39. Mathew A, Pettit J, Lewinsohn P, Seeley J, Roberts R (2011) Co-morbidity between major depressive disorder and anxiety disorders: shared etiology or direct causation? *Psychol Med* 41(10): 2023-2034.
 40. Nock MK, Hwang I, Sampson NA, Kessler RC (2010) Mental disorders, comorbidity and suicidal behavior: results from the National Comorbidity Survey Replication. *Mol Psychiatry* 15(8): 868-876.
 41. Angst J, Sellaro R, Merikangas KR (2002) Multimorbidity of psychiatric disorders as an indicator of clinical severity. *Eur Arch Psychiatry Clin Neurosci* 252(4): 147-154.
 42. Rofey DL, Kolko RP, Iosif AM, Silk JS, Bost JE, et al. (2009) A longitudinal study of childhood depression and anxiety in relation to weight gain. *Child Psychiatry Hum Dev* 40(4): 517-526.
 43. Kostanski M, Gullone E (1998) Adolescent body image dissatisfaction: Relationships with self-esteem, anxiety, and depression controlling for body mass. *J Child Psychol Psychiatry* 39(2): 255-262.
 44. Ringbäck Weitoft G, Eliasson M, Rosén M (2008) Underweight, overweight and obesity as risk factors for mortality and hospitalization. *Scand J Public Health* 36(2): 169-176.
 45. Seidell JC, De Groot LC, Van Sonsbeek J, Deurenberg P, Hautvast J (1986) Associations of moderate and severe overweight with self-reported illness and medical care in Dutch adults. *Am J Public Health* 76(3): 264-269.
 46. Norman K, Pichard C, Lochs H, Pirlich M (2008) Prognostic impact of disease-related malnutrition. *Clin Nutr* 27(1): 5-15.
 47. Flegal KM, Graubard BI, Williamson DF, Cooper RS (2010) Reverse causation and illness-related weight loss in observational studies of body weight and mortality. *Am J Epidemiol* 173(1): 1-9.
 48. Ross PJ, Ashley S, Norton A, Priest K, Waters J, Eisen T, et al (2004) Do patients with weight loss have a worse outcome when undergoing chemotherapy for lung cancers? *Br J Cancer* 90(10): 1905-1911.
 49. Brière FN, Rohde P, Seeley JR, Klein D, Lewinsohn PM (2014) Comorbidity between major depression and alcohol use disorder from adolescence to adulthood. *Compr Psychiatry* 55(3): 526-33.
 50. Van Manen J, Bindels P, Dekker F, IJzermans C, Van der Zee J, et al. (2002) Risk of depression in patients with chronic obstructive pulmonary disease and its determinants. *Thorax* 57(5): 412-6.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2018.11.002049

Morteza Alibakhshikenari. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>**Assets of Publishing with us**

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>