

# Effect of Some Weaning Foods on Weanling Albino Rats

Umaru Hauwa Aduwamai<sup>1\*</sup> and Mahmud Bilkisu Abubakar<sup>2</sup>

<sup>1</sup>Department of Biochemistry, School of Life Sciences Modibbo Adama University of Technology Yola, Nigeria

<sup>2</sup>Department of Biochemistry, Faculty of Science, Adamawa State University Mubi, Nigeria

\*Corresponding author: Umaru Hauwa Aduwamai, Department of Biochemistry, School of Life Sciences Modibbo Adama University of Technology Yola, Nigeria



## ARTICLE INFO

**Received:** 📅 September 03, 2019

**Published:** 📅 September 10, 2019

**Citation:** Umaru Hauwa A, Mahmud Bilkisu A. Effect of Some Weaning Foods on Weanling Albino Rats. Biomed J Sci & Tech Res 21(2)-2019. BJSTR. MS.ID.003570.

**Keywords:** Malnutrition; Weaning; Groundnut; Soya Beans; Tigernut; Beniseed; Crayfish

## ABSTRACT

Inadequate intake of essential nutrients to replace breast milk during weaning is a major cause of malnutrition in young. The effect of some weaning foods on weanling albino rats was determined. Thirty weanling albino rats were randomly divided into six groups of five rats each. Sixty percent (60%) tiger nut, soya bean, groundnut, beniseed, and crayfish were incorporated into the normal diet and administered to the weanling albino rats for twenty-eight days. Results obtained showed significantly higher feed efficiency ratio in groups fed soya bean and groundnut compared to other groups. The least feed efficiency ratio was observed in group fed tiger nut. Body weight was found to be highest in groups fed soya beans and ground nut (67.47 and 63.83% increase respectively) compared to other groups. Haematological parameters revealed significantly higher increase in PCV ( $P < 0.05$ ) in groups fed soya bean and ground nut compared to other groups. Result obtained from this study indicates that soya beans and groundnut should be preferred in selecting weaning foods.

## Introduction

Malnutrition among children is a major public health concern in the developing countries. It affects all aspects of children's life, its effects are not limited to physical health but extend to mental, social, and spiritual well-being [1]. The world health organization estimates that malnutrition accounts for 54 percent of child mortality worldwide [2], while for children under the age of five years, childhood underweight accounts for 35.0% of all deaths worldwide [3].

Deficiency in energy balance and protein are characteristics of malnutrition, which may impair the proper functioning of various physiological processes [4]. Many factors can cause malnutrition, inadequate food intake, infections, psychosocial deprivation, and insanitary environment as well as lack of hygiene, social inequality, and possibly some genetic contribution. Reports from different studies report that children who live in households lacking access to sufficient clean healthy food are more likely to be predisposed to

poor nutrition and health related problems than children from food secure households [5]. Malnutrition increases the risk of infection and infectious disease, and moderate malnutrition weakens every part of the immune system [6]. It is a major risk factor in the onset of active tuberculosis [7].

Weaning is a major cause of malnutrition in the world. Weaning is influenced by socioeconomic status, cultural and religious beliefs and practices [8]. Adequate early nutrition is the most essential element in assuring adequate growth and appropriate cognitive and functional development of the infant. Severe protein energy malnutrition results in the clinical syndromes of marasmus, kwashiorkor or marasmic-kwashiorkor. Malnutrition arises due to poverty, ignorance, dietary practices, and agricultural productivity [9]. The aim of this study therefore is to determine the effect of some weaning foods on weanling albino rats with a view to finding the best weaning food.

## Materials and Methods

### Collection and Preparation of Samples

*Glycine max*, *Arachis hypogaea*, *cyperus esculentus*, *Sesamum indicum L.* and *Procambarus clarkii* were obtained from Mubi market, samples were washed, air dried and ground to powder.

### Experimental Animals

Thirty weanling male albino rats (aged 3 weeks) were obtained from National veterinary research institute Jos. The rats were housed under standard laboratory conditions.

### Methods

**Feed Efficiency Ratio (FER):** Feed efficiency ratio was calculated using the following formula:

$$\text{Feed efficiency ratio (FER)} = \frac{\text{Gain in body weight (g)}}{\text{Food intake (g)}}$$

### Body Weight Determination

Rats were weighed on weekly bases to ascertain weight gain or weight loss for a period of four weeks.

### Experimental Design

Thirty weanling albino rats were randomly divided into 6 groups of five rats each. Group 1 served as normal control, Group 2 served as tiger nut group, Group 3 served as soya beans group, Group 4 served as groundnuts group, Group 5 served as beniseed group, and Group 6 serves as crayfish group. About 60% of the experimental diet (tiger nut, soyabeans, groundnuts, beniseed, and crayfish) was added to 40% of the laboratory diet and fed to the animals for 4 weeks as shown in the Table 1.

**Table 1:** At Termination of experiment on the 28<sup>th</sup> day, the rats were sacrificed using chloroform vapor as an anesthesia and blood samples were collected by cardiac puncture into sample bottles.

Group	Description	Treatment
I	Diet 1	Normal feed (100%)
II	Diet 2	Normal feed (40%) + Tiger nut (60%)
III	Diet 3	Normal feed (40%) + soya bean (60%)
Iv	Diet 4	Normal feed(40%)+Ground nut (60%)
V	Diet 5	Normal feed (40%) + Beniseed (60%)
VI	Diet 6	Normal feed (40%) + cray fish (60%)

### Haematological Parameters

Haematological parameters were determined using automated analyzer. The Parameters include: PCV (Parked cells volume on weekly basis), Red blood cell count, Haemoglobin, Haematocrit, Mean Cell Volume (MCV), Mean Cell Haemoglobin (MCH), Mean Cell Haemoglobin Concentration (MCHC), Red Cell Distribution Width (RDW), White Blood Cell Count (WBC), Differential White Cell, Neutrophils Lymphocytes, Monocytes, Eosinophils, Basophils and Platelet count.

**Table 2:** Feed efficiency ratio of rats.

Groups	WEEK 1	WEEK 2	WEEK 3	WEEK 4	% Increase in feed efficiency ratio
1(Normal Control)	8.22 ±2.03 <sup>ad</sup>	15.00±3.05 <sup>ac</sup>	18.31±3.40 <sup>a</sup>	23.14±2.26 <sup>a,c</sup>	64.48
2 (Tiger nut)	7.80 ±1.53 <sup>ad</sup>	9.48 ± 2.00 <sup>ad</sup>	13.97 ± 1.01 <sup>b</sup>	15.71 ± 1.73 <sup>c</sup>	50.35 <sup>b</sup>
3 (Soya bean)	3.71±0.98 <sup>bd</sup>	5.37 ± 1.31 <sup>bc</sup>	10.77 ± 2.07 <sup>b</sup>	20.67 ± 2.11 <sup>c</sup>	82.04 <sup>a</sup>
4 (G/nut)	5.47±1.42 <sup>bd</sup>	9.22 ± 1.72 <sup>ad</sup>	14.15 ± 1.14	19.65 ± 3.90 <sup>c</sup>	72.16
5 (Beniseed)	4.22±0.65 <sup>bd</sup>	6.71 ± 1.19 <sup>bd</sup>	10.94 ± 2.41 <sup>b</sup>	14.65 ± 2.34 <sup>bc</sup>	71.19
6 (Crayfish)	5.08±1.50 <sup>bd</sup>	8.92±2.26 <sup>a</sup>	11.28±1.55 <sup>b</sup>	13.05 ± 2.54 <sup>bc</sup>	61.07

Note: Values are Mean ± SEM for 5 determinations

a = significantly higher (p<0.05) compared to other groups

b = significantly lower (p<0.05) compared to other groups

c = significantly higher (p<0.05) compared t other weeks

d = significantly lower (p<0.05) compared to other weeks

## Results

### Table 2

The feed efficiency ratio of rats fed with different supplement diets for four weeks is as presented in Table 2. The highest percentage of feed efficiency ratio (82.14%) was obtained in group fed with soya bean supplement diet, followed by group fed with groundnut (72.16%). Group fed with tiger nut had the least feed efficiency ratio (50.35%).

**Table 3**

Results of this table revealed a significant increase in body

weight (67.47%) of group three rats (Soya bean), followed by group fed with groundnut (63.83%) Tiger nut group had the least weight gain (41.90%) (Table 3).

**Table 3:** Body weight of rats in grams (g).

Groups	WEEK 1	WEEK 2	WEEK 3	WEEK 4	% Increase in weight
1 (Normal Control)	76.60 ± 8.01 <sup>a,d</sup>	97.00 ± 9.92 <sup>a</sup>	114.00 ± 11.42 <sup>a</sup>	155 ± 15.92 <sup>a,c</sup>	50.58
2 (Tiger nut)	61.00 ± 3.67 <sup>d</sup>	67.00 ± 4.63 <sup>b</sup>	73.20 ± 6.76	105.00 ± 5.11 <sup>c</sup>	41.90 <sup>b</sup>
3 (Soya bean)	42.00 ± 2.34 <sup>b,d</sup>	79.00 ± 4.63 <sup>a</sup>	91.60 ± 8.08 <sup>a</sup>	109.00 ± 11.51 <sup>c</sup>	67.47 <sup>a</sup>
4 (G/nut)	51.20 ± 2.58 <sup>d</sup>	81.00 ± 6.16 <sup>a</sup>	101.00 ± 7.1 <sup>a</sup>	141.00 ± 14.9 <sup>c</sup>	63.83 <sup>a</sup>
5 (B/seed)	51.20 ± 2.58 <sup>d</sup>	65.80 ± 3.89 <sup>b</sup>	85.40 ± 6.18	109.00 ± 13.22 <sup>c</sup>	53.03
6 (Crayfish)	46.40 ± 2.30 <sup>b,d</sup>	61.20 ± 4.14 <sup>b</sup>	74.00 ± 12.30 <sup>b</sup>	98.40 ± 17.38 <sup>b</sup>	52.85

Note: Values are Mean ± SEM for 5 determinations

a = significantly higher (p<0.05) compared to other groups

b = significantly lower (p<0.05) compared to other groups

c = significantly higher (p<0.05) compared to other weeks

d = significantly lower (p<0.05) compared to other weeks

**Table 4**

The Result of the packed Cell Volume (PCV) is as shown in Table 4. Results obtained showed significantly higher increase in PCV

levels in soya bean group (18.25%) and groundnut group (17.62%) compared to other groups. Tiger nut group had the least PCV result (11.02%).

**Table 4:** PCV of experimental rats in percentage (%).

GROUPS	WEEK 1	WEEK 2	WEEK 3	WEEK 4	% Increase in PCV
1 (Normal Control)	44.16 ± 1.40 <sup>b,d</sup>	46.33 ± 2.51 <sup>a</sup>	48.58 ± 7.07	50.91 ± 2.08 <sup>c</sup>	13.26
2 (Tiger nut)	49.83 ± 2.11 <sup>ad</sup>	51.00 ± 3.46 <sup>a</sup>	53.30 ± 2.12 <sup>a</sup>	56.00 ± 1.73 <sup>c</sup>	11.02 <sup>b</sup>
3 (Soya bean)	46.60 ± 5.78 <sup>d</sup>	48.92 ± 1.50 <sup>a</sup>	52.00 ± 3.46 <sup>a</sup>	57.00 ± 1.84 <sup>c</sup>	18.25 <sup>a</sup>
4 (Groundnut)	44.49 ± 1.44 <sup>b,d</sup>	46.66 ± 5.77 <sup>a</sup>	48.93 ± 1.41	54.00 ± 0.57 <sup>c</sup>	17.62 <sup>a</sup>
5 (Beniseed)	42.97 ± 1.18 <sup>b</sup>	44.66 ± 3.05 <sup>a</sup>	48.93 ± 9.89	51.03 ± 1.15 <sup>c</sup>	16.73
6 (Crayfish)	43.33 ± 2.01 <sup>b,d</sup>	45.33 ± 3.05 <sup>b</sup>	47.50 ± 3.53	49.78 ± 1.73 <sup>c</sup>	13

Note: Values are Mean ± SEM for 5 determinations

a = significantly higher (p<0.05) compared to other groups

b = significantly lower (p<0.05) compared to other groups

c = significantly higher (p<0.05) compared to other weeks

d = significantly lower (p<0.05) compared to other weeks

**Table 5**

The hematological parameters of rats fed with normal feed, tiger nut, soya bean, groundnut, beniseed and crayfish supplemented feeds is presented in Table 5. Hemoglobin, PCV, platelet, RBC, MCH, MCHC and MCV were studied. Hemoglobin levels were relatively higher in group three (18.66 ± 1.02) fed with soya bean supplement diet and group four (18.67 ± 0.74) fed with groundnut supplement diet when compared with the control (16.97 ± 0.84). Tiger nut

group had the least haemoglobin (15.54 ± 1.43). PCV level was significantly (p<0.05) higher in group three (57.00 ± 1.84) and four (54.00 ± 0.57) compared to control (50.91 ± 2.08). A slight increase in platelets was obtained for all groups when compared with the control group. Increase in RBC was recorded in group three (5.90 ± 0.10) and four (5.83 ± 0.41) but significantly lower in groups five (4.50 ± 0.60) and six (4.03 ± 0.05) fed with beniseed and crayfish supplement diet respectively.

**Table 5:** Haematological parameters of rat fed with different weaning foods.

Groups	Hb	PCV	WBC	RBC	MCH	MCHC	MCV	Platelets	Neutrophils	Eosinophil	Lymphocytes	Monocytes	Basophils
1 (Control)	16.97 ± 0.86 <sup>b</sup>	50.91 ± 2.08 <sup>b</sup>	5.43 ± 0.55 <sup>a</sup>	5.70 ± 0.17	17.00 ± 1.67 <sup>b</sup>	33.40 ± 1.57	51.80 ± 1.60 <sup>b</sup>	1.71 ± 0.02 <sup>b</sup>	50.66 ± 3.78 <sup>a</sup>	1.66 ± 0.02 <sup>b</sup>	44.33 ± 1.52	0.66 ± 0.01	0.33 ± 0.01
2 (Tiger nut)	15.54 ± 1.43 <sup>b</sup>	56.63 ± 0.57 <sup>a</sup>	5.03 ± 0.05	5.16 ± 0.00 <sup>b</sup>	18.80 ± 0.00 <sup>b</sup>	35.33 ± 1.53 <sup>a</sup>	51.93 ± 2.68 <sup>b</sup>	1.78 ± 0.04	49.33 ± 1.15 <sup>b</sup>	2.66 ± 0.05	44.66 ± 0.57	1.66 ± 0.02 <sup>b</sup>	0.33 ± 0.01

3 (Soya bean)	18.66± 1.02 <sup>a</sup>	57.00± 1.84 <sup>a</sup>	4.63± 0.46 <sup>b</sup>	5.90± 0.10 <sup>a</sup>	21.70± 1.13	32.60± 1.21	65.53± 1.10	1.80± 0.01 <sup>a</sup>	54.00± 3.60 <sup>a</sup>	2.33± 0.57	43.33± 4.16	0.02± 0.00 <sup>b</sup>	0.00± 0.00 <sup>b</sup>
4 (Groundnut)	18.67± 0.74 <sup>a</sup>	54.00± 0.57 <sup>a</sup>	5.13± 0.41	5.83± 0.41 <sup>a</sup>	19.53± 1.33	32.86± 0.92	59.60± 5.10	1.83± 0.01 <sup>a</sup>	58.66± 1.52 <sup>a</sup>	2.00± 0.00	37.33± 3.21 <sup>b</sup>	1.00± 0.00 <sup>b</sup>	0.00± 0.00 <sup>b</sup>
5 (Beniseeb)	17.01± 1.00 <sup>a</sup>	51.03± 1.15 <sup>b</sup>	5.00± 0.10	4.50± 0.60 <sup>b</sup>	24.33± 4.36 <sup>a</sup>	33.06± 0.60	75.26± 14.33 <sup>a</sup>	1.77± 0.02	48.66± 0.57 <sup>b</sup>	1.33± 0.57 <sup>b</sup>	47.66± 0.57 <sup>a</sup>	1.00± 0.00 <sup>b</sup>	1.00± 0.00 <sup>a</sup>
6 (Arayfish)	16.50± 0.46 <sup>b</sup>	49.78± 1.73 <sup>b</sup>	5.23± 0.11 <sup>a</sup>	4.03± 0.05 <sup>b</sup>	25.86± 1.85 <sup>a</sup>	33.40± 0.79	77.13± 4.81 <sup>a</sup>	1.79± 0.01	55.33± 3.05	3.33± 0.57 <sup>a</sup>	39.00± 5.56	2.66± 0.20 <sup>a</sup>	1.00± 0.00 <sup>a</sup>

Note: Values are Mean±SEM for 5 determinations

a = significantly higher, compared to other groups

b = significantly lower, compared to other groups

## Discussion

The high percentage increase in feed efficiency ratio observed in Table 2 for group fed with soya bean supplement diets could be as a result of the high protein content (39.44 g) of soya bean that is required for growth and development by the weanling rats. Soya bean also contains high amount of folic acid (204.16 g) that is required for brain development in weanling children. Increase body weight observed in group fed groundnut could be as a result of all the essential amino acid content in the diet. Peanuts contain all the essential amino acids necessary for normal body growth and metabolism [10]. From Table 3 above, the gain in weight of rats fed with soya bean and groundnuts supplement diets was significantly higher than the control and other groups fed with different supplement diets. The study reveals a significant increase in body mass of the rats studied. Increased in body weights is greatly associated with feed intake and constituent of feed consumed. This increased in body weight is greater with soya bean and groundnuts supplement diets than with the standard feed. This could result from the composition of diets, particularly its high proteins, (required to complement that of breast milk), carbohydrates, and fats contents which ensures an adequate intake favorable to the growth of the rats [11]. Soya bean and groundnuts are commonly mixed with other cereals for animal feeds, and is the main source of energy for the diet because of its high levels of starch and fatty acids [12]. Globally, soya beans and groundnuts are among the largest sources of dietary protein. This study agrees with the findings of Daniels et al. [13] who stated that among plant proteins consumed, those from soya beans and groundnuts are considered to have a high nutritional value. Decreased in groups fed tiger nut may be due to the fiber content of tiger nut which interferes with iron absorption. The high content of fiber content of tiger nut may have a good effect on digestion in adults but not in infants [14].

The low body weights of rats fed tiger nuts may also be due to protease inhibitors which affects protein digestibility. Protease inhibitors are globular proteins that bind irreversibly to protease enzymes, thereby inhibiting their proteolytic activities in the digestive tract of both humans and animals. The presence of protease inhibitors in foods decrease protein digestibility, pancreatic hypertrophy, growth inhibition and increase in demand for sulphur containing amino acids [15]. The packed cell volume

showed significant increased across all groups. The increased observed was highest for groups fed with soya bean and groundnut (56.00 ± 1.73 and 57.00 ± 1.84) after 4 weeks respectively when compared with the control. Increase in hematological parameters in the animals in this study could be explained by the presence of the various nutritive components contained in the diets. The increase in hematological parameters in some experimental and control groups could be associated with constituent of diets particularly soya bean and groundnut fed diets. Iron is an important co-factor for cytochrome oxidase enzymes at cellular level metabolisms and required for Red Blood Cell (RBC's) productions [16]. Magnesium plays a structural role in bone, cell membranes, and chromosomes [17].

From Table 5 above, the soya bean and groundnut fed group showed significant increase in most of the haematological parameters when compared with the control. There was no significant increase in WBC in all the groups when compared with the control. White blood cell (leucocytes) are known to increase sharply when infection occurs, as one of the first line of defense of the body. The increase in PCV, Haemoglobin, RBC, Platelets, neutrophils, counts following tiger nut, soya bean, groundnut, beniseed and crayfish supplements feeding for 4 weeks confirms the nutritional properties of these feeds.

## References

- (2016) World health organization The WHO child growth standards.
- Duggan C, Watkins JB, Walker WA (2008) Nutrition in pediatrics: Basic science, clinical application. BC Decker, Hamilton, pp. 127-141.
- Pruss-Ustun A, Bos R, Gore F, Bartram J (2008) Safer water, better health - Costs, benefits and sustainability of interventions to protect and promote health. World Health Organization (WHO), Geneva, Switzerland, p. 53.
- Bertin C (2006) Report of the standing committee on Nutrition at its 33<sup>rd</sup> session. World Health Organization, Geneva, Switzerland, p. 1-21.
- Bhargava M, Aggarwal PSD, Kandpal, SD, Semwal J (2015) Magnitude of undernutrition in urban and rural school going children using World health organization reference standards. Ntl J Comm Med 6(4): 452-457.
- Stillwaggon E (2008) Race, sex, and the neglected risks for women and girls in Sub-Saharan Africa. Feminist Eco 14(4): 67-86.
- Schaible UE, Kaufmann SHE (2007) Malnutrition and infection: Complex mechanisms and global impacts. PLoS Med 4(5): e115.

8. Amuna P, Zotor F, Chinyana YR (2000) The role of traditional cereals/legumes/fruit based multi mix in weaning in developing countries. *J Fd Sci Nutr* 30: 116-122.
9. Fotso JC, Kuate-Defo B (2005) Measuring socioeconomic status in health research in developing countries: Should we be focusing on households, communities or both?. *Soc Indic Res* 72(2): 189-237.
10. Huffman SL, Martin LH (2001) First feedings: Optimal feeding of infants and toddlers. *Nutr Res* 14: 127-159.
11. Dean J, Edwards DG (1985) The availability to the rat of energy from various diets ingredients. *Lab Ani* 19(4): 305-310.
12. Bouafou KG M (2007) Study of the production of maggots from household refuse and the nutritional value of dried maggot meal (FAS) in growing rats. PhD Thesis Animal Physiology, University of Cocody, Abidjan, Ivory Coast, pp. 145.
13. Daniels JL, Longnecker MP, Rowland AS, Golding J (2004) Fish intake during pregnancy and early cognitive development of offspring. *Epidemiol* 15(4): 394-402.
14. Sanchez-Zapata E, Fernandez-Lopez JA, Perez-Alvarez J (2012) Tiger Nut (*Cyperus esculentus*) commercialization: Health aspects, composition, properties, and food applications. *Comprehen Rev Fd Sci Fd Safe* 1(4): 366-377.
15. Adediran AM, Karim R, Oyeyinka SA, Oyeyinka AT, Awonorin SO (2013) Physico-chemical properties and akara making potentials of preprocessed Jack beans *Canavalia ensiformis* and Cowpea (*Vigna unguiculata L. Walp*) composite flour. *Croatian J Fd Tech Biotech Nutri* 8: 102.
16. Shafir T, Angulo-Barroso RS, (2009) Iron deficiency anemia in infancy and reach and grasp development. *Infant Behav Dev* 32: 366-375.
17. Maurice Edward Shils, Moshe Shike (2006) Modern nutrition in health and disease, (10<sup>th</sup> edn). Lippincott Williams & Wilkins, Baltimore, pp. 223- 247.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2019.21.003570

Umaru Hauwa Aduwamai. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>