

Cefoperazone with Antacid, Metal Complexation studied by Spectrophotometrically and Exploration of Antimicrobial Activity, *In-Vitro* Investigation

Joushan Ara*, Syeda Ridita Sharif, Farjana Nawrin, and Md Shahidul Islam

Department of pharmacy, University of Science and Technology Chittagong (USTC), Chattogram, Bangladesh

Corresponding author: Joushan Ara, Lecturer, Department of Pharmacy, University of Science & Technology Chittagong (USTC), Chattogram, Bangladesh



ARTICLE INFO

Received: 📅 October 09, 2020

Published: 📅 October 20, 2020

Citation: Joushan Ara, Syeda Ridita Sharif, Farjana Nawrin, Md Shahidul Islam. Cefoperazone with Antacid, Metal Complexation studied by Spectrophotometrically and Exploration of Antimicrobial Activity, *In-Vitro* Investigation. Biomed J Sci & Tech Res 31(2)-2020. BJSTR. MS.ID.005081.

Keywords: Cefoperazone; Spectrophotometry; Complexation; Job's Plot; Antimicrobial activity

ABSTRACT

Cefoperazone as a Cefoperazone sodium is a semisynthetic broad-spectrum cephalosporin antibiotic for which is used as a parenteral drug. Cefoperazone is quite effective against a wide variety of gram positive and gram-negative organisms including staphylococcus, Enterobacter, streptococcus, klebsiella and salmonella species. Cefoperazone is widely used in the treatment of Urinary tract infections, pelvic inflammatory diseases, respiratory tract infections etc. The principal purpose of present follow a line of investigation was to explore *in vitro* communications of Cefoperazone with crucial metal salt furthermore antacid to settle on the bioavailability in view of antimicrobial movement of Cefoperazone following drug metal communications at the pH7.4. Cefoperazone is one of the cephalosporin drugs to is exercised to treat numerous different sorts of infections originated by bacteria. Since the existence of compelling the ligand may influence bioavailability of metal in blood and tissues, for that reason with the intention of study the handy interaction of the Cefoperazone with vital elements in attendance in the body. Cefoperazone has been related with Zn, Mg as an *in-vitro* examination. Besides, the anti-microbial movement of drug along with the complexes was resolved. It has scrutinized that Cefoperazone interrelates with metal and antacid on the pH 7.4 by plotting various UV spectrophotometric methods. This examine authenticates that there was the interaction between Cefoperazone and metal, antacid which was authenticated by job's plot technique furthermore by antimicrobial exploration. This was authenticated that region of inhibition of Cefoperazone with Metal in addition to antacid condensed from 16mm to 14mm and 13mm correspondingly. The typical Cefoperazone disk also checked in opposition to *Staphylococcus aureus*.

Introduction

Cefoperazone sodium is a third-generation semisynthetic antibiotic that is used in the treatment of infections caused by susceptible microorganisms [1]. It is official in United States Pharmacopoeia, Chemically, cefoperazone sodium is 7-[R{2-(4-ethyl-2,3- dioxopiperazin-1-yl carboxamide)-2-(4-hydroxyphenyl) acetamide}-3-[1-methyl-1H-tetrazol-5-yl-thiomethyl]]- 3-cephem-4-carboxilate. Its molecular formula is $C_{25}H_{26}N_9NaO_8S_2$. Cefoperazone is wide spectrum and third generation of cephalosporin occupied in management of respiratory tract in addition to urinary infections [2]. Now drug interaction be capable of simply be distinguished as interaction

connecting a drug in addition to other substance with the intention of prevents drug commencing the stage as predictable. These connections may happen due to be short of knowledge regarding the main ingredients implicated in the related substances [3]. Two medicines are antagonistic at what time their interaction occurs a reduce in special effects of one or else both of medicines. The dissimilar responses of receptor to act of a medicine has affected in number of arrangements, which use expressions such as "partial agonist", "competitive agonist" etc. This is even expected that a lot of authors would mismanage any given arrangement [4] Commonly

iron complexes were used in convey of oxygen in human blood as well as tissues. A mature at rest takes 250ml of uncontaminated oxygen for each minute, these oxygens conceded by metal complex convey system recognized heame, alloying oxygen to go away the blood at what time it arrives the tissue [5].

Antibiotic and metal interaction and successfully found there after interaction result. Also, the antimicrobial activity of drug and the metals complexes were determined. This has been seen that the antibiotic interacts along with metal at the pH 7.4. The different essential metal intricate of the many drugs has been synthesized in addition to characterized by the techniques like NMR, UV, atomic absorption, FT-IR as well as elemental analysis. Spectroscopic, IR Spectroscopic, disk diffusion method, Biological assay studies of complexes [6]. On the other hand, in this interaction there uses metals complex which are interaction with drug into our body. And have been use their various type of bacteria for finding zone of inhibition. Some paper there uses methanol and ethanol for disk diffusion result clearly showing. Cefoperazone in a single way then it has a high bioavailability and proportionally increasing the absorption with concentration. But if we use its interaction with metals then its bioavailability and absorption decrease and less half-life [7]. By using there, a various of methods and study such as Antifungal study, Antibacterial study, Biological study with different type of spectroscopic analysis, Conductometric, IR spectrometric and Disk-diffusion methods its shows that's less pharmacological action [8]. So that after all discussion it can state that if it is taken antibiotic then it will be used the metals complex before two hours and after one hour.

Materials and Methods

Cefoperazone stock solution 250milliliter of 1×10^{-2} Molar was ready by softening 1.386gm of Cefoperazone solution in the 250 milliliter of the demineralized water (DMW) in a 250milliliter volumetric flask. The reserve solution was thinned to desired potency by the buffer solution [9] (Tables 1 & 2).

Table 1: List of chemicals and reagents.

Serial No	Name	Source
1	Cefoperazone	Gift samples from Square pharmaceuticals Ltd.
2	ZincSulphate(Metal)	Merck ltd, Mumbai, India
3	Magnesium Hydroxide(Antacid)	Merck ltd, Mumbai, India
4	Sodium di-hydrogenphosphate	USTC, Foys lake, Chittagong, dept of pharmacy
5	Disodium hydrogen phosphate	USTC, Foys lake, Chittagong, dept of pharmacy
6	PhosphateBuffer	USTC, Foys lake, Chittagong, dept of pharmacy

Table 2: List of instruments & equipment's.

Name	Model	Source
pHMeter	P ^H -211	Hanna,Romania
UVspectrophotometer	T80	PG instrument Ltd, England
Electronic Balance	AL-204	Mettlertoledo, Switzerland
Pipette		Fischer scientific,Germany

Preparation of Metal Solutions

For the grounding of 0.01 Molar metal solution like zinc sulfate heptahydrate (exactly 0.28754gm) was evaluated exactly in addition to initiated with the assist of the funnel in a 100ml volumetric flask, softened in DM water in addition to framework to the mark by the similar solvent. These prime solutions were additional diluted ten creases in the equivalent solvent along with the concluding solutions were 0.0001 M concentration.

Preparation of Antacid Solutions

For the preparation of 0.01M antacid solution, $Mg(OH)_2$ (0.0740gm) was evaluated accurately in addition to initiated with the facilitate of funnel in 100milliliter volumetric flask, softened in the demineralized water furthermore construct up to mark by the equal solvent. These primary solutions were further diluted ten folds in the same solvent and the final solutions were 0.0001 M concentration [10].

Grounding of the Buffer Solutions

To get ready buffer solution 1.76gram of the disodium hydrogen phosphate were softened in the demineralized water by 2.43 gram of the solution dihydrogen phosphate in addition to pH was altered to 7.4 as well as volume was completed to 1000 milliliter with the equivalent solution [11].

Grounding of the Typical Curve of the Cefoperazone

Cefoperazone reserve solution at the pH 7.4 in addition to concentration of the 1×10^{-5} Molar was inserted in unlike concentrations to the 10 test tubes, to have subsequent concentrations like 9×10^{-5} Molar, 8×10^{-5} Molar, 7×10^{-5} Molar, 6×10^{-5} Molar, 5×10^{-5} Molar, 4×10^{-5} Molar, 3×10^{-5} Molar, 2×10^{-5} Molar, 1×10^{-5} Molar. Now the solutions were suitably mixed. Then the absorbance values of solutions were established at 585nm through UV spectrometer [12].

Using Disc Diffusion Process

Solution of celebrated concentration ($3 \mu g/ml$) of trial samples are completed through dissolving measured quantity of the trials in calculated amount of solvents. Dried in addition to sterilized filter term paper discs (6mm as a diameter) are subsequently impregnated with recognized amounts of trial substances by using micropipette. Discs holding the test substances are positioned on the nutrient agar media homogeneously seeded by the test

organism, typical antibiotic discs along with blank discs were utilized as positive furthermore negative control. These plates now are kept at the low temperature like 4°C for the 24 hours to permit maximum diffusion. Throughout this time desiccated discs suck up water from surrounding medium furthermore the trial materials dissolve along with spread out of the trial disc. The diffusion happens according to physical law to controls diffusion of the molecules during agar gel. Consequently, there was a gradual alter of trial substances concentrations in medium adjoining the disc [13].

Results and Discussions

From the subsequent Table 3, this can monitor to the absorbance of the Cefoperazone enhances with the enhancing concentration in keeping with Beer Lambert's equation (Table 4). From the Table 5, it can observe that the absorbance of Cefoperazone is different when it interacts with Mg (OH)₂. From the Table 6, it can observe that interaction between drug and metal may lead to form complexes which have different light absorption capacity and spectrum pattern is altered. So, any alteration and spectrum behavior are regarded as a tool for primary interaction from the spectral studies.

Table 3: Standard curve of Cefoperazone.

Mx10 ⁻⁵	Absorbance
1	0.472
2	0.480
3	0.494
4	0.503
5	0.516
6	0.523
7	0.530
8	0.539
9	0.544

Table 4: Absorbance of Cefoperazone at different wavelength.

Wavelength	Absorbance
200	0.037
210	0.039
220	0.33
230	0.179
240	0.106
250	0.029
260	0.034
270	0.065
280	0.088
290	0.095
300	0.125
310	0.136
320	0.164

330	0.197
340	0.216

Table 5: Spectral analysis of Cefoperazone with Mg(OH)₂.

Wavelength/nm	Absorbance of Cefoperazone	Absorbance of Cefoperazone with Mg(OH) ₂
200	0.037	0.543
210	0.039	0.776
220	0.033	0.662
230	0.179	0.323
240	0.106	0.182
250	0.029	0.092
260	0.034	0.135
270	0.065	0.147
280	0.088	0.169
290	0.095	0.178
300	0.125	0.189
310	0.136	0.193
320	0.164	0.198
330	0.197	0.214
340	0.216	0.219

Table 6: Combined Spectral analysis of Cefoperazone with ZnSO₄·7H₂O.

Wavelength/nm	Absorbance of Cefoperazone	Absorbance of Cefoperazone with ZnSO ₄ ·7H ₂ O
200	0.037	0.342
210	0.039	0.675
220	0.033	0.561
230	0.179	0.222
240	0.106	0.081
250	0.029	0.091
260	0.034	0.104
270	0.065	0.116
280	0.088	0.130
290	0.095	0.147
300	0.125	0.159
310	0.136	0.163
320	0.164	0.170
330	0.197	0.184
340	0.216	0.203

Outcome of the Metals on Cefoperazone by Job's Technique of Incessant Variation:

The molar proportions of complexes of the Cefoperazone by metal salt were calculated approximately through Job's method. The examined absorbance rates were calculated in the pH 7.4 at diverse concentration like 1x10⁻⁵ to 9x10⁻⁵ Molar of Cefoperazone by metal salt. Then Job's plots at the pH were acquired through plotting absorbance variation against mole portion of drug (Table

7). From the above we can observe that Cefoperazone forms strong 1:1 complex with zinc sulfate heptahydrate which is indicated as inverted 'V' shaped curve. From the above we can observe that

Cefoperazone forms strong 1:1 complex with Mg (OH)₂ which is indicated as inverted 'v' shaped curve (Table 8).

Table 7: Values of job's plot of Cefoperazone with ZnSO₄.7H₂O.

Concentration of Cefoperazone Mx10 ⁻⁵	Absorbance of Cefoperazone A	Concentration of ZnSO ₄ .7H ₂ O Mx10 ⁻⁵	Absorbance of ZnSO ₄ .7H ₂ O B	Absorbance of mixture C	Absorbance difference D=(A+B)- C
1	0.472	9	0.187	0.110	0.549
2	0.480	8	0.186	0.109	0.557
3	0.494	7	0.187	0.109	0.572
4	0.503	6	0.188	0.112	0.579
5	0.516	5	0.189	0.111	0.594
6	0.523	4	0.192	0.128	0.587
7	0.530	3	0.181	0.125	0.586
8	0.539	2	0.179	0.136	0.582
9	0.544	1	0.177	0.150	0.571

Table 8: Values of Job plot of Cefoperazone and Mg(OH)₂.

Concentration of Cefoperazone Mx10 ⁻⁵	Absorbance of Cefoperazone A	Concentration of Mg(OH) ₂ Mx10 ⁻⁵	Absorbance of Mg(OH) ₂ B	Absorbance difference D=(A+B)-C	Absorbance difference D=(A+B)-C
1	0.472	9	0.090	0.114	0.448
2	0.480	8	0.086	0.112	0.454
3	0.494	7	0.093	0.118	0.469
4	0.503	6	0.096	0.118	0.481
5	0.516	5	0.103	0.120	0.499
6	0.523	4	0.097	0.132	0.488
7	0.530	3	0.092	0.136	0.486
8	0.539	2	0.085	0.147	0.477
9	0.544	1	0.083	0.162	0.465

Antimicrobial Reading

The antimicrobial effectiveness of the trial agents is calculated by their action to avoid the enlargement of microorganisms neighboring discs which provides clear precinct of inhibition. Later than incubation, the antimicrobial actions of the trial materials were concluded through measuring diameter of zones of the inhibition inside millimeter through a crystal-clear mm scale (Table 9). The trial samples were examined in opposition to Staphylococcus aureus. The typical Cefoperazone disk also checked in opposition to Staphylococcus aureus. The consequences of the antimicrobial action calculated in view of diameter of the zone of the inhibition in the mm were demonstrated in Table 10. Antimicrobial sensitivity testing of Cefoperazone against Staphylococcus aureus after interacting with ZnSO₄.7H₂O and Mg (OH)₂ solution, respectively. Now it was proved that zone of the inhibition of the Cefoperazone through Metal and antacid Zn, Mg reduced from 16mm to 14mm

& 13mm correspondingly due to the metal, antacid and drug interaction.

Table 9: Combined absorbance of drug with different metal & antacid.

Cefoperazone	Cefoperazone with ZnSO ₄ .7H ₂ O	Cefoperazone with Mg(OH) ₂
0.472	0.187	0.090
0.480	0.186	0.086
0.494	0.187	0.093
0.503	0.188	0.096
0.516	0.189	0.103
0.523	0.192	0.097
0.530	0.181	0.092
0.539	0.179	0.085
0.544	0.177	0.083

Table 10: Diameter of zone of inhibition.

Bacteria used	Standard disk (zone of inhibition/mm)	Sample disk (zone of inhibition)
Staphylococcus aureus	16 mm	Cefoperazone + ZnSO ₄ .7H ₂ O
		14mm
Staphylococcus aureus	16 mm	Cefoperazone with Mg(OH) ₂
		13mm

Conclusion

Though, there were some limitations in this research work like proper financial support from any side, active pure drug collection was not easy in Bangladesh. Eventually it has been overcome. The sophisticated spectrophotometric technique is easy, straight as well as worthwhile for the fortitude of drug. Commencing this spectral reading, this has been observed that Cefoperazone provides a spiky peak at the 585nm. At what time Zinc Sulfate furthermore antacid solution, Mg (OH)₂ combined with Cefoperazone 1:1 ratio and the strength of peak alters extremely and absorption features are changed due to the interaction although the position of complex do not change. Then antimicrobial viewing of a mediator is crucial to find out its spectrum in opposition to various natures of the pathogenic microorganisms. The Job's plot has provided the molar percentage of the complexes of Cefoperazone through Zinc Sulfate in addition to antacid solution like Mg (OH)₂. At the pH 7.4 Cefoperazone structures tough 1:1 complex through Zinc Sulfate in addition to antacid solution like Mg (OH)₂ designated as 'Λ' natured curves. These curves can indicate well-built kinetics of the complexation between Cefoperazone by Zinc Sulfate along with antacid solution like magnesium hydroxide. The trial samples were checked in opposition to *Staphylococcus aureus*. The typical Cefoperazone disk also checked in opposition to *Staphylococcus aureus*. It was examined that antimicrobial action of the Cefoperazone reduces when it structures complexes through ZnSO₄.7H₂O in addition to antacid solution like magnesium hydroxide. Thus, through antimicrobial examination it was proved that zone of the inhibition of the Cefoperazone through Metal and antacid Zn, Mg reduced from 16mm to 14mm & 13mm correspondingly.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2020.31.005081

Joushan Ara. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>

References

- Narasimhamurthy LN, Basaya PS, Pai, Sanjay PN (2009) A Stability-Indicating HPLC Method for determination of Cefoperazone. Eurasian Journal of Analytical Chemistry 4(1): 10-18.
- Parra A, Garcia-Villanova J, Rodenas V, Gomez MD (1994) First and second derivative spectrophotometric determination of cefoperazone and sulbactam in injections. Journal of Pharmaceutical and Biomedical Analysis 12(5): 653-657.
- Ratna Kumari K, Devala Rao G (2010) New Spectrophotometric Methods for the Determination of Cefoperazone in Pharmaceutical Formulation. International Journal of Pharmaceutical Research and Development 2(1): 1-5.
- Saleh GA, Askal HF, Radwan MF, Omar MA (2001) Use of charge-transfer complexation in the spectrophotometric analysis of certain cephalosporins. Talanta 54(6): 1205-1215.
- El-Azazy MS, Shalaby A, EL-Bolkiny MN, Khalil HM (2003) Spectrophotometric determination of cefepime hydrochloride, cefoperazone Na, ceftazidime pentahydrate, cefuroxime sodium and etamsylate using ammonium molybdate. Sci Pharm 71(3): 211-228.
- S Koppiset V, Chandra N (2011) Influence of Alcohol and Smoking on Drug Action: A Step for better utilization of drugs. Journal of Chemical and Pharmaceutical Research 3(1): 242-248.
- Chou TC (2006) Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. Pharmacological reviews 58(3): 621-681.
- C Sadowski D (2012) Drug Interactions with Antacids Mechanisms and Clinical Significance. Springer International Journal 11(6): 395-407.
- Tallarida RJ (2001) Drug synergism: its detection and applications. Journal of Pharmacology and Experimental Therapeutics 298(3): 865-872.
- Aronson JK (2004) Drug interactions-information, education, and the British National Formulary. British Journal of Clinical Pharmacology 57: 371-372.
- Sanket P, Satish P (2011) Dual wavelength spectrophotometric method for simultaneous estimation of ofloxacin and cefpodoxime proxetil in tablet dosage form. Asian J Pharm Life Sci 1(3): 261-268.
- Patil V, Chaudari RY (2012) Spectrophotometric method for estimation of cefpodoxime proxetil and ofloxacin in tablet dosage form by simultaneous equation method. Int J Pharm Life Sci 3(9): 1982-1984.
- Billow S, Kizek R, Jelen F, Novotn P (2003) Square-wave voltammetric determination of cefoperazone in bacterial culture, pharmaceutical drug, milk, and urine. Analytical and Bioanalytical Chemistry 377(2): 362-369.



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>