Sleeve Gastrectomy is the Most Common Cause of Gastro-Esophageal Reflux Disease in Comparison with Other Bariatric Operations

Ahmad Elnasasra², Ahmad Jaber², Hasan Kais² and Midhat Abu Sneineh^{1*}

¹Department of General Surgery, Bariatric Unit, AZ sint jan Medical Center, Belgium

²Department of Surgery A, Shamir (Assaf Harofeh) Medical Center, Israel

*Corresponding author: Midhat Abu Sneineh, Department of General Surgery, Bariatric Unit, AZ sint jan Medical Center, Belgium

ARTICLE INFO

Received: November 05, 2020

Published: November 09, 2020

Citation: Ahmad Jaber, Ahmad Elnasasra, Hasan Kais, Midhat Abu Sneineh. Sleeve Gastrectomy is the Most Common Cause of Gastro-Esophageal Reflux Disease in Comparison with Other Bariatric Operations. Biomed J Sci & Tech Res 31(5)-2020. BJSTR. MS.ID.005153.

Keywords: GERD; Bariatric operations; Gastrectomy; Gastric Reflux; Bariatric Surgery

Abbreviations: GERD: Gastro-Esophageal Reflux Disease; RNYGB: Roux En Y Gastric Bypass; SG: Sleeve Gastrectomy; LAGB: Laparoscopic Adjustable Gastric Band; IHD: Ischemic Heart Disease; IBD: Inflammatory Bowel Disease; COPD: Chronic Obstructive Pulmonary Disease

ABSTRACT

Purpose: GERD is one of the complications of bariatric operations that may affect quality of life. Our aim is to perform a retrospective cohort study to determine the incidence of symptomatic GERD following different types of bariatric surgery and which operations are considered a contraindication of GERD. In addition, we are attempting to identify the risk factors of GERD after bariatric surgery.

Materials and Methods: Medical records of 729 patients hospitalized for bariatric surgery between January 2010 and June 2019 in Shamir (Assaf Harofeh) Medical Center were reviewed.

Results: There was significant difference between the type of the bariatric procedure and the incidence of GERD symptoms after the operation. The incidence of symptomatic GERD in patients who underwent SG was 39.9% ($p = 0.0131$). This was significantly higher comparing to 16.4% following roux en y gastric bypass, 23.4% following LAGB, and 11% following OAGB. 113 patients out of 718 had a positive swallow test and of these patients 71 developed GERD symptoms post-operatively without correlation to the degree of reflux at the swallow test but with statistically significant correlation to the type of operation especially for SG (P value < 0.001) and to our knowledge this was never reported in the literature.

Conclusion: SG is a good bariatric procedure option but should be contraindicated in asymptomatic reflux contrast swallow study and symptomatic GERD patients preoperatively because of high levels of symptomatic GERD post-operatively. Asymptomatic reflux at contrast swallow study pre-operatively should be considered a risk factor for GERD after the operation.

Introduction

Bariatric operations are considered the best option for losing weight in morbidly obese patients, these operations include Sleeve Gastrectomy (SG), Roux En Y Gastric Bypass (RNYGB), One Anastomosis Gastric Bypass (OAGB) and Laparoscopic Adjustable Gastric Band (LAGB) [1,2], sleeve gastrectomy is one of the most common bariatric procedure worldwide, it is still considered a safe bariatric operation [3], epidemiological data shows that maintaining a normal BMI may reduce the likelihood of developing Gastro-Esophageal Reflux Disease (GERD) and its potential complications

[4], so bariatric operations if they decrease overall weight will lead to improvement in GERD symptoms but unfortunately that is not always the case. The causes of gastric reflux in general not fully understood but there are theories about abnormal relaxation of the lower esophageal sphincter, increased frequency of transient sphincter relaxation, or from increased pressure from the stomach secondary to a hiatus hernia or increased intra-abdominal pressure [5]. The literature suggests that different surgeries have different impacts on the symptomatology of GERD and that careful

assessment may be needed before performing bariatric surgery in patients with GERD [6]. GERD as a complication after bariatric surgery ranges between mild that responds well to lifestyle modifications, to intermediate that improves with daily treatment with proton pump inhibitor, to devastating complication that may affect quality of life and will need surgical treatment, thus choosing the appropriate operation according to GERD symptoms prior to the surgery is an important step.

El-Hadi et al reported that SG has been associated with an increased incidence of GERD following the procedure, however gastric banding and RNYGB has been shown to improve the symptoms of GERD in the short-term but but a new reflux symptoms and esophagitis in the long-term results [7]. Currently the data about GERD after bariatric surgery and the impact of bariatric operations on GERD is scarce so our aim is to perform a retrospective cohort study to determine the incidence of symptomatic GERD following different types of bariatric surgery and which operations are considered a contraindication of GERD. In addition, we are attempting to identify the risk factors of GERD after bariatric surgery.

Methods

Medical records of 729 patients hospitalized for elective bariatric surgery between January 2010 and June 2019 at Shamir (Assaf Harofeh) Medical Center were reviewed. Twenty-one patients were excluded because of missing data. The study was approved by the Institutional Review Board/Ethics Committee of the Shamir (Assaf Harofeh) Medical Center. Ages from 18 until 65 years and both sexes included in the study. We collected data from outpatient clinic records and telephone interview questionnaire. Available variables included demographic information, age, gender,

and primary and secondary diagnoses. Categorical variables were summarized with frequencies and percentages. Continuous variables were summarized using sample size, mean, standard deviation, median, minimum, maximum, and 95% CI. Correlations between categorical variables and incidence of GERD after surgery were tested for significance using the chi-square test. Correlations between continuous variables and incidence of GERD after surgery were tested for significance using the independent t-test. Baseline characteristics of interest included sex, age, diagnosis of GERD, and Body Mass Index (BMI) group.

Results

We retrospectively evaluated 718 patients with an average follow-up of 24 months (range 12–36 months), we collected the information from outpatient clinic as well as telephone interview using (GERD-Q questionnaire) [8] as an assessment for GERD symptoms (Table 1). GerdQ is a self-administered 6-item questionnaire that was recently developed as a tool to improve and standardize symptom-based diagnosis and evaluation of treatment response in patients with GERD [9]. According to outpatient clinic records, 236 (32.9%) patients have GERD symptoms (heartburn, regurgitation, dysphagia, odynophagia, increased salivation and chest pain) according to medical history before the operation (Table 2). Mean age was 48 ± 19 years (76% females). There were 218 laparoscopic SG (30.4%), 175 laparoscopic LAGB (24.4%), 61 RYGB (8.5%), and 264 (36.8%), OAGB procedures during the study period (Table 2). At the follow-up period, 96 of 718 patients (13.4%) developed new symptomatic GERD and 71 of 236 pre-operative GERD patients (30%) continued to have the same or worse GERD symptoms after the surgery (Table 3) so a total of 167 after the surgery for the period of the follow-up (Table 4).

Table 1: GerdQ self-assessment questionnaire.

S. No	Symptoms in the previous week	Symptoms presence			
		0 days	1 day	2-3 days	4-7 days
1.	How often did you have a burning feeling behind your breastbone (heartburn)?	0	1	2	3
2.	How often did you have stomach contents (liquid or food) moving upwards to your throat or mouth (regurgitation)?	0	1	2	3
3.	How often did you have a pain in the center of the upper stomach?	3	2	1	0
4.	How often did you have nausea?	3	2	1	0
5.	How often did you have difficulty getting a good night's sleep because of your heartburn and/or regurgitation?	0	1	2	3
6.	How often did you take additional medication for your heartburn and/or regurgitation other than what the physician told you to take (such as Maalox)?	0	1	2	3

Table 2: *Chi-square p value is <0.001., **chi-square p value is 0.0341.

Type of operation	Number/percentage of patients	Number of symptomatic GERD patients before the operation	Number of patients persist to have GERD symptoms after the operations
Sleeve Gastrectomy	218(30.4%)	42	39 (92.9%)*
Roux en-Y gastric bypass	61(8.5%)	41	7 (17.1%)
One anastomosis gastric bypass	264(36.8%)	98	30 (30.6%)

LAGB	175(24.4%)	56	20 (35.7%)
Total	718	236	96 (40.7%)**

Table 3: *chi-square p value is < 0.001, **Chi-square p value is 0.0341.

Type of operation	Number of patients with asymptomatic reflux at contrast swallow study	Number of these patients with New onset GERD symptoms after the operation
Sleeve Gastrectomy	23	19 (82.6%)*
Roux en-Y gastric bypass	33	18 (54.4%)
One anastomosis gastric bypass	28	15 (53.6%)
LAGB	29	19 (65.5%)
Total	113	71 (62.8%)**

Table 4: *chi-square p value is =0.0131.

Type of operation	Number of patients	Symptomatic GERD after the operation total
Sleeve Gastrectomy	218(30.4%)	87 (39.9%)*
Roux en-Y gastric bypass	61(8.5%)	10 (16.4%)
One anastomosis gastric bypass	264((36.8%)	29 (11%)
LAGB	175(24.4%)	41 (23.4%)
Total	718	167 (23.3)

Our protocol that all patients should pass contrast swallow study before the surgery [10,11], we examined which patients had a pre-operative asymptomatic reflux at contrast swallow study and a total of 113 patients out of 718 (15.7%) had a positive swallow test (mild, moderate or severe) and of these patient 71 (62.8%) developed GERD symptoms post-operatively without correlation to the degree of reflux at the swallow test but with statistically significant correlation to the type of operation especially for SG (P value <0.001) (Table 3) and to our knowledge this was never reported in the literature. There was a significant difference between the type of the bariatric procedure and the incidence of GERD symptoms after the operation. The incidence of symptomatic GERD in patients who underwent SG was 39.9% (p =0.0131). This was significantly higher comparing to 16.4% following roux en y gastric bypass, 23.4% following LAGB, and 11% following OAGB (Table 4). There was no correlation between the incidence of GERD and; BMI at surgery; change in BMI; co-morbidities such as diabetes, hyperlipidemia, hypertension, and COPD (Table 5), we also examined which of the symptomatic per-operative GERD patients persist to have symptoms and we found that 92.9% of the patients that underwent SG with diagnosis of GERD pre-operatively persist to have symptoms after the operation (Table 2).

Table 5: Medical history and GERD after operation.

		Total	GERD after surgery
Smoker	Yes	231	58 (25.1%)
	No	487	109 (22.4%)
Dyslipidemia	Yes	156	42 (26.9%)
	No	562	125 (22.2%)
Diabetes mellitus	Yes	161	34 (21.1%)
	No	557	111 (19.9%)

Hypertension	Yes	193	46 (23.8%)
	No	525	121 (23%)
IHD	Yes	21	7 (33.3%)
	No	697	160 (23%)
IBD	Yes	4	1 (25%)
	No	714	166 (23.2%)
COPD	Yes	31	6 (19.4%)
	No	687	161 (23.4%)
Cirrhosis	Yes	5	1 (20%)
	No	713	166 (23.3%)
Renal Failure	Yes	2	0 (0%)
	No	716	167 (23.3%)

Discussion

We found that 23.3% of our patients persist to have or developed symptomatic GERD symptoms after the bariatric operation. We found a significant difference between the type of the bariatric procedure and the risk of GERD with the highest prevalence after LSG (39.9%) compared to 11% after OAGB, 23.4% after LAGB, and 16.4% after RNYGB, respectively. Of these patients 159 patients responds well to PPI and 8 patients needed further investigation with contrast swallow and gastroscopy and of them 2 patients operated for hiatal hernia repair, one after sleeve gastrectomy and the other after OAGB. Our findings is consistent with some previous studies as Howard et al [12], Frezza et al [13], however our finding that smoking is not related to symptomatic GERD is inconsistent with Kohata et al which said smoking cessation reduced the prevalence of GERD and the severity of reflux [14], but in our study only bariatric patients examined which is not the same population in other studies about relation between smoking and GERD. We think

according to our results that every patient should be questioned for GERD symptoms before the operation. Sleeve gastrectomy should be contraindicated in bariatric patients with symptoms of GERD or in asymptomatic patients with reflux on contrast swallow study. Moreover, although currently major guidelines make no definite statement about sleeve gastrectomy in patients with asymptomatic reflux [15], our findings are not correlated with Petersen WV et al that states SG significantly increased lower esophageal pressure and may protect obese patients from gastro-esophageal reflux [16], however they are consistent with Stenard F, et al [15].

None of the parameters in our study apart from the type of the bariatric surgery, symptoms of GERD before the operation and reflux at the contrast study were found to be of value in predicting the development of GERD postoperatively. The limitations of our study include its single-center focus, the relatively small number of SG patients, and the retrospective study design.

Conclusion

Sleeve gastrectomy is a good bariatric procedure option but should be contraindicated in asymptomatic reflux contrast swallow study and symptomatic GERD patients preoperatively because of high levels of symptomatic GERD post-operatively. Asymptomatic reflux at contrast swallow study pre-operatively should be considered a risk factor for GERD after the operation.

References

1. Lemaitre F, Léger P, Nedelcu M, Nocca D (2016) Laparoscopic sleeve gastrectomy in the South Pacific. Retrospective evaluation of 510 patients in a single institution. *Int J Surg* 30: 1-6.
2. Iannelli A, Dainese R, Piche T, Facchiano E, Gugenheim J (2008) Laparoscopic sleeve gastrectomy for morbid obesity. *World J Gastroenterol* 14(6): 821-827.
3. Sneineh MA, Harel L, Elnasasra A, Razin H, Rotmensh A, et al. (2020) Increased Incidence of Symptomatic Cholelithiasis After Bariatric Roux-En-Y Gastric Bypass and Previous Bariatric Surgery: a Single Center Experience. *Obes Surg* 30(3): 846-850.
4. Mustafa El Hadi, Daniel W Birch, Richdeep S Gill, Shahzeer Karmali (2014) The effect of bariatric surgery on gastroesophageal reflux disease. *Can J Surg* 57(2): 139-144.
5. Orlando RC (2001) Overview of the mechanisms of gastroesophageal reflux. *Am J Med* 111 Suppl 8A: 174S-177S.
6. Suter M, Dorta G, Giusti V, Calmes JM (2004) Gastro-esophageal reflux and esophageal motility disorders in morbidly obese patients. *Obes Surg* 14(7): 959-966.
7. El Hadi M, Birch DW, Gill RS, Karmali S (2014) The effect of bariatric surgery on gastroesophageal reflux disease. *Can J Surg* 57(2): 139-144.
8. Suzuki H, Matsuzaki J, Okada S, Hirata K, Fukuhara S, et al. (2013) Validation of the GerdQ questionnaire for the management of gastro-oesophageal reflux disease in Japan. *United European Gastroenterology Journal* 1(3): 175-183.
9. Jones R, Junghard O, Dent J, N Vakil, K Halling, et al. (2009) Development of the GerdQ, a tool for the diagnosis and management of gastro-oesophageal reflux disease in primary care. *Aliment Pharmacol Ther* 30(10): 1030-1038.
10. Katz PO, Gerson LB, Vela MF (2013) Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol* 108(3): 308-328.
11. Lemanowicz A, Serafin Z (2014) Imaging of patients treated with bariatric surgery. *Pol J Radiol* 79: 12-19.
12. Howard DD, Caban AM, Cendan JC, Ben David K (2011) Gastroesophageal reflux after sleeve gastrectomy in morbidly obese patients. *Surg Obes Relat Dis* 7(6): 709-713.
13. Frezza EE, Ikramuddin S, Gourash W, T Rakitt, A Kingston, et al. (2002) Symptomatic improvement in gastroesophageal reflux disease (GERD) following laparoscopic Roux-en-Y gastric bypass. *Surg Endosc* 16(7): 1027-1031.
14. Kohata Y, Fujiwara Y, Watanabe T, Masanori Kobayashi, Yasuhiko Takemoto et al. (2016) Long-Term Benefits of Smoking Cessation on Gastroesophageal Reflux Disease and Health-Related Quality of Life. *PLoS One* 11(3): e0150554.
15. Stenard F, Iannelli A (2015) Laparoscopic sleeve gastrectomy and gastroesophageal reflux. *World J Gastroenterol* 21(36): 10348-10357.
16. Petersen WV, Meile T, Küper MA, Zdichavsky M, Königsrainer A, et al. (2012) Functional importance of laparoscopic sleeve gastrectomy for the lower esophageal sphincter in patients with morbid obesity. *Obes Surg* 22(3): 360-366.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2020.31.005153

Midhat Abu Sneineh. Biomed J Sci & Tech Res

This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>