

Davis Buckley Hyper Immunoglobulin E Syndrome Associated with IgA deficiency: The second Case

Aamir Jalal Al Mosawi*

Advisor doctor and expert trainer, Baghdad Medical City and the National Center of Training and Development, Head, Iraq
Headquarter of Copernicus Scientists International Panel, Baghdad, Iraq



*Corresponding author: Aamir Jalal Al Mosawi, Advisor doctor and expert trainer, Baghdad Medical City and the National Center of Training and Development, Head, Iraq Headquarter of Copernicus Scientists International Panel, Baghdad, Iraq

ARTICLE INFO

Received: 📅 November 10, 2020

Published: 📅 November 19, 2020

Citation: Aamir Jalal Al Mosawi. Davis Buckley Hyper Immunoglobulin E Syndrome Associated with IgA deficiency: The second Case. Biomed J Sci & Tech Res 32(1)-2020. BJSTR. MS.ID.005185.

Keywords: Davis Buckley Syndrome; Hyper Immunoglobulin E (IgE); Recurrent Infection Syndromes; IgA Deficiency

ABSTRACT

Background: Davis Buckley hyper immunoglobulin E (IgE) recurrent infection syndromes are very rare group of heterogeneous disorders of variable genetic basis and manifestations, but they are essentially characterized by significant elevation of serum IgE levels occurring in association with recurrent infections with or without coarse facial features and eosinophilia. The aim of this paper is to report the occurrence of an extraordinary case of Davis Buckley syndrome associated with associated with IgA deficiency.

Patients and Methods: An Iraqi boy of Turkish origins was referred at the age of 10 years to the pediatric psychiatry clinic at the Children Teaching Hospital of Baghdad Medical City because of poor school performance and poor attention span was studied.

Results: The boy didn't have coarse facial features or facial dysmorphism. During the previous few years the boy was experiencing urinary tract infections and recurrent ear infections associated with discharge and were being treated with several courses of antibiotics. The boy didn't have growth retardation or bony or dental abnormalities. The parents were health relative, and family history was negative for similar condition. Blood tests showed eosinophilia and reactive thrombocytosis. Serum IgE was significantly elevated at the level of 0.4mg/dL (normal 0.01-0.04 mg/dL). The patient also had very low IgA level which was 18 mg/dL (normal 90-450 mg/dL). Mutation analysis was positive for Stat 3 mutation.

Conclusion: There has been only one case of Davis Buckley syndrome associated with low IgA levels reported by Mazzone et al (1996). The case in this paper is most probably the second case of the syndrome associated with IgA deficiency.

Short Communication

Davis Buckley hyper immunoglobulin E (IgE) recurrent infection syndromes are very rare group of heterogeneous disorders of variable genetic basis and manifestations, but they are essentially characterized by significant elevation of serum IgE levels occurring in association with recurrent infections with or without coarse facial features and eosinophilia. The syndrome may also be associated with eczema, skeletal and dental abnormalities [1-4]. In 1966, Davis et al described two red-haired, fair-skinned girls who had infection sino-pulmonary, staphylococcal cutaneous infections, Elevated levels of IgE and defective of neutrophil chemo taxis. Davis et al thought that the condition could be similar to the affliction of

Job, a biblical prophet whose body was covered with sore boils [1]. Buckley [2] described two boys had severe dermatitis, recurrent cutaneous, pulmonary, joint abscesses, growth retardation, and coarse facial features. The boys also had with significant elevation serum IgE levels and eosinophilia [2]. The aim of this paper is to report the occurrence of an extraordinary case of Davis Buckley syndrome associated with associated with IgA deficiency.

Patients and Methods

An Iraqi boy of Turkish origins was referred at the age of 10 years to the pediatric psychiatry clinic at the Children Teaching Hospital

of Baghdad Medical City because of poor school performance and poor attention span was studied.

Results



Figure 1: The boy didn't have coarse facial features or facial dysmorphism.

Table 1.

ESR	22 mm/hour
Hemoglobin	12.6 g/dL
Hematocrit (HCT)	35.1%
WBC	7.99x10 ⁹ /L
Neutrophil	2.07 (36.3%)
Lymphocyte	3.95 (52.4%)
Monocyte	.471 (6.24%)
Eosinophil	.344 (4.55%)
Basophile	.111 (1.47%)
RBC	4.58 x10 ¹² /L
Mean corpuscular volume (MCV)	76.5 FL
Mean corpuscular hemoglobin (MCH)	28.2 pg
Mean corpuscular hemoglobin concentration (MCHC)	30.0 g/dL
Red blood cell distribution width (RDW)	13.1%
Platelet count	619 x 10 ⁹ /L
Mean platelet volume (MPV)	4.99 FL

The boy didn't have coarse facial features or facial dysmorphism (Figure 1). During the previous few years the boy was experiencing urinary tract infections and recurrent ear infections associated with discharge and were being treated with several courses of

antibiotics. The boy didn't have growth retardation or bony or dental abnormalities. The parents were health relative, and family history was negative for similar condition. Blood tests showed eosinophilia and reactive thrombocytosis (Table 1). Serum IgE was significantly elevated at the level of 0.4mg/dL (normal 0.01-0.04 mg/dL). IgG level was 1065 mg/dL (normal 800-1800 mg/dL), and IgM level was 144 mg/dL (normal 60-180 mg/dL). The patient also had very low IgA level which was 18 mg/dL (normal 90-450 mg/dL). Mutation analysis was positive for Stat 3 mutation.

Discussion

Szczawinska Poplonyk [4] emphasized the complexity of the syndrome, the diversity of clinical manifestations, and the heterogeneous genetic nature. They suggested the possibility of the existence of an autosomal dominant classical form of the syndrome, and an autosomal recessive form. The autosomal dominant hyper-IgE syndrome is associated with facial, dental, skeletal, and connective tissue abnormalities which are not reported in association with the recessive type [4]. During childhood, platelet counts range between 250 x 10⁹/L and x 10⁹/L. An elevated platelet count more than 500, 000/10⁹/L indicates the presence of thrombocytosis. Primary thrombocytosis is an extremely rare during childhood, while secondary or reactive thrombocytosis is commonly a reactive process caused by infection, chronic inflammation, and iron deficiency [5]. The dominant form Davis Buckley syndrome has generally been attributed to STAT-3 mutation with loss of function of the signal transducer and activator of transcription 3, while autosomal recessive forms are generally attributed to mutations in dedicator of cytokines 8 (DOCK-8) [6].

In this case, the occurrence of Davis Buckley syndrome without coarse facial features, but with eosinophilia and the presence of Stat 3 mutation suggest an overlap between the autosomal dominant and recessive forms. Mazzone [7] emphasized the rarity IgA deficiency in association with Davis Buckley syndrome. They reported a young woman with Davis Buckley syndrome who had recurrent urinary tract infections, genital bacterial infections, generalized erythematous eczematous patches and stomatitis of oral mucosa and fever. The patient had high immunoglobulin IgE and low IgA levels [7].

Conclusion

There has been only one case of Davis Buckley syndrome associated with low IgA levels reported by Mazzone [7]. The case in this paper is most probably the second case of the syndrome associated with IgA deficiency.

Acknowledgement

The author would like to acknowledge his for the parents for willing accepting publishing the photo of the child.

References

1. Davis SD, Schaller J, Wedgwood RJ (1966) Job's Syndrome. Recurrent, "cold", staphylococcal abscesses. *Lancet* 1(7445): 1013-1015.
2. Buckley RH, Wray BB, Belmaker EZ (1972) Extreme hyperimmunoglobulinemia E and undue susceptibility to infection. *Pediatrics* 49(1): 59-70.
3. Hill HR, Quie PG (1974) Raised serum-IgE levels and defective neutrophil chemotaxis in three children with eczema and recurrent bacterial infections. *Lancet* 1(7850): 183-187.
4. Szczawinska Poplonyk A, Kycler Z, Pietrucha B, Heropolitanska Pliszka E, Breborowicz A, et al. (2011) The hyperimmunoglobulin E syndrome-clinical manifestation diversity in primary immune deficiency. *Orphanet J Rare Dis* 6: 76.
5. Chiarello P, Magnolia M, Rubino M, Liguori SA, Miniero R (2011) Thrombocytosis in children. *Minerva Pediatr* 63(6): 507-513.
6. Alcántara Montiel JC, Vega Torres BI (2016) Síndrome hiper-IgE. Lecciones de la función y defectos de STAT-3 o DOCK-8 *Rev Alerg Mex* 63(4): 385-396.
7. Mazzone A, Girola S, Fossati G, Mazzucchelli I, Ricevuti G (1996) Síndrome di Giobbe (Iper-IgE) ed ipo-IgA. Rara associazione di immunodeficienze. *Recenti Prog Med* 87(2): 71-74.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2020.32.005186

Aamir Jalal Al Mosawi. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>