

# Isolation and Characterization of Detergent Compatible Alkaline Protease Producing *Bacillus Subtilis* APO-1

Ami Vishnusinh Sayaniya and Pritesh Patel\*

CG Bhakta Institute of Biotechnology, Uka Tarsadia University, India

\*Corresponding author: Pritesh Patel, Assistant Professor, C. G. Bhakta Institute of Biotechnology, Uka Tarsadia University, Bardoli, Maliba Campus, Surat, Gujarat, India



## ARTICLE INFO

Received: 📅 May 03, 2021

Published: 📅 May 12, 2021

**Citation:** Ami Vishnusinh Sayaniya, Pritesh Patel. Isolation and Characterization of Detergent Compatible Alkaline Protease Producing *Bacillus Subtilis* APO-1. Biomed J Sci & Tech Res 35(5)-2021. BJSTR. MS.ID.005750.

**Keywords:** Alkaline Protease; Bacteria; Biomedical Waste; *Bacillus Subtilis*; Enzyme

## ABSTRACT

Alkaline Proteases catalyze the hydrolysis of peptide bond between proteins have main applications in detergent industry. Therefore, production of alkaline protease with useful attributes of biotechnological interest from efficient microbial strain is significant. In present study, an extracellular alkaline protease enzyme producing bacteria *Bacillus Subtilis* APO-1 was isolated from soil of industrial area near Surat. The casein hydrolysis agar medium was used for isolation, cultivation and maintenance of alkaline protease producing strain. Bacteria with 22 mm clear zone of hydrolysis around the colony were selected for its further morphological, biochemical, and molecular level. The culture dependent approach based on PCR amplified partial subunit 16S rDNA sequence analysis was submitted to NCBI to obtain accession number MN093333. Molecular phylogenetic analysis suggested that the isolated bacterium is in homology with *Bacillus Subtilis*. Alkaline protease production was evaluated in a broad pH range 7.0–11.5 and found that organism was capable of producing enzyme at all pH range studied. Alkaline protease isolated from *Bacillus Subtilis* APO-1, recorded highest enzyme activity of 48.27U/ml at 37 °C. Further, it was found that among the various conditions of washing tried, the combination of alkaline protease of APO-1 and detergent was able to significantly remove blood stain from white cotton cloth. Our research suggests that the *Bacillus Subtilis* APO-1 is a promising strain for alkaline protease production for its possible commercial use as a constituent of detergent formulations.

## Introduction

Proteases are a complex group omnipresent enzyme in nature which catalyze the cleavage of the peptide bonds among the amino acid residues in different proteins; has a place with a class of hydrolases with massive physiological and commercial importance. In light of their acid-base behavior, proteases are characterized in to three groups, that is, acid, neutral and alkaline proteases. Acid proteases perform best at pH range of 2.0-5.0 and are mostly produced by fungi. Proteases with pH optima in the range of 7.0 or around are called neutral proteases. Neutral proteases are predominantly produced by plants. While proteases optimum activity at pH range of 8 or more are classified as alkaline proteases [1]. Though categorized into three classes on the bases of mode of action, alkaline proteases alone comprise about 89% of the entire

protease market because of its broad applications in detergents, leather processing, protein recovery, meat tenderization, food industry, pharmaceuticals, and bioremediation [2]. Alkaline proteases, with good activity and stability in high alkaline range, are interesting for several bioengineering and biotechnological applications. However, their predominant application is in the detergent industry [3]. Since the pH of laundry detergents is in the range of 9.0–12.0, alkaline proteases are utilized in detergent formulations, with other hydrolytic enzymes, as additives to work with the breakdown and release of proteins [4,5].

Many microorganisms produce proteases; however, the enormous extent of commercial alkaline proteases is generally gotten from *Bacillus* sp., as a result of its capacity to secrete large

amount of alkaline proteases having significant proteolytic activity, stability at high temperature and wide range of pH. Number of studies has been reported on alkaline protease production by *Bacillus* like *Bacillus cereus* [6], *Bacillus infantis* [7], *Bacillus subtilis* [8], *Bacillus pumilus* [9] etc. Microbial proteases, especially from *Bacillus subtilis* are the most widely exploited industrial enzymes [10]. Advances in analytical methods have exhibited that proteases direct profoundly explicit specific and selective modifications of proteins, for example, activation of zymogenic types of enzymes by limited proteolysis, blood coagulating and lysis of fibrin clusters, and processing and transport of secretory proteins across the membranes [11]. Removal of proteinaceous stains like blood, milk, egg, grass and chocolate is very difficult utilizing traditional detergent approach. Vishalakshi et al. reported the successful removal of blood stain by alkaline protease from *Streptomyces gulbargensis* [12]. Notwithstanding, removal of such stain is accomplished by utilizing alkaline proteases [13,14]. The alkaline protease enzymes are acquiring a massive significance in the detergent industries because of their inherent capacity to sustain the broad pH with effective specificity in various cleanser detergent additives. The latest trend is to utilize detergent-compatible proteases that are steady over a wide pH range. This enzyme also has inherent benefits and cost-effectiveness over the ordinary synthetic catalysts. The finding for detergent-compatible proteases with better properties is a continuous exercise. The present study was initiated to with the aim to collect soil sample from dump side near Surat for the isolation of alkaline protease producing alkaliphilic bacteria and qualitative and quantitative analysis of protease production. Further as an application part work was intended for blood stain removal and detergent compatibility of crude enzyme.

## Material and Methods

### Isolation of Alkaline Protease Producing Bacteria

Soil sample were collected from Surat dump side area below 5-6 cm depth and stored in sterile plastic bags and taken to the lab aseptically. Isolation of alkaline protease producing alkaliphilic bacteria was carried out using alkaline solid agar medium contained glucose (5 g/L), yeast extract (3 g/L), peptone (5 g/L), and agar (20 g/L), (pH 7 to 10.5) in addition to 20% (w/v) casein powder, as an indicator of alkaline protease production [15]. 1 gm of soil sample was weighted and serial dilution from 10<sup>-1</sup> to 10<sup>-6</sup> was carried out. 0.1 ml of each aliquot was spread on above mentioned media plate and incubated at 37 °C for 24 to 48 hours. The zone of hydrolysis was noted on all the pH range evaluated and the colony showing highest zone of inhibition was selected for further study. The colony was grown on casein agar plate repeatedly and preserved on Nutrient agar slant at 4°C for further study.

### Morphological and Biochemical Characterization

The morphological characterization of the isolated bacterium was performed through Gram staining. The different biochemical tests were performed to determining the metabolic activities of bacteria are indole production test, methyl red test, Voges - Proskauer test, urea hydrolysis test, hydrogen sulphide production test, nitrate reduction test, ammonia production test, citrate utilization test, phenylalanine deamination test, triple sugar iron agar test, starch hydrolysis test, lipid hydrolysis test, casein hydrolysis test, hemolysin production test, oxidase test, catalase test, gelatin hydrolysis test, dehydrogenase test, sugar utilization test.

### Molecular Identification and Phylogenetic Analysis

The genomic DNA was isolated according to Sambrook [16]. Total genomic DNA was extracted were used for PCR amplification of 16S rDNA in combination with the following sets of universal primers 1) 8F AGA GTT TGA TCM TGG CTC AG and 2) 1492R CGG TTA CCT TGT TAC GAC TT). DNA sequencing of PCR amplicon was carried out with 8F & 1492R primers using BDT v3.1 Cycle sequencing kit on ABI 3730xl Genetic Analyzer. The PCR amplified products were then subjected to 16s rRNA sequencing for molecular identification. Analyses of 16s rDNA gene sequence was performed with bioinformatics tool BLAST. Sequences were aligned against the database of National Centre for Biotechnology Information ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). The sequences were aligned using multiple sequence alignment program CLUSTAL W followed by MEGA X for phylogenetic analysis. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) were shown next to the branches and phylogenetic tree generated with Neighbor-joining method.

### pH Stability

Solid agar medium similar as isolation media was used to evaluate the ability of *Bacillus Subtilis* APO-1 to produce alkaline protease at different pH value i.e 7, 8, 9, 10, 11 and 11.5. 24-hour active culture was centrally inoculated on solid agar medium and plates were incubated at 37 °C. Alkaline protease production was observed after 48 hours.

### Protease Activity Assay

200µl of 24hr old bacterial culture was inoculated in 60ml casein hydrolysate medium and kept in incubator at 37°C. After 48-hour incubation, the bacterial culture was centrifuged at 3000 rpm for 12 min and the supernatant was utilized for alkaline protease enzyme assay. Protease action was dictated by addition of 0.65% casein in 50mM phosphate buffer (pH 7.5) as substrate. 1ml enzyme and casein substrate was incubated at 37°C for 20 min. To stop the reaction 2.5ml Trichloroacetic acid was utilized.

One unit of protease activity was characterized as the 0.1 unit increase in optical density upon 1 hr incubation. At that point solution was filtered by utilizing a 0.45  $\mu\text{m}$  filter and 2ml of filtrate was mixed with 2.5ml of 500mM sodium carbonate and incubated for 20 min at 37°C. After that, 0.5ml of Folin Ciocalteu's reagent was added, and the absorbance was measured at 660 nm utilizing Spectrophotometer [17]. The measure of protease produced by bacteria was assessed and expressed in microgram of tyrosine released under standard test conditions.

### Stain Removal and Detergent Compatibility of Bacterial Enzyme

Activity of protease enzyme as a detergent additive was analyzed through different combinations of treatment on blood sample-stained cloth. Four white cotton clothes (5×5cm) with blood separately treated in flasks as followings.

- Blood-stained cloth treated with (2ml) distilled water.
- Blood-stained cloth treated with (1ml) distilled water+ 1ml detergent (0.01mg/ml).
- Blood-stained cloth treated with (2ml) enzyme solution.
- Blood-stained cloth treated with (1ml) detergent (0.01mg/ml) + (2ml) enzyme solution.

All four flasks were incubated at 37°C for 15 min. All four flasks were incubated at 37°C for 15 min. After incubation, cloth pieces were taken out, washed with normal water and dried. Visual assessment of cloth pieces displayed the impact of enzymes in removal of blood stains (Figure 2). Untreated cloth pieces stained with blood were taken as control.

## Result and Discussion

### Isolation of Alkaline Protease Producing Bacteria




**Figure 1:** Zone of casien solubilization by *Bacilli subtilis* APO-1 isolated from industrial soil sample collected from Surat.

In the present study, an attempt was made to isolate most efficient extracellular alkaline protease producing bacteria from soil sample from industrial field present on outskirts of Surat region Gujarat. Based on its Casein hydrolysate activity through zone of solubilization, one potential isolate APO-1 was selected for further study (Figure 1). Bacterial colony appeared on the casein agar plates were observed to be 2-3 mm in diameter, lobate, flat and white and exhibited around 22 mm clear zone of hydrolysis around the colony. The isolate APO-1 was stored in 40% glycerol at 4°C for further experimental studies.

### Morphological, Biochemical and Molecular Characterization

The results of morphological characterization reveal that *Bacillus Subtilis* APO-1 was gram-positive, facultative anaerobe, motile and short rod. An extensive biochemical characterization of APO-1 was carried out and is tabulated in Table 1. Genomic DNA of the isolate APO-1 was amplified and obtained the product length was 1050 bp. The PCR amplification using forward primer and reverse primer generated a single band amplicon size of 1 kb on 1 % agarose gel (Figure 2).



**Figure 2:** PCR amplification of 16S rDNA gene of the isolated bacterium.

### Morphological, Biochemical and Molecular Characterization

The results of morphological characterization reveal that *Bacillus Subtilis* APO-1 was gram-positive, facultative anaerobe, motile and short rod. An extensive biochemical characterization of APO-1 was carried out and is tabulated in Table 1. Genomic DNA of the isolate APO-1 was amplified and obtained the product length


was 1050 bp. The PCR amplification using forward primer and reverse primer generated a single band amplicon size of 1 kb on 1 % agarose gel (Figure 2).

Neighbor-joining phylogenetic tree showing the relatedness between the 16S rRNA gene sequence of the isolated bacterium and their closest relative sequences retrieved from microbial databases (Figure 3). The phylogenetic tree constructed by the neighbor-joining method indicated that the isolate presented

on separate cluster. The numbers in parentheses correspond to the accession number. The optimal tree with the sum of branch length = 0.23477701 is shown. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The analysis involved 11 nucleotide sequences. That is, fewer than 5% alignment gaps, missing data, and ambiguous bases were allowed at any position. All positions with less than 95% site coverage were eliminated. Evolutionary analyses were conducted in MEGA X.

**Table 1:** Biochemical Characterization of *Bacilli subtilis* APO-1.

Biochemical Test	Result	Biochemical Test	Result
Glucose	Positive	Maltose	Negative
Xylose	Positive	Sucrose	Positive
Lactose	Negative	Fructose	Positive
Indole production	Positive	Triple sugar iron test	Negative
MR test	Positive	Starch hydrolysis test	Positive
VP test	Positive	Lipid hydrolysis test	Positive
Urea hydrolysis test	Negative	Casein hydrolysis test	Positive
H2S Production test	Negative	Hemolysin test	Negative
Nitrate reduction test	Positive	Oxidase test	Positive
Ammonia production test	Negative	Catalase test	Positive
Citrate utilization test	Positive	Gelatine hydrolysis test	Positive
Phenyl alanine test	Negative	Dehydrogenase test	Positive



**Figure 3:** Phylogenetic tree generated with Neighbour-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) were shown next to the branches and phylogenetic tree.

## pH Stability

pH stability is an essential characteristic for the use of alkaline protease as detergent additives. In our study protease of APO-1 was found active in broad pH range 7.0-11.5. Organism APO-1 grown well on all the pH studied with clear zone of casein solubilization.

## Protease Activity Assay

When the protease is testing digests casein, the amino acid tyrosine is liberated along with other amino acids and peptide fragments. The amount of protease produced was estimated and expressed in microgram of tyrosine released under standard assay conditions. In this assay, casein acts as a substrate. Folin's reagent primarily reacts with free tyrosine to produce a blue-colored chromophore which is quantifiable and measured as an absorbance value on the spectrophotometer. The more tyrosine that is released from casein, the more the chromophores are generated and the stronger the activity of the protease. From the standard curve the activity of protease samples can be determined in terms of Units, which is the amount in micromoles of tyrosine equivalents released from casein per minute. Micromoles of tyrosine divided by time in minutes yields measurement of protease activity called "units". The result showed in the micromoles of tyrosine liberated during this particular proteolytic reaction. Alkaline protease activity estimated in culture filtrate of APO-1 with 0.5, 0.7 and 1 ml of enzyme solution


was 32.11 U/ml, 40.20 U/ml, and 48.27 U/ml, respectively (Table 2).

**Table 2:** Extracellular enzyme production by *Bacilli subtilis* APO-1.

Isolate	Various amount of enzyme solution (ml)	Conc. ( $\mu\text{g/ml}$ )	Enzyme Unit/ml
Bacilli subtilis APO-1	1	183.5	48.27
	0.7	175.9	40.2
	0.5	160.9	32.11

## Stain Removal and Detergent Compatibility of Crude Enzyme

Cleansing property of protease was checked for the removal of blood stain from the white cotton cloth by mixing the enzyme with detergent solution. Additionally, correlation was established with the commercially available detergents, and results show that introduction of enzyme to the detergent increases the washing performance of the detergent. It was clearly illustrated the cleaning action of APO-1 protease, blood stain was not removed completely with detergent alone, while the mixture of enzyme with commercial detergent removes the blood stain more efficiently. Reflectance measurement showed that about 85-90 % blood stain was removed (Figure 4).



**Figure 4:** Visual examination of cloth pieces exhibited the effect of enzymes in removal of blood stains.

## Discussion

In this study, the microorganism *Bacillus Subtilis* APO-1 isolated from the soil is evaluated for the alkaline protease production. The protease from this organism is active at alkaline pH so this is alkaline protease. Alkaline proteases are a physiologically and commercially important group of enzymes used primarily as detergent additives. They play a specific catalytic role in the hydrolysis of protein. The capacity of *Bacillus* strains to produce large amounts of the alkaline

protease with high activity recognized them as significant sources of commercial proteases and in this way are excellent candidates for large scale proteases production. Numerous scientists revealed the production of proteases from *Bacillus* species and their applications in detergent industries [18]. El-Safey and Abdul-Raouf worked on production, purification and characterization of extracellular protease enzyme isolated from *B. Subtilis* and studied the optimum conditions for maximum protease activity which is

comparable to our study [19]. In our study the strain APO-1 has been identified as *Bacillus Subtilis* on the basis of morphological and biochemical and molecular characterization. Casein was used as the substrate in nutrient agar medium for qualitative analysis and showed maximum zone of activity after overnight incubation. Protease of APO-1 was found active in broad pH range 7.0-11.5. These findings are in accordance with earlier reports showing the optimum pH 10.0-10.5 reported by Banik et al. [20]. The similar results were reported by Genckal and Tari who found novel alkaline protease from alkalophilic *Bacillus* sp L21 showed optimal stability at pH 11.0 [21].

The protease form *Bacillus Subtilis* APO-1 gave better result in removal of blood stain from cotton cloth as detergent additives. Numerous investigations exhibited the applicability and suitability alkaline protease in commercial detergents, which served an amazing execution to speed up the cleaning capacity of detergents. Shaikh et al (2018) reported the detergent-compatible alkaline protease from the bacterial strain *Bacillus* sp. APP-07 isolated from soil [22]. Haddar et al. also reported detergent-compatible protease enzyme from *Bacillus* *mojavensis* A21, which exhibited excellent stability in the presence of solid and liquid detergents [3]. Mala et al. also reported biocompatibility of alkaline protease from *Bacillus* sps. with commercial detergents such as Kite, Tide and Aerial and reported that the enzyme showed 70% activity even after 3 hours [23].

## Conclusion

This is first time we are reporting the strain of *Bacillus Subtilis* capable of producing alkaline protease at pH11.5. This feature is promising for further detail characterization for its large-scale production. The alkaline protease form *Bacillus Subtilis* APO-1 presented excellent stain removal potential in combination with detergent solution. The blood stain was completely removed from the cotton fabric and might serve several possible applications as detergent additives.

## Acknowledgment

None.

## Conflict of Interest

None.

## References

- Gupta R, Beg Q, Khan S, Chauhan B (2002) An overview on fermentation, downstream processing and properties of microbial alkaline proteases. *Applied microbiology and biotechnology* 60(4): 381-395.
- Mukherjee AK, Adhikari H, Rai SK (2008) Production of alkaline protease by a thermophilic *Bacillus Subtilis* under solid-state fermentation (SSF) condition using *Imperata cylindrica* grass and potato peel as low-cost medium: characterization and application of enzyme in detergent formulation. *Biochemical Engineering Journal* 39(2): 353-361.
- Haddar A, Agrebi R, Bougateg A, Hmidet N, Sellami-Kamoun A, Nasri M (2009) Two detergent stable alkaline serine-proteases from *Bacillus mojavensis* A21: purification, characterization and potential application as a laundry detergent additive. *Bioresource Technology* 100(13): 3366-3373.
- Jain D, Pancha I, Mishra SK, Shrivastav A, Mishra S (2012) Purification and characterization of haloalkaline thermoactive, solvent stable and SDS-induced protease from *Bacillus* sp.: a potential additive for laundry detergents. *Bioresource Technology* 115: 228-236.
- Joshi S, Satyanarayana T (2013) Characteristics and applications of a recombinant alkaline serine protease from a novel bacterium *Bacillus lehensis*. *Bioresource technology* 131: 76-85.
- Lakshmi B, Kumar DM, Hemalatha K (2018) Purification and characterization of alkaline protease with novel properties from *Bacillus cereus* strain S8. *Journal of Genetic Engineering and Biotechnology* 16(2): 295-304.
- Saggu SK, Mishra PC (2017) Characterization of thermostable alkaline proteases from *Bacillus infantis* SKS1 isolated from garden soil. *PLoS one* 12(11): e0188724.
- Shafique T, Shafique J, Zahid S, Kazi M, Alnemer O, et al. (2021) Screening, selection and development of *Bacillus Subtilis* apr-IBL04 for hyper production of macromolecule alkaline protease. *Saudi Journal of Biological Sciences* 28(2): 1494-1501.
- Özçelik B, Aytar P, Gedikli S, Yardımcı E, Çalışkan F, et al. (2014) Production of an alkaline protease using *Bacillus pumilus* D3 without inactivation by SDS, its characterization and purification. *Journal of enzyme inhibition and medicinal chemistry* 29(3): 388-396.
- Oda K (2012) New families of carboxyl peptidases: serine-carboxyl peptidases and glutamic peptidases. *The Journal of Biochemistry* 151(1): 13-25.
- Rao MB, Tanksale AM, Ghatge MS, Deshpande VV (1998) Molecular and biotechnological aspects of microbial proteases. *Microbiology and molecular biology reviews* 62(3): 597-635.
- Vishalakshi N, Lingappa K, Amena S, Prabhakar M, Dayanand A (2009) Production of alkaline protease from *Streptomyces gulbargensis* and its application in removal of blood stains.
- Raj A, Khess N, Pujari N, Bhattacharya S, Das A, et al. (2012) Enhancement of protease production by *Pseudomonas aeruginosa* isolated from dairy effluent sludge and determination of its fibrinolytic potential. *Asian Pacific Journal of Tropical Biomedicine* 2(3): S1845-S1851.
- Chaudhari S, Chaudhari S, Khobragade R, Deshmukh A (2013) Studies on production of alkaline proteases from *Bacillus* species and its application in detergent industry. *DAV International Journal of Science* 2: 44-48.
- Sundararajan S, Kannan CN, Chittibabu S (2011) Alkaline protease from *Bacillus cereus* VITSN04: potential application as a dehairing agent. *Journal of bioscience and bioengineering* 111(2): 128-133.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*. Cold spring harbor laboratory press, United States.
- Das G, Prasad M (2010) Isolation, purification & mass production of protease enzyme from *Bacillus Subtilis*. *Int Res J Microbiol* 1(2): 26-31.
- Ibrahim AS, Al-Salamah AA, Elbadawi YB, El-Tayeb MA, Ibrahim SSS (2015) Production of extracellular alkaline protease by new halotolerant alkaliphilic *Bacillus* sp. NPST-AK15 isolated from hyper saline soda lakes. *Electronic Journal of Biotechnology* 18(3): 236-243.
- El-Safey E, Abdul-Raouf U (2004) Production, purification and characterization of protease enzyme from *Bacillus Subtilis*. In: *International Conferences for Development and the Environment in the Arab World*, Assiut Univ 2: 23-25.

20. Banik RM, Prakash M (2004) Laundry detergent compatibility of the alkaline protease from *Bacillus cereus*. *Microbiological research* 159(2): 135-140.
21. Gençkal H, Tari C (2006) Alkaline protease production from alkalophilic *Bacillus* sp. isolated from natural habitats. *Enzyme and Microbial Technology* 39(4): 703-710.
22. Shaikh I, Dixit P, Shaikh T (2018) Purification and characterization of alkaline soda-bleach stable protease from *Bacillus* sp. APP-07 isolated from Laundromat soil. *Journal of Genetic Engineering and Biotechnology* 16(2): 273-279.
23. Mala M, Srividya S (2010) Partial purification and properties of a laundry detergent compatible alkaline protease from a newly isolated *Bacillus* species Y. *Indian journal of microbiology* 50(3): 309-317.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2021.35.005750

Prittesh Patel. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>