

Advantages of the Topical Application of Ozoile in the Healing of Venous Ulcers of the Lower Limbs. A Randomized Clinical Study

Gaia Zagolin¹, Greta Loss¹, Matilde Tasinato¹, Roberto Cassino¹, Gaia Degli Angeli¹ and Rolando Tasinato^{1*}



¹General and Vascular Surgery Department, AULSS 3, Mirano General Hospital (Venice) Italy

²RSA Korian Sacra Famiglia – Pieve del Cairo (Retorbido (Pavia) Italy

*Corresponding author: Rolando Tasinato MD, RSA Korian Sacra Famiglia – Pieve del Cairo (Retorbido (Pavia) Italy

ARTICLE INFO

Received: 📅 November 29, 2022

Published: 📅 December 08, 2022

Citation: Gaia Zagolin, Greta Loss, Matilde Tasinato, Roberto Cassino, Gaia Degli Angeli, et al. Advantages of the Topical Application of Ozoile in the Healing of Venous Ulcers of the Lower Limbs. A Randomized Clinical Study. Biomed J Sci & Tech Res 47(3)-2022. BJSTR. MS.ID.007516.

ABSTRACT

This paper reports a randomized clinical trial about 81 patients with phlebostatic ulcers of the lower limbs who underwent treatment with topical application of Ozoile. The results were compared with a control group which did not receive the treatment. The trial lasted 9 months overall. Over this period of time, patients treated with Ozoile showed a statistically meaningful reduction (P value > 0.007) of mean re-epithelialization time compared to the control group. Biopsies of the floor and margins of the ulcers treated topically with Ozoile were performed and analyzed histologically and demonstrated that topical stable ozonides from biological olive oil application stimulates fibroblasts proliferation and neoangiogenesis. Such mechanisms, combined with debridement and compression therapy, make for an efficient treatment strategy for phlebostatic ulcers, providing treated patients with a reduction of healing time and better quality of life.

Keywords: Wound Care; Venous Ulcers; Ozone Therapy; Ozonized Oil; Bacteria; Neoangiogenesis; Chronic Cardiopathy; Sodium Hypochlorite; Infection; Phlebostatic; Neoangiogenesis

Introduction

During the latest years, several experiences regarding topical ozone application for the treatment of chronic cutaneous lesions were reported in scientific literature. Ozone's antiseptic properties have been exploited to reduce bacterial and microbial load mostly in bacteria-infected wounds and at the floor level of chronic cutaneous ulcers, both in acute and chronic phases [1]. In dermatology, ozone is used for the treatment of skin infections, especially those caused by pathogens that have developed resistance to conventional therapeutic approaches with antibiotics or antimycotic medications [2,3]. In addition, ozone stimulates regenerative processes that facilitate re-epithelialization of chronic skin ulcers. When bound to oil-rich substances, it remains firmly in contact with tissues,

releasing molecular oxygen which can then exert its bacteriostatic and antiseptic activity on bacteria, anaerobes especially [4]. It was demonstrated that the presence of ozone in cultured tissues induces the production of endothelial growth factors, such as VEGF (Vascular Endothelial Growth Factor), that lead to the formation of new vessels, which improve blood flow to the floor and margins of the lesions. In vitro, ozone was also shown to induce fibroblasts migration towards the floor of the ulcer and facilitate neoangiogenesis. Different mixing techniques suggested in the past, although efficient, were often unable to supply stable preparations with progressive and controlled ozone release. Topical application of ozone-releasing preparations caught.

The interest of many teams working with acute and chronic skin lesions in the field of wound care [5,6]. Recently, we evaluated the use of Ozoile, a biological inducer obtained from a patented mixing process in which ozone is bound to olefinic bonds of fatty acids of a biological olive oil forming stable ozonides. The formula we used for the study is patented, produced and distributed in Italy by Erbagil S.r.l.

Materials and Methods

Between August 2021 and May 2022, a total of 81 patients was evaluated. These patients (43 men and 38 women aged 64.7 years on average, in a 49-83-year range) presented with chronic skin ulcers of the lower limbs of vascular origin. The ulcers had been active for at least 6 weeks and originated from venous vessels. The venous etiology of the lesions was documented in each case by clinical examination, Color-Doppler imaging and reflection photoplethysmography. Inclusion criteria comprised: being over 18 years of age, suffering from severe chronic venous insufficiency graded C6 on the CEAP classification and absence of other relevant comorbidities (such as NYHA class III or IV chronic cardiopathy, insulin-dependent diabetes mellitus, obesity with a BMI > 40, active neoplasm, COPD, concurrent treatment with corticosteroid or immunosuppressor drugs, legs arteriopathy with an ABI index < 0.7). Patients were subsequently randomized into two groups. Patients in group A (41 patients) were treated following the TIME protocol according to EWMA's (European Wound Management Association) directions. Besides ulcer debridement, patients in this group underwent both compression therapy and application of Ozoile oil spray and Ozoile cream. Ozoile oil spray was applied to the floor of the ulcer, while the Ozoile-containing emulsion was applied to the perilesional skin. Group B patients were likewise treated with debridement according to the TIME model and cleansing with repeated washes of saline solution.

Followed by application of sodium hypochlorite 0.05% solution for about 3 minutes and compression therapy, but they did not undergo application of the Ozoile products. Medications were performed every 3 days on average. Short-stretch elastic cohesive bandages were applied, using the multi-layer technique, starting from the metatarsal region and proceeding upwards towards the knee, excluding the latter. After being applied the bandage, each patient was asked to walk, being given no functional limitation to his or her motor activity. The two groups of patients

were treated by the same team, composed of 2 physicians and 3 nurses. Once the medication which was object of randomization had been performed, elastic bandages were applied following the same technique, with no differences between group A and group B. Patients randomization and data collection was performed independently through the use of a FH VMa Electronic Case Report Form (eCRF). All patients consented to the treatment and the two groups proved to be homogeneous in anthropological features and biographical data at statistical analysis. The patients randomized in the study were subjected to two weekly evaluations on average, every 3-4 days, to check on the status of their lesions scarring process and renew medication and bandages. Every week, ulcers size, depth and exudate entity were measured for each patient, and the acquired data set was reported in the medical records. The results of these periodic measurements were then inserted into an electronic database accompanied by photographic evidence.

The aim of the study was to investigate the presence of statistically significant differences in ulcers healing rate among the two groups.

Results

Patients were observed for a period of 10 months overall, during which 2 patients abandoned the study because of intercurrent SARS-CoV-2 infection. Final data refer to 79 patients (40 from group A, 39 from group B) who successfully concluded the therapeutic protocol showing healing of the ulcer. The trend for re-epithelialization time observed in the two groups is reported in (Figure 1). Phlebostatic ulcers healing rate was found to be higher in group A patients (patients treated with Ozoile) than in group B patients (control group). Independent statistical analysis of the data set shows that at 12 weeks past T0 mean healing time was reduced by 24.2%, at 16 weeks from T0 it was reduced by 26.3% and at 18 weeks by 15.1%. Overall, compared to the control group, the group of patients treated with both compression therapy and Ozoile topical application showed a mean reduction of about 23.8%. Mild atopic reaction to the Ozoile-containing cream was observed in group A, but treatment suspension was not necessary. In addition, in group A patients relied on analgesic therapy to a lesser extent than patients in group B and exudate control was more efficient. This aspect led the subjects to a faster return to their working activity and gave both patients and caregivers a better perception of their own quality of life.




Figure 1: Diagram analyzing the trend of the percentage of chronic venous ulcers that healed comparing group A and group B: patients in group A treated with Ozoile and patients in group B treated with standard treatment.

Discussion

In a first phase, Ozoile application to the floor of the ulcers shows faster debridement of the treated lesions. Ozone release into the cellular medium favors cleansing of the floor from bacterial and fungal populations in the biofilm both via stimulation of autolytic debridement and via antiseptic action of ozone itself [7,8]. In our study, this action characterizes the first 2-3 weeks of treatment (Figure 2). Some patients underwent histological examination with a 5 mm punch biopsy sampling the floor and margins of the lesion. Optical microscope examination then allowed to highlight the efficacy of Ozoile-containing preparations in promoting reparative processes, mostly granulation tissue formation and neoangiogenesis accompanying the process of re-epithelialization [9,10]. Evaluation of the histological images shows that when applied to the ulcer floor and to perilesional skin Ozoile is capable of stimulating activation of cellular lines responsible for tissue repair in chronic ulcerative lesions [11] (Figure 3). Moreover,

qualitative and quantitative examinations of the cell populations revealed through microscopy confirm that progressive application of the Ozoile-based preparation has anti-inflammatory effect, followed by a remarkable reduction in the number of inflammatory response-related cells. This effect can be observed starting from the third week of application on average and lasts up until the scarring process is complete [12-14]. By stimulating repair processes, Ozoile application is responsible for a temporary flareup of the lesion.

This temporary transformation of the lesion from chronic to acute paves the way for the initialization of the re-epithelialization process carried out mainly through fibroblasts migration and subsequent formation of newly-vascularized epithelium [15-19]. Stimulation of the repair processes is also noticeable in the histological examinations of patients in group B, who were not treated with Ozoile, but cell populations involved in scar formation and neoangiogenesis appear later and less remarkably than in group A (Figure 4).




Figure 2: Optical microscopy images obtained from incisional biopsies of the perilesional skin of a patient after 4 applications of Ozoile oil Spray to the floor of the lesion and Ozoile cream to the perilesional skin. The biopsies were performed 21 days after application of the medication (Fig. A and B T0 – Fig. C and D T1): After 3 weeks, the epithelium appears more structured and stratified and several newly-formed small vessels appear in the connective tissue under the derma, indicating the presence of active neon angiogenesis.




Figure 3: Histological sample with hematoxylin and eosin stain. The biopsy was obtained from the floor of a chronic ulcerative skin lesion of phlebostatic origin. The patient (male, 63 years old) was treated with sequential applications of Ozoile oil Spray and Ozoile cream to the perilesional skin. Observation at T0 is reported in Fig. A: besides tissue dismantlement, notable bacterial contamination can be seen. In Fig. B, representing a biopsy obtained after 4 weeks of treatment, there appear areas of granulation tissue and neo angiogenesis.




Fig A




Fig B

Figure 4: Higher magnification detail. Active fibroblasts proliferation and neoangiogenesis can be observed.

Conclusion

Debridement and compression therapy are regarded as the gold standard treatment for chronic venous ulcers [20,21]. The duration of the treatment is influenced by the quality of the medication. In our experience, the use of stable ozonides from biological olive oil (Ozoile)-containing preparations offers remarkable advantages by means of increasing ulcer re-epithelialization. In addition, Ozoile stimulates epithelial regenerative processes [22,23] and reduces bacterial load at the floor of the ulcer.

References

- Fitzpatrick E, Holland OJ, Vanderlelie JJ (2018) Ozone therapy for the treatment of chronic wounds: A systematic review. *Int Wound J* 15(4): 633-644.
- Zeng J, Lu J (2018) Mechanisms of action involved in ozone-therapy in skin diseases. *Int Immunopharmacol* 56: 235-241.
- Lim Y, Lee H, Woodby B, Valacchi G (2019) Ozonated Oils and Cutaneous Wound Healing. *Curr Pharm Des* 25(20): 2264-2278.
- Borges GÁ, Elias ST, Da Silva SM, Magalhães PO, Macedo SB, et al. (2017) In vitro evaluation of wound healing and antimicrobial potential of ozone therapy. *Craniomaxillofac Surg* 45(3): 364-370.
- Krkl C, Yiğit MV, Özercan İH, Aygen E, Gültürk B, et al. (2016) The Effect of Ozonated Olive Oil on Neovascularization in an Experimental Skin Flap Model. *Adv Skin Wound Care* 29(7): 322-327.
- Kim HS, Noh SU, Han YW, Kim KM, Kang H, et al. (2009) Therapeutic effects of topical application of ozone on acute cutaneous wound healing. *J Korean Med Sci* 24(3): 368-74.
- Silva V, Peirone C, Amaral JS, Capita R, Alonso-Calleja C, et al. (2020) High Efficacy of Ozonated Oils on the Removal of Biofilms Produced by Methicillin-Resistant *Staphylococcus aureus* (MRSA) from Infected Diabetic Foot Ulcers. *Molecules* 25(16): 3601.
- Xiao W, Tang H, Wu M, Liao Y, Li K, et al. (2017) Ozone oil promotes wound healing by increasing the migration of fibroblasts via PI3K/Akt/mTOR signaling pathway. *Biosci Rep* 37(6): BSR20170658.
- Izadi M, Kheirjou R, Mohammadpour R, Aliyoldashi MH, Moghadam SJ, et al. (2019) Efficacy of comprehensive ozone therapy in diabetic foot ulcer healing. *Diabetes Metab Syndr* 13(1): 822-825.
- Valacchi G, Zanardi I, Lim Y, Belmonte G, Miracco C, et al. (2013) Ozonated oils as functional dermatological matrices: effects on the wound healing process using SKH1 mice. *Int J Pharm* 458(1): 65-73.
- Monzillo V, Lallitto F, Russo A, Poggio C, Scribante A, et al. (2020) Ozonized Gel Against Four *Candida* Species: A Pilot Study and Clinical Perspectives. *Materials (Basel)* 13(7): 1731
- Solovăstru LG, Stîncanu A, De Ascentii A, Capparé G, Mattana P, et al. (2015) Randomized, controlled study of innovative spray formulation containing ozonated oil and α -bisabolol in the topical treatment of chronic venous leg ulcers. *Adv Skin Wound Care* 28(9): 406-409.
- Zeng J, Lu J (2018) Mechanisms of action involved in ozone-therapy in skin diseases. *Int Immunopharmacol* 56: 235-241
- Xiao WR, Wu M, Bi XR (2021) Ozone oil promotes wound healing via increasing miR-21-5p-mediated inhibition of RASA1. *Wound Repair Regen* 29(3): 406-416.
- Pietrocola G, Ceci M, Preda F, Poggio C, Colombo M, et al. (2018) Evaluation of the antibacterial activity of a new ozonized olive oil against oral and periodontal pathogens. *J Clin Exp Dent* 10(11): e1103-e1108.
- Ugazio E, Tullio V, Binello A, Tagliapietra S, Dosio F, et al. (2020) Ozonated Oils as Antimicrobial Systems in Topical Applications. Their Characterization, Current Applications, and Advances in Improved Delivery Techniques. *Molecules* 25(2): 334.
- Xiao W, Tang H, Wu M, Liao Y, Li K (2017) Ozone oil promotes wound healing by increasing the migration of fibroblasts via PI3K/Akt/mTOR signaling pathway. *Biosci Rep* 37(6): BSR20170658.
- Tualzik T, Chopra R, Gupta SJ, Sharma N, Khare M, et al. (2021) Effects of ozonated olive oil and photo biomodulation using diode laser on gingival depigmented wound: A randomized clinical study. *Indian Soc Periodontol* 25(5): 422-426.
- Cho KH, Kang DJ, Nam HS, Kim JH, Kim SY, et al. (2021) Ozonated Sunflower Oil Exerted Protective Effect for Embryo and Cell Survival via

- Potent Reduction Power and Antioxidant Activity in HDL with Strong Antimicrobial Activity Antioxidants (Basel) 10(11): 1651.
20. Zerillo L, Polvere I, Varricchio R, Madera JR, D'Andrea S, et al. (2022) Antibiofilm and repair activity of ozonated oil in liposome. Microb Biotechnol 15(5): 1422-1433.
21. Prządka P, Kuberka M, Skrzypczak P, Kiełbowicz Z (2022) Healing of a large skin defect in a dog with concurrent ozonated olive oil application. J Small Anim Pract 63(6): 492.
22. Cipriani F, Lucattelli E, De Rosa M, Di Lonardo A (2021) Effectiveness Of Reactive Oxygen Species In An Oil-Based Medication For Healing Burn Wounds: A Case Series. Ann Burns Fire Disasters 34(2): 170-174.
23. Wen Q, Liu D, Wang X, Zhang Y, Fang S, et al. (2022) A systematic review of ozone therapy for treating chronically refractory wounds and ulcers. nt Wound J 19(4): 853-870.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2022.47.007516](https://doi.org/10.26717/BJSTR.2022.47.007516)

Rolando Tasinato. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>