

Usefulness of Procalcitonin in the Suspension of Antibiotics

Elisa Quintero Rincones^{1*}, Jairo Puello Gonzales¹, Josmar Cera Diaz², Juan Castilla Ortega³, Elisa De La Rosa Povea³, Jesús De Oro Acevedo¹ and Katherine Contreras Padilla¹

¹General Practitioner, Corporación Universitaria Rafael Nuñez, Colombia

²General Physician, Universidad del Sinu, Colombia

³General Physician, Universidad de Sucre, Colombia

***Corresponding author:** Elisa Quintero Rincones, General Practitioner, Corporación Universitaria Rafael Nuñez, Cartagena de Indias, Colombia

ARTICLE INFO

Received: 📅 February 26, 2024

Published: 📅 March 07, 2024

Citation: Elisa Quintero Rincones, Jairo Puello Gonzales, Josmar Cera Diaz, Juan Castilla Ortega, Elisa De La Rosa Povea, Jesús De Oro Acevedo and Katherine Contreras Padilla. Usefulness of Procalcitonin in the Suspension of Antibiotics. Biomed J Sci & Tech Res 55(3)-2024. BJSTR. MS.ID.008703.

ABSTRACT

Background: Procalcitonin is a promising biomarker for the early detection of systemic bacterial infections; its use as a predictor for antibiotic suspension has been studied and has become relevant.

Methodology: A systematic review was carried out through various databases from January 2011 to February 2022; The search and selection of articles was carried out in indexed journals in English. The following keywords were used: Procalcitonin; infections; Sepsis; Bacteria; Virus.

Results: Currently there are different types of inflammatory markers, among which we can highlight procalcitonin, which has become a new biomarker for the early detection of bacterial infections, especially systemic infections. Procalcitonin levels have been shown to increase over 6 to 12 hours, and these values continue to steadily increase 2 to 4 hours after the onset of sepsis. After identifying the condition, it is advisable to measure procalcitonin every 6 to 24 hours, in order to identify the need to change the antibiotic.

Conclusions: The present review offers updated and detailed information on the importance of procalcitonin as a discriminating agent for viral diseases of bacteria and thus whether or not to suspend the antibiotic.

Keywords: Procalcitonin; Infections; Sepsis; Bacteria; Viruses; Bacteria

Introduction

Today there are different laboratory tests for the detection of infectious diseases, including PCR, but they lack the specificity to accurately diagnose bacterial versus non-bacterial infections [1,2]. Procalcitonin is a promising biomarker for the early detection of systemic bacterial infections. Procalcitonin is a 116 its diagnostic significance was not recognized until 1993 [3,4]. In 1993, Assicot, et al. demonstrated a positive correlation between high serum Procalcitonin levels and patients with positive results for bacterial infection and sepsis (e.g. positive blood cultures). It was also shown that these levels do not increase with viral infection, but decrease with antibiotic administration [5,6]. Under normal conditions of homeostasis, procalcitonin undergoes initial synthesis by thyroid C cells. Later, this peptide is transformed into procalcitonin through cleavage of a

25 amino acid signal sequence by endopeptidases. The final product, calcitonin, the 32 amino acid hormones responsible for the regulation of serum calcium, is formed after conversion by the enzyme prohormone convertase [7,8]. Infectious diseases remain a major cause of death, disability, and social and economic disruption for millions of people worldwide. Prevention and treatment strategies for infectious diseases are derived from a thorough understanding of the complex interactions between specific viral or bacterial pathogens and the human host [9,10].

Viruses bind to host cells as a prerequisite for intracellular entry and replication. Because cell surfaces are especially enriched with glycans, numerous viruses target glycans for attachment and entry into cells [8,11]. Children with infectious diseases, both bacterial and viral, are often treated with empirical antibiotics as the first line. Consider-

ing both the threat of microorganisms and the toxicity of antibiotics, it is imperative to develop tests at the point of primary care as well as at the in-hospital level to discriminate bacterial from viral infections and to define the indications for antibiotic treatment [7,10-12]. It is therefore appropriate to perform this work in order to identify the importance of procalcitonin as a discriminating agent of viral diseases from bacteria and thus the suspension or not of the antibiotic.

Materials and Methods

A systematic review was carried out, searching PubMed, Scielo and ScienceDirect databases, among others. The collection and selection of articles was carried out in English- language indexed journals from 2011 to 2022. As keywords, the following terms were used in the databases according to DeCS and MeSH methodology: Procalcitonin; infections; Sepsis; Bacteria; Viruses. In this review, 102 original and review publications related to the topic studied were identified, of which 32 articles met the specified inclusion requirements, such as articles that were not less than the year 2011, that were full-text articles and that reported on procalcitonin as a predictor of the interruption or not of antibiotic therapy. Exclusion criteria took into account that the articles did not have sufficient information and that they did not present the full text at the time of review.

Results

Procalcitonin as an Inflammatory Marker

Currently there are different types of inflammatory markers, among which we can highlight procalcitonin, which has become a new biomarker for the early detection of bacterial infections, especially systemic infections. Procalcitonin does not increase in viral infections, decreasing its values when appropriate antibiotic therapy is administered [13,14]. One of the markers or acute phase reactants most frequently sent to health centers is C- reactive protein, which has a disadvantage in that it lacks the specificity needed to accurately diagnose bacterial versus non-bacterial infections [15,16]. At blood level we can identify normal procalcitonin levels (values lower than 0.05 ng/mL), being elevated or influenced by endotoxins or cytokines such as interleukin 6 (IL-6), tumor necrosis factor (TNF)-alpha and IL-1b, increasing procalcitonin levels up to 100 to 1000 times their normal value amino acid residue that was first explained by Le Moullec, et al. in 1984; however, [17,18].

The main source of procalcitonin is found at the thyroid level, in thyroid C cells, but it can also be synthesized by extra-thyroid tissues [10,15,18]. In contrast, cytokines, such as interferon (INF)-gamma, which are released after viral infection, lead to a negative regulation

of PCT, highlighting another advantage of PCT assays [19,20]. After an initial bacterial infection, procalcitonin levels have been shown to increase from 6 to 12 hours, and these values continue to rise steadily from 2 to 4 hours after the onset of sepsis. Twenty to 24 hours is the half-life that has been recorded so far. One of the ways to demonstrate the efficacy of antibiotics is to identify procalcitonin every so often, since if the antibiotic is effective these levels decrease by 50% every 24 hours [20,21]. It must be taken into account that not in all sepsis states it is advisable to identify the procalcitonin level if it is associated to a factor that may hinder its identification, among these factors we can identify the following: [22-24].

1. Trauma
2. Burns
3. Carcinomas (medullary C-cell, small cell lung and bronchial carcinoid)
4. Immunomodulatory therapy that increases pro-inflammatory cytokines
5. Cardiogenic shock
6. First 2 days of life of the newborn
7. During peritoneal dialysis treatment
8. In cirrhotic patients.

Procalcitonin as a Predictor of Antibiotic Initiation and Discontinuation

As previously stated, procalcitonin levels will decrease by 50% in 24 hours after the initiation of the antibiotic, thus demonstrating that, if the antibiotic is not effective in the patient, procalcitonin levels will not decrease to the value necessary to conclude that the antibiotic is effective in the patient. Therefore, this method or these values could be considered as physiological control of the systemic infection presented by the patient [14,20]. Most research has shown that PCT levels show clinical significance when they are in the range of 0.1 to 0.5 ng/mL. In addition, research has shown that PCT levels below 0.1 ng/mL have a high negative predictive value (96.3 %) for excluding bacterial infections [25]. Procalcitonin has several cut-off levels applicable in different conditions or diseases that could predict the course of the disease or the cause of the disease. Table 1 shows the main conditions that could be identified or ruled out by procalcitonin [13,20,22,26,27]. After identifying the condition, it is advisable to measure procalcitonin every 6 to 24 hours, in order to identify the need to change the antibiotic, increase the antibiotic dose, or discontinue the antibiotic implemented for the condition.

Table 1: Conditions that could be identified or ruled out by procalcitonin.

Systemic Disease or Condition	Procalcitonin Levels	Function of Procalcitonin
Arthritis	0.1 to 0.25 ng/mL	Discriminate infectious (septic) arthritis from non-infectious arthritis.
Infections Bacteremia	0.25 ng/mL	To rule out bacteremic infections
Bloodstream infection (primary):	0.1 ng/mL	Differentiating between a true infection and a contaminated sample
Bronchitis (acute)/ Chronic Obstructive Pulmonary Disease (COPD) Exacerbations:	0.1 to 0.5 ng/mL	Reduce (unnecessary) antibiotic exposure in the emergency department and hospital setting without adverse outcomes.
Endocarditis	2.3 ng/mL	High diagnostic accuracy for predicting acute endocarditis.
Meningitis	0.5 ng/mL	Differentiate viral from bacterial meningitis and subsequently reduce antibiotic exposure.
Neutropenia	0.1 to 0.5 ng/mL	To identify systemic bacterial infections in neutropenic patients.
Pneumonia	0.1 to 0.5 ng/mL	Reduce antibiotic exposure during hospitalization without adverse outcomes.
Postoperative fever	0.1 to 0.5 ng/mL	Differentiate postoperative infections from non-infectious fever.
Postoperative infections	0.5 to 1.0 ng/mL	Minimize antibiotic treatment in the surgical ICU without detrimental results.
Severe sepsis/shock	0.25 to 0.5 ng/mL	Limit antibiotic treatment in the ICU without detrimental results.
Upper Respiratory Tract Infections	0.1 to 0.25 ng/mL	Limit antibiotic treatment in the ICU without detrimental results.
Urinary tract infections	0.25 ng/mL	Determine the extent of renal involvement.
Ventilator-associated pneumonia	0.1 to 0.25 ng/mL	Minimize antibiotic treatment without detrimental results.

If procalcitonin levels decrease by 50% after the condition and the initiation of the antibiotic, it is considered that the antibiotic therapy is taking effect [26,27]. It has also been proposed that one of the criteria for discontinuing the antibiotic is when procalcitonin levels decrease or fall below 0.1 ng/ml or 80-90% below the initial measurement [28]. Efficient diagnosis of bacterial infections allows physicians to initiate antibiotic therapy when appropriate, thus avoiding misuse and overuse of antibiotics. As antibiotic resistance continues to increase, it has become increasingly important for physicians to determine different algorithms and laboratory tests to help maintain current antibiotic parameters [29]. Unfortunately, most first-line tests for determining infection, such as blood cultures and C-reactive protein (CRP), lack the efficiency and specificity needed to treat patients promptly. Therefore, laboratory tests are still being implemented to adjust cut-off points for procalcitonin to be used promptly and with greater efficacy in clinical practice [28-30].

Discussion

The study by Joshua et al, in which they conducted a systematic search of databases using an a priori strategy. Systematic reviews were included that reported an outcome related to antibiotic initiation and/or duration in the setting of respiratory infection. Data extraction was performed by the first author and independently verified by a second author. This paper concluded that the use of procalcitonin leads to a reduction in antibiotic initiation for respiratory illness. It also results in a decrease in antibiotic duration [31]. Another study by Anna et al, concluded that although there are several randomized clinical trials and observational trials evaluating the usefulness of

procalcitonin in the early diagnosis of septic events in different clinical settings, there is still uncertainty as to its use in initiating and suspending antibiotic therapy. Since there are different factors that could stimulate the production of procalcitonin and thus obtain false-positive results, the skill of the treating physician plays an important role in this part [32].

All these trials demonstrate that procalcitonin alone would not be advisable to use, as the physician's skill in dealing with the different types of infections and factors that contribute to increased procalcitonin levels in the blood plays an important role here. A strength of the current study is the methodology implemented, with respect to the literature search, and steps in the selection of relevant articles, quality assessment and data extraction. However, this study has several limitations, which should be taken into account before reaching a conclusion, among these are the lack of evidence from clinical analysis trials to determine with certainty the efficacy of procalcitonin with respect to the interruption or not of antibiotic therapy in different types of infections, both viral and bacterial, so more studies are needed to answer these questions.

Conclusion

Currently there are different types of inflammatory markers, among which we can highlight procalcitonin, which has become a new biomarker for the early detection of bacterial infections, especially systemic infections. Procalcitonin does not increase in viral infections, decreasing its values when appropriate antibiotic therapy is administered. The main source of procalcitonin is found at the thyroid level, in thyroid C cells, but it can also be synthesized by extra-thyroid

tissues, such as the liver, pancreas, kidneys, lungs, intestine and leukocytes. After an initial bacterial infection, procalcitonin levels have been shown to increase from 6 to 12 hours, and these values continue to rise steadily from 2 to 4 hours after the onset of sepsis. Twenty to 24 hours is the half-life that has been recorded so far. One of the ways to demonstrate the efficacy of antibiotics is to identify procalcitonin every so often, since if the antibiotic is effective these levels decrease by 50% every 24 hours. After identifying the condition, it is advisable to measure procalcitonin every 6 to 24 hours, in order to identify the need to change the antibiotic, increase the dose of the antibiotic, or discontinue the antibiotic implemented for the condition. If procalcitonin levels decrease by 50% after the condition and the initiation of the antibiotic, it is considered that the antibiotic therapy is working.

References

- Riedel S, Melendez JH, An AT, Rosenbaum JE, Zenilman JM (2011) Procalcitonin as a marker for the detection of bacteremia and sepsis in the emergency department. *Am J Clin Pathol* 135(2): 182-189.
- Lippi G, Sanchis-Gomar F (2017) Procalcitonin in inflammatory bowel disease: drawbacks and opportunities. *World J Gastroenterol* 23(47): 8283-8290.
- Schuetz P, Albrich W, Mueller B (2011) Procalcitonin for diagnosis of infection and guide to antibiotic decisions: past, present and future. *BMC Med* 9: 107.
- Grace E, Turner RM (2014) Use of procalcitonin in patients with various degrees of chronic kidney disease including renal replacement therapy. *Clin Infect Dis* 59(12): 1761-1767.
- Salinas M, López-Garrigós M, Flores E, Uris J, Leiva-Salinas C (2018) Pilot Group of the Appropriate Utilization of Laboratory Tests (REDCONLAB) working group. Procalcitonin in the Emergency Department: A potentially expensive over-request that can be modulated through institutional protocols. *Am J Emerg Med* 36(1): 158-160.
- Kip MMA, Van Oers JA, Shajiei A, Beishuizen A, Berghuis AMS, Girbes AR, et al. (2018) Cost-effectiveness of procalcitonin testing to guide antibiotic treatment duration in critically ill patients: results from a randomised controlled multicentre trial in the Netherlands. *Crit Care* 22(1): 293.
- Wedzicha JA, Miravittles M, Hurst JR, Calverley PM, Albert RK, et al. (2017) Management of COPD exacerbations: a European Respiratory Society/American Thoracic Society guideline. *Eur Respir J* 49(3).
- Koropatkin NM, Cameron EA, Martens EC (2012) How glycan metabolism shapes the human gut microbiota. *Nat Rev Microbiol* 10: 323-335.
- Lu Q, Li S, Shao F (2015) Sweet talk: protein glycosylation in bacterial interaction with the host. *Trends Microbiol* 23: 630-641.
- Peng W, De Vries RP, Grant OC, Thompson AJ, McBride R, et al. (2017) Recent H3N2 viruses have evolved specificity for extended, branched human-type receptors, conferring potential for increased avidity. *Cell Host Microbe* 21: 23-34.
- Zingg W (2017) Health-care-associated infections in neonates, children, and adolescents: an analysis of paediatric data from the European Centre for Disease Prevention and Control point-prevalence survey. *Lancet Infect Dis* 17: 381-389.
- Van Houten CB (2017) A host-protein based assay to differentiate between bacterial and viral infections in preschool children (OPPORTUNITY): a double-blind, multicentre, validation study. *Lancet Infect Dis* 17: 431-440.
- Philipp Schuetz, Robert Balk, Matthias Briel, Alexander Kutz, Mirjam Christ-Crain, et al. (2015) Economic evaluation of procalcitonin guided antibiotic therapy in acute respiratory infections: a US health system perspective. *Clin Chem Lab Med* 53: 583-592.
- Yutaka Kondo, Yutaka Umemura, Kei Hayashida, Yoshitaka Hara, Morio Aihara, et al. (2019) Diagnostic value of procalcitonin and presepsin for sepsis in critically ill adult patients: a systematic review and meta-analysis. *J Intensive Care* 7: 22.
- Y van der Does, M Limper, K E Jie, S C E Schuit, H Jansen, et al. (2018) Procalcitonin-guided antibiotic therapy in patients with fever in a general emergency department population: a multicenter non-inferiority randomized clinical trial (HiTEMP study). *Clin Microbiol Infect* 24: 1282-1289.
- Fei Peng, Wei Chang, Jian-Feng Xie, Qin Sun, Hai-Bo Qiu, et al. (2019) Ineffectiveness of procalcitonin-guided antibiotic therapy in severely critically ill patients: a meta-analysis. *Int J Infect Dis* 85: 158-166.
- Kyeongman Jeon, Jae Kyung Suh, Eun Jin Jang, Songhee Cho, Ho Geol Ryu, et al. (2019) Procalcitonin-guided treatment on duration of antibiotic therapy and cost in septic patients (PRODA): a multi-center randomized controlled trial. *J Korean Med Sci* 34: e110.
- Yahya Shehabi, Martin Sterba, Peter Maxwell Garrett, Kanaka Sundaram Rachakonda, Dianne Stephens, et al. (2014) Procalcitonin algorithm in critically ill adults with undifferentiated infection or suspected sepsis. A randomized controlled trial. *Am J Respir Crit Care Med* 190: 1102-1110.
- Iankova I, Thompson-Leduc P, Kirson NY, et al. (2018) Efficacy and safety of procalcitonin guidance in patients with suspected or confirmed sepsis: a systematic review and meta-analysis. *Crit Care Med* 46: 691-698.
- Yannick Wirz, Marc A Meier, Lila Bouadma, Charles E Luyt, Michel Wolff, et al. (2018) Effect of procalcitonin-guided antibiotic treatment on clinical outcomes in intensive care unit patients with infection and sepsis patients: a patient-level meta-analysis of randomized trials. *Crit Care* 22: 191.
- Evelien de Jong, Jos A van Oers, Albertus Beishuizen, Piet Vos, Wytze J Vermeijden, et al. (2016) Efficacy and safety of procalcitonin guidance in reducing the duration of antibiotic treatment in critically ill patients: a randomized, controlled, open-label trial. *Lancet Infect Dis* 16: 819-827.
- Nicolas Bréchet, Guillaume Hékimian, Jean Chastre, Charles-Edouard Luyt (2015) Procalcitonin to guide antibiotic therapy in ICU. *Int J Antimicrob Agents* 46: S19-24.
- Albrich WC, Harbarth S (2015) Pros and cons of using biomarkers versus clinical decisions in start and stop decisions for antibiotics in the critical care setting. *Intensive Care Med* 41:1739-1751.
- Filippo Pieralli, Lorenzo Corbo, Arianna Torrigiani, Dario Mannini, Elisa Antonielli, et al. (2017) Usefulness of procalcitonin in differentiating Candida and bacterial blood stream infections in critically ill septic patients outside the intensive care unit. *Intern Emerg Med* 12: 629-635.
- Dominique J Pepper, Junfeng Sun, Chanu Rhee, Judith Welsh, John H Powers, et al. (2019) Procalcitonin-guided antibiotic discontinuation and mortality in critically ill adults. *Chest* 155: 1109-1118.
- Michelle M A Kip, Ron Kusters, Maarten J IJzerman, Lotte M G Steuten, et al. (2015) A PCT algorithm for discontinuation of antibiotic therapy is a cost-effective way to reduce antibiotic exposure in adult intensive care patients with sepsis. *J Med Econ* 18: 944-953.
- Philipp Schuetz, Yannick Wirz, Ramon Sager, Mirjam Christ-Crain, Daiana Stolz, Michael Tamm, et al. (2017) Procalcitonin to initiate or discontinue antibiotics in acute respiratory tract infections. *Cochrane Database Syst Rev* (10): CD007498.
- David T Huang, Donald M Yealy, Michael R Filbin, Aaron M Brown, Chung-

- Chou H Chang, et al. (2018) Procalcitonin-guided use of antibiotics for lower respiratory tract infection. *N Engl J Med* 379: 236-249.
29. Ishan S Kamat, Vignesh Ramachandran, Harish Eswaran, Danielle Guffey, Daniel M Musher (2020) Procalcitonin to distinguish viral from bacterial pneumonia: a systematic review and meta-analysis. *Clin Infect Dis* 70: 538-542.
30. Nikolaos-Dimitrios Pantzaris, Diamantina-Xanthi Spilioti, Aikaterini Psaromyalou, Ioanna Koniari, Dimitrios Velissaris (2018) The use of serum procalcitonin as a diagnostic and prognostic biomarker in chronic obstructive pulmonary disease exacerbations: a literature review update. *J Clin Med Res* 10: 545-551.
31. Joshua, V Maithili, R Joseph, B Gavin (2020) Procalcitonin to Guide Antibiotic Therapy for Respiratory Infection: A Systematic Review of Systematic Reviews. *Open Forum Infect Dis* 7(Suppl 1): S73.
32. Anna Maria Azzini, Romolo Marco Dorizzi, Piersandro Sette, Marta Vecchi, Ilaria Coledan, et al. (2020) A 2020 review on the role of procalcitonin in different clinical settings: an update conducted with the tools of the Evidence Based Laboratory Medicine. *Ann Transl Med* 8(9): 610.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2024.55.008703

Elisa Quintero Rincones. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>