

Methotrexate Involving Drug-Drug Interaction Causing Myelosuppression: A Case Report

Dervershi Trivedi*, Jinkal Mistry, Nirali Kayastha, Harikrishna Thulaseedheran, Bansari Parikh, Shuchi Shukla and Jayesh Trivedi

Research Scholar, Department of Pharmacy Practice, Sal Institute of Pharmacy, Sola-Bhadaj road, Ahmedabad, Gujarat, Ahmedabad, India

*Corresponding author: Dervershi Trivedi, Assistant Professor of Surgery, Dhiraj Hospital, Vadodara. Sumandeep Vidyapeeth, Vadodara, India

ARTICLE INFO

Received: 📅 June 05, 2024

Published: 📅 June 14, 2024

Citation: Dervershi Trivedi, Jinkal Mistry, Nirali Kayastha, Harikrishna Thulaseedheran, Bansari Parikh, Shuchi Shukla and Jayesh Trivedi. Methotrexate Involving Drug-Drug Interaction Causing Myelosuppression: A Case Report. Biomed J Sci & Tech Res 57(1)-2024. BJSTR. MS.ID.008944.

ABSTRACT

Methotrexate is commonly prescribed antimetabolite for rheumatoid arthritis. There are certain side effects at higher doses such as bone marrow suppression, pulmonary toxicity, hematologic toxicity, nephrotoxicity and increased risk of infection. This case reports discuss about the elderly patient who developed a rare presentation about the methotrexate induced pancytopenia. As, patient was on methotrexate since 2 years, leflunomide and etoricoxib, consequently, there can be a possible drug-drug interaction between these three drugs and which may enhance the risk of myelosuppression as well as manifests oral ulcers, diarrhoea, and infection. During the admission to hospital, patient was symptomatically treated with the symptoms of toxicity and also with the antidote of methotrexate, injection leucovorin. Even though the exact mechanism of MTX-induced haematological toxicity remains uncertain but possible cause can be MTX-induced hematopoietic toxicity has been found as excess unbound extracellular MTX. In conclusion, Patients on MTX therapy should be regularly monitored with liver function tests and CBC to identify myelosuppression and avoid the sequelae of pancytopenia.

Keywords: Methotrexate; Pancytopenia; Antidote; Drug – drug interaction; Therapy

Introduction

Methotrexate (MTX) is antimetabolite commonly used in various doses to treat malignancy, severe psoriasis, and rheumatoid arthritis [1]. MTX is prescribed to more rheumatoid arthritis patients than any other biologic medication now in use. It is the most studied and prescribed medication used in combination with other DMARDs, demonstrating clear additive therapeutic efficacy [2]. MTX has been used to treat rheumatoid arthritis (RA) and psoriasis since 1951. However, MTX therapy for RA was not widely used until the 1980s [3]. Although higher doses of MTX, defined as doses of 500 mg/m² or greater are often more beneficial, oral therapy at doses ≥ 15 mg/week, as well producing toxicity and side effects such as bone marrow suppression, pulmonary toxicity, nephrotoxicity, hepatotoxicity, hematologic toxicity, and an increased risk of infections [3]. The combination of MTX and either leflunomide or sulphasalazine (DMARDs) in non-responders RA patients reduce the symptoms but tempt an increase in serious ADRs [4].

Case Presentation

A 73 years old male presented with complains severe watery diarrhoea for 3 days, mouth ulcer, back pain and swelling also with blood report of a week ago showing Hb: 8.2, RBCs: 3.42, WBCs: 500 and Platelets: 10,000. On admission, oral mucositis grade II was examined and personal history was suggestive of increase bowel movement, decrease in appetite. Also on systemic examination, generalized mild tenderness in alimentary system was evident.

Medical and Medication History

Patient had past medical history of right shoulder rheumatoid arthritis for 2 years taking tablet Methotrexate (7.5mg) twice a week, tablet Leflunomide (10mg) once a day, tablet Methylprednisolone (4mg) twice a day, tablet Folic acid (5mg) once a day for 5 days in a week, Etoricoxib (60mg) twice a day, tablet Acetaminophen (650mg) once a day. Additionally, surgical procedure total Knee replacement on left knee was done 5 years ago.

Investigation

Besides, Stool culture test resulted to be positive.

Diagnosis

Patient was diagnosed with Methotrexate-induced Pancytopenia through suggestive symptoms, drug-drug interaction among past medication and abnormal haematological levels.

Treatment

On admission patient was treated with prophylactic antibiotics like injection meropenem (1gm) infused with 10cc NS thrice a day (TDS), injection metronidazole (400mg) TDS, Injection teicoplanin (400mg) once a day (OD), injection fluconazole (200mg) OD; supportive therapies such as anti-emetic like injection ondansetron (4mg) TDS, PPI like injection pantoprazole (40mg) twice a day (BD), normal saline (100ml/hr); anti-diarrheal like capsule racecadotril (100mg) TDS, capsule rifaximin (400mg) TDS; MTX antidote injection leucovorin (50mg with 100cc NS) four times day (QDS). Patient was also transfused with 4 units of platelet rich concentration (PRC). On the 2nd day injection fluconazole was replaced with tablet voriconazole (200mg) twice a day. As well injection romiplostim (500mcg) was given subcutaneously. As stool routine microscopy test resulted positive, injection nitazoxanide was prescribed, injection potassium chloride. On day 3, injection leucovorin was reduce to (25mg) QDS with 100cc NS. Injection PCV (Packed cell volume) was given on 4th day.

Additionally, magnesium serum level was found to be 1.58mg/dl, consequently treated with magnesium supplement, and to treat neutropenia injection filgrastim was given subcutaneously. Betadine gargle was suggested to oral mucositis. Besides, methotrexate was omitted and its level were normal to 0.01 on 4th day of admission, also, haematological parameters were normal in 8 days. Patient was discharge with modified prescription for arthritis that is: capsule pantoprazole+ domperidone, capsule rifaximin, tablet voriconazole, tablet folic acid, tablet potassium, tablet acetaminophen (Table 1).

Discussion

Patients treated for Rheumatoid arthritis with MTX may experience haematological damage, such as myelosuppression, leukopenia, neutropenia, and megaloblastic anemia [3,5]. Pancytopenia is one of the MTX toxicities that is challenging to prevent because it might occur suddenly during therapy [6]. Even though the exact mechanism of MTX-induced haematological toxicity remains uncertain, it has been linked to the development of Arthritis. One cause of MTX-induced hematopoietic toxicity has been found as excess unbound extracellular MTX. Along with age, infections, folic acid insufficiency, hypoalbuminemia, and concomitant drugs are all factors to consider [3]. MTX is a folate antagonist that inhibits dihydrofolate reductase (DHFR), preventing the conversion of dihydrofolate to tetrahydrofolate and thus blocking the synthesis of purines and pyrimidines and, therefore, inhibiting DNA, RNA, and protein synthesis. In addition to inhibiting

DHFR, other mechanisms that are thought to be involved in its effect against autoimmune diseases include selective downregulation of B cells, inhibition of T-cell activation, and inhibition of methyltransferase activity [7]. Polyglutamylation of this drug prolongs its intracellular presence form MTX-PG [8]. Therefore, an increase in polyglutamylation increases the risk of toxicity due to direct and prolonged intracellular exposure. Myeloid megakaryocytes and epithelial polyglutamylation elevate the intracellular levels of MTX and consequently the patients may have experienced ulcers.

The same is for WBC and RBC, which may manifest as neutropenia, infection, also present in this case [8,9]. The most potential adverse reaction is severe myelosuppression, which causes most of the relatively rare fatalities caused by MTX [3]. Looking at pharmacokinetics of MTX has been reported that 95% of MTX is eliminated through renal excretion within 30 hours of administration. MTX and its metabolites 7-hydroxy-methotrexate are eliminated with distal renal tubules with hydrogen ions produced by the hydrogen/potassium ATPase pump in the renal tubules [10]. Moreover, it is rapidly absorbed after oral administration, maximum plasma concentrations (Cmax) being reached within 1.5 hours under fasting conditions and absorbed in the proximal jejunum. Methotrexate binds almost exclusively to the albumin component of plasma proteins. RA patients often have a mean binding percentage of 42-57% [11]. Apart from this, DDIs are the main contributors to ADRs, especially among the elderly, Marinella Patanè et al. says, the combination of MTX and either leflunomide or sulphasalazine in non-responders RA patients reduced pain symptoms but induce serious ADRs [4]. Also, Australia's voluntary adverse drug reaction reporting scheme suggests the Australian Adverse Drug Reactions Advisory Committee (ADRAC) has received 9 reports of well-documented pancytopenia in patients taking leflunomide. Out of which 7 patients were taking concomitant methotrexate 7.5–12.5 mg/week [7]. The mechanism of interaction between MTX and leflunomide is uncertain. Leflunomide inhibits OAT3, BCRP, and OATP1B1/1B3 transporters that contribute to methotrexate disposition and increases the level of MTX in the cells [12,13].

Besides, Human organic anion transporters hOAT1 (SLC22A6) and hOAT3 (SLC22A8) are thought to be mediate renal tubular uptake of methotrexate, because they are expressed in the basolateral membrane of proximal epithelial cells, and mediate transport of methotrexate. Selective cyclooxygenase-2 like etoricoxib inhibits methotrexate transport by hOAT3 affecting decreased excretion of MTX with an increase in risk of Adverse events [4,11-12]. As revealed by the case, medication history of elder patient informs MTX 7.5mg twice a week, leflunomide 10mg once a day and etoricoxib 60mg twice a day; consequently, there can be a possible drug-drug interaction between these three drugs and which may enhance the risk of myelosuppression as well as manifests oral ulcers, diarrhoea, and infection. For antidote of methotrexate toxicity, MTX works by blocking the synthesis of reduced folates; folic acid (leucovorin), which is a fully reduced folate, can restore marrow toxicity. Patients receiving methotrexate are al-

ready encouraged to notify their healthcare provider immediately if they experience any symptoms suggestive of infection. Furthermore, we recommend that patients discontinue methotrexate immediately if such symptoms occur [14].

Conclusion

It is crucial that primary care physicians are aware of these complications and recommendations, because the majority of these serious complications can be detected on time and even prevented before any serious consequences. Patients taking MTX therapy should be regularly monitored with liver function tests and CBC to identify myelosuppression and avoid the sequelae of pancytopenia. Also, renal function must be monitored as this drug uses mainly the kidneys for excretion, especially in elderly patient and educating patient on the importance of MTX and its side effect is an vital task of clinical pharmacist.

References

1. Tao D, Wang H, Xia F, Ma W (2022) Pancytopenia Due to Possible Drug-Drug Interactions Between Low-Dose Methotrexate and Proton Pump Inhibitors. *Drug Healthc Patient Saf*14: 75-78.
2. Joel M Kremer (2004) Toward a Better Understanding of Methotrexate. *Arthritis & Rheumatism* 50(5): 1370-1382.
3. Hamed KM, Dighriri IM (2022) Overview of methotrexate toxicity: A comprehensive literature review. *Cureus* 14(9): e29518.
4. Patanè Marinella, Ciriaco Miriam, Chimirri Serafina, Ursini Francesco, Naty Saverio, et al. (2013) Interactions among Low Dose of Methotrexate and Drugs Used in the Treatment of Rheumatoid Arthritis. *Advances in Pharmacological Sciences*, p. 1-8.
5. Preet Singh Y, Aggarwal A, Misra R, Agarwal V (2007) Low-dose methotrexate-induced pancytopenia. *Clin Rheumatol* 26: 84-7.
6. Gutierrez Ureña S, Molina JF, García CO, Cuéllar ML, Espinoza LR, et al. (1996) Pancytopenia secondary to methotrexate therapy in rheumatoid arthritis. *Arthritis Rheum* 39(2): 272-276.
7. Kanderi T, Janet Chan Gomez, Max M Puthenpura, Keerthi Yarlagadda, Mounika Gangireddy et al. (2020) Pancytopenia as a complication of low-dose methotrexate in a septuagenarian: A rare presentation. *Cureus* 12(6): e8492.
8. JM Kremer, RA Hamilton (1995) The effects of nonsteroidal antiinflammatory drugs on methotrexate (MTX) pharmacokinetics: Impairment of renal clearance of MTX at weekly maintenance doses but not at 7.5 mg, *Journal of Rheumatology* 22(11): 2072-2077.
9. Gonzalez Ibarra F, Eivaz Mohammadi S, Surapaneni S (2014) Methotrexate induced pancytopenia, *Case Reports in Rheumatology*, p. 1-4.
10. Tao D, Hui Wang, Fangfang Xia, Wenlu Ma (2022) Pancytopenia due to possible drug-drug interactions between low-dose methotrexate and proton pump inhibitors, *Drug, Healthcare and Patient Safety* 14: 75-78.
11. Anaya JM, Fabre D, Bressolle F, C Bologna, R Alric, et al. (1994) Effect of etodolac on methotrexate pharmacokinetics in patients with rheumatoid arthritis. *J Rheumatol* 21: 203-208.
12. Bridgewater N], Sanofi Aventis US LLC (2016). Arava (leflunomide) [prescribing information].
13. <https://online.lexi.com/lco/action/interact#analysis-tab-content>.
14. Bernard Bannwarth, Fabienne Péhourcq, T Schaefferbeke, J Dehais (1996) Clinical Pharmacokinetics of Low-Dose Pulse Methotrexate in Rheumatoid Arthritis. *Clinical Pharmacokinetics* 30(3): 194-210.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2024.57.008944

Dervershi Trivedi. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>