

Multimodal Integrative Therapy in a Former NFL Player with Post-Concussive Syndrome and Early-Onset Cognitive Decline: A Case Report

Justin Dearing¹, Melanie Luther², Seamus Kelly³ and Oscar Coetzee^{3*}

¹The Dearing Clinic, Brentwood, TN, USA

²Nutrition with Mel, CO, USA

³Georgetown University, Washington DC

⁴Notre Dame of Maryland University, Baltimore, MD, Designs for Health, Palm Coast FL, USA

*Corresponding author: Justin Dearing, The Dearing Clinic, Brentwood, TN, USA

ARTICLE INFO

Received: 📅 June 29, 2026

Published: 📅 July 08, 2026

Citation: Justin Dearing, Melanie Luther, Seamus Kelly and Oscar Coetzee. Multimodal Integrative Therapy in a Former NFL Player with Post-Concussive Syndrome and Early-Onset Cognitive Decline: A Case Report. Biomed J Sci & Tech Res 66(1)-2026. BJSTR.MS.ID.010291.

ABSTRACT

A middle-aged male presented with severe neck pain, forgetfulness, decreased mental clarity, weight gain, and gastrointestinal discomfort, which led to a reliance on pain medications. His lifestyle choices did not involve regular drinking or smoking. His treating physicians made diagnoses of early-onset dementia, post-concussive syndrome, obesity, cervical spinal enthesopathy, and myalgia in other sites. After diagnostic testing, the patient was advised by a licensed health care provider to follow a personalized nutraceutical regimen, physical rehabilitation, chiropractic, intravenous (IV) regenerative therapy, IV nutrient therapy, and hyperbaric oxygen therapy (HBOT). The supplement regimen consisted of Designs for Health Berberine Synergy™, IgGI Shield™, GI Revive®, Homocysteine Supreme™, Hepatone Plus™, Libidostim-M™, OmegaVail™ Hi-Po, Whole Body Collagen®, Vascanox HP®, Adrenotone™, Tri-Butyrin, Allicillin™, and GI Microb-X™. The protocol was associated with clinically meaningful improvements in cognitive function, quality of life, gut microbiome markers, nutrient status, and neurotransmitter status. This case report describes the clinical outcomes associated with a multimodal integrative treatment approach.

Abbreviations: IV: Intravenous; HBOT: Hyperbaric Oxygen Therapy; TBI: Traumatic Brain Injury; NFL: National Football League; FRQ: Fall Risk Questionnaire; CBC: Complete Blood Count; CNS: Central Nervous System; OAT: Organic Acids Test; LPS: Lipopolysaccharide; SNPs: Single-Nucleotide Polymorphisms; qEEG: Quantitative EEG; SCFAs: Short-Chain Fatty Acids; DATS: Diallyl Trisulfide; DADS: Diallyl Disulfide; DAS: Diallyl Sulfide; NAD⁺: Nicotinamide Adenine Dinucleotide; MSCs: Mesenchymal Stem Cells; PRP: Platelet-Rich Plasma; BDNF: Brain-Derived Neurotrophic Factor; VEGF: Vascular Endothelial Growth Factor; SOD: Superoxide Dismutase; LDL-C: Low-Density Lipoprotein Cholesterol

Introduction

Traumatic brain injury (TBI) and its long-term sequelae represent a major clinical and socioeconomic challenge, particularly among retired professional athletes with a history of repetitive head and neck trauma [1]. Post-concussive syndrome, cognitive decline, and associated metabolic and musculoskeletal disorders frequently coexist, producing a complex clinical picture that is often refractory to conventional therapies. This case presents a 50-year-old male, a former National Football League (NFL) lineman with more than a decade of professional play, who developed severe cervical pain, progressive cognitive impairment, gastrointestinal dysfunction, and significant

metabolic dysfunction following cumulative sub-concussive impacts and a recent fall. The patient's presentation was consistent with multifactorial pathology, including early-onset dementia, post-concussive syndrome, obesity, use of opioid medication, metabolic dysfunction, and cervical enthesopathy. At the beginning of this case, the patient presented with a height of 6'5" and a weight of 320 lb. (BMI 37.9). This baseline measurement was obtained on the clinic scale before Fit3D technology was introduced. Despite standard medical management and pain-control strategies, his condition progressed to functional impairment, with loss of independence in daily life and an inability to care for his family.

This case is clinically significant because it illustrates the potential for a comprehensive, multimodal integrative approach, including targeted nutraceutical supplementation, IV nutrition and regenerative therapy, hyperbaric oxygen therapy, chiropractic rehabilitation, and personalized physical therapy, to achieve substantial improvements in cognition, pain, gastrointestinal health, and overall quality of life. Existing literature on post-concussive and neurodegenerative syndromes highlights chronic neuroinflammation, oxidative stress, mitochondrial dysfunction, and gut-brain axis disruption as key pathophysiologic drivers [2-6]. Prior studies have demonstrated partial benefits of nutritional and rehabilitative interventions; however, reports detailing coordinated, precision-based regimens incorporating high-quality nutraceuticals and adjunctive IV regenerative medicine remain limited [7-10]. This case adds to the growing body of literature describing clinically meaningful improvements in cognitive and systemic symptoms associated with a multimodal integrative treatment approach.

Narrative

Patient Information

A 50-year-old male with a history as a professional NFL lineman presented with severe acute onset of left neck pain, post-concussive syndrome, weight gain, and gastrointestinal upset. The patient reported a history of numerous sub-concussive impacts to the head and neck during his football career. The patient arrived at the clinic after a fall that resulted in a neck and foot injury. Upon examination, a history of mild cognitive impairment and early-onset dementia was uncovered. He presented with memory and cognitive processing problems to the point that he was unable to care for his children and support his wife. The patient had previously received a diagnosis of early-onset dementia. Repetitive head trauma was considering a potential contributing factor, and was likely compounded by regular consumption of pain medication.

Clinical Findings

Physical examination revealed muscle spasms of the right paraspinal musculature with soft tissue trigger points at the right scalene, right suboccipital, and right upper trapezius.

Mental health was stable with no signs of depression, anxiety, or stress, as indicated by the DASS-21 assessment (Depression: 0, Anxiety: 2, Stress: 4 — all within normal range) on March 16, 2023.

Sleep quality was reported as good, supported by an Epworth Sleepiness Scale score of 3 (scores below 6 are considered within normal limits) on March 16, 2023.

The Fall Risk Questionnaire (FRQ) indicated a significant risk for falling with a score of 9, where scores above 4 suggest increased risk (March 16, 2023).

Laboratory Work

The patient completed standard CBC and CMP panels at the beginning of this case, in 2023, as part of routine primary care screening. Repeat CBC/CMP testing was performed in 2024 during an NFL routine player check-up. The final round of testing was completed in 2025 for life insurance purposes; however, only liver markers were provided to his physician to compare to 2023 and 2024 tests (Table 1). These results demonstrated improvements in immune cell stability as well as overall cardiovascular and hepatic function (see Tables 1 & 2). Comparison of the patient's complete blood count (CBC) results reflect improved systemic metabolic and inflammatory status following 18 months of therapy (see Tables 1 & 2). The improvement in white blood cell distribution reflects a normalization of immune function and health of the bone marrow. Neutrophils increased from 47.6% to 60.2%, which indicates a stronger and more efficient first-line defense response.¹¹ Neutrophils are essential for combating acute infections and clearing cellular debris; a rise within the optimal range suggests improved innate immune readiness and recovery capacity [11,12]. Lymphocytes stabilized at 30%, representing balanced adaptive immunity [13]. Stable lymphocyte levels imply that the body is no longer mounting an excessive or prolonged immune response and that both antibody production and cellular immune regulation are functioning appropriately [14]. Meanwhile, monocytes declined from 10% to 7%, suggesting a reduction in chronic inflammatory signaling.

Elevated monocytes often accompany long-standing tissue stress or inflammation, so their normalization supports the interpretation that the prior chronic inflammatory burden has been improved and that tissue repair processes are now predominating over inflammatory ones [15,16]. Collectively, these findings suggest a transition from a state of chronic immune activation to one of immunological balance and restored homeostasis. Because mitochondrial integrity underpins bone marrow vitality, driving energy metabolism, redox balance, and stem cell renewal, the observed hematologic improvements are consistent with enhanced mitochondrial function within marrow and hematopoietic cells. Baseline findings were consistent with immune fatigue and low metabolic activity, as evidenced by a WBC count of 3.7 and mild monocytosis (10.2%). After 18 months, total white blood cell counts normalized (3.7 → 5.1), and differential values balanced (see Table 2). The latest neutrophil-to-lymphocyte ratio of approximately 2:1, in 2024, reflects improved physiologic stress adaptability, from 1.2:1 in 2023 [17]. Collectively, these findings suggest a transition from a state of chronic immune activation to one of immunological balance and mitochondrial resilience. This connection is made because mitochondrial integrity underpins bone marrow vitality, driving energy metabolism, redox balance, and stem cell renewal [18,19]. The observed hematologic improvements are consistent with enhanced mitochondrial function within bone marrow and hematopoietic cells [18,19].

Neurological and Cognitive Testing

In March 2023, a series of Central Nervous System (CNS) Vital Signs assessments was initiated to track brain and nervous system performance. Results indicated scores very low range for reaction time, low for verbal memory, and low-average across neurocognitive index, composite memory, psychomotor speed, cognitive flexibility, processing speed, and executive function — highlighting multiple areas for improvement (see Figure 1). In May 2024, the patient improved to exhibit no areas of very low scoring, with low composite memory and verbal memory, and a low average for visual memory and reaction time. The rest of his scores were in the average or above-average range (Figure 1).

Functional Testing

In December 2023, the patient showed limited improvement, and additional testing was pursued to identify underlying factors contributing to suboptimal progress. A Metabolomics Spotlight Organic Acids Test (OAT) and a Gastrointestinal Spotlight Functional Stool Test were conducted (see Tables 3 & 4). Findings revealed elevated secretory IgA, suggesting heightened immune activity likely in response to fungal marker *Candida* spp [20-22]. *Candida* spp. can promote aldehyde formation in the gut, and shares a relationship with depleted B vitamin status, which was observed in the OAT results [23-25]. Elevated *Pseudomonas* spp. was also detected, an opportunistic histamine-producing gram-negative bacterium that produces lipopolysaccharide (LPS), a compound contributing to neuroinflammation [26,27]. Evidence of *Staphylococcus aureus* overgrowth was also noted, likely secondary to digestive insufficiency [28,29]. Gut barrier integrity was generally good, though elevated Zonulin levels were detected (Table 3). Commensal bacterial populations appeared within normal ranges. The OAT results confirmed low B-vitamin status and neurotransmitter imbalance involving dopamine and norepinephrine, associated with deficiencies in vitamins B2, B3, and magnesium (Table 7). Detoxification pathways were generally adequate, with the exception of low butyric acid levels, again linked to vitamin B3 deficiency. The low butyric acid level observed on the OAT suggests reduced production

of short-chain fatty acids and possible dysbiosis affecting the colonic environment.

Butyrate is critically important for maintaining gut barrier integrity, modulating inflammation, and providing energy to colonocytes. Deficiency can impair mucosal healing, weaken immune tolerance, and promote intestinal permeability [30,31]. Restoring butyrate levels through dietary fiber diversification, probiotic support, and correction of digestive insufficiencies should be a central focus in improving gut health and systemic resilience.

Genetic Findings

On December 28, 2023, a Genetic Spotlight assessment was performed to identify relevant single-nucleotide polymorphisms (SNPs). Results indicated reduced methylation capacity, with *MTHFR* enzyme activity at approximately 30%. The patient demonstrated increased genetic risk for neurodegeneration associated with the *APOE* ϵ 4 allele and *GAB2* variant. *APOE* ϵ 4 — the principal genetic risk factor for late-onset Alzheimer’s disease — also influences lipid metabolism and cardiovascular risk, [32] whereas *GAB2* variants are thought to modify Alzheimer’s risk specifically in ϵ 4 carriers through their role in intracellular signaling [33]. Additional genes associated with elevated neurodegenerative risk were identified, including *IDO1*, *TPH2*, *SOD1*, *SOD2*, *ESR1*, and *HTR2A*. The patient also exhibited the *HLA-DQ2.2* haplotype, conferring moderate risk for generalized inflammation and potential autoimmune reactivity.

Symptoms

The patient reported gastrointestinal dysfunction, dizziness with head and neck movements following a fall in his home, and significant limitation in cervical range of motion. New-onset cognitive symptoms included forgetfulness, slowed reaction time, and reduced verbal memory recall. Myalgia presented as severe acute left-sided neck pain with associated restriction in cervical mobility.

Diagnostics

(Tables 1-6), (Figures 1-3).

Table 1: CBC and Liver Markers.

Marker	2023	2024	2025	% Change	Functional Ideal	Interpretation
WBC	3.7	5.1	—	37.8%	5.0-8.0	Marrow and immune activity normalized
RBC	4.7	4.76	—	1.3%	4.4-4.9	Stable red-cell mass
Hemoglobin	14.8	15.2	—	2.7%	14-15	Improved oxygen-carrying capacity
Hematocrit	45.6	44.3	—	-2.9%	39-55	Stable viscosity
MCV	97	93.2	—	-3.9%	85-92	Closer to ideal, better B-vitamins
MCH	31.5	32	—	1.6%	27-32	Normal hemoglobin loading
MCHC	32.5	34.3	—	5.5%	32-36	Normalized iron and B6 status
Platelets	177	192	—	8.5%	150-300	Enhanced marrow turnover
ALT	36	18	20	-44.4%	<25-30 U/L	Improved liver, detoxification, and methylation
AST	22	17	19	-13.6%	<20-25 U/L	Improved liver and lowered systemic inflammation
ALP	77	55	63	-18.2%	50-90 U/L	Improved bile flow

Table 2: White Blood Cell Differential Markers and Functional Interpretation.

Marker	2023	2024	Functional Ideal	Interpretation
Neutrophils %	47.6	60.2	40-60	From borderline low to optimal innate immune activity
Lymphocytes %	38.1	30.4	25-40	Normalized, less chronic viral activation
Monocytes %	10.2	7.1	4-7	Resolution of chronic inflammatory load
Eosinophils %	2.7	1.8	3-4	Slightly below ideal, mild residual stress tone
Basophils %	0.8	0.5	<1	Stable, no mast-cell activation pattern

Table 3: Gastrointestinal Stool Analysis Findings at Baseline and Follow-up.

Marker	12/28/23	8/23/24	Reference Range	Interpretation
Secretory IgA	3042	1272	510-2010 µg/g	High. Imbalance in keystone diversity, food intolerance. Lowered on follow-up.
Zonulin	158.2	21.0	<175 ng/g	Higher than ideal. Gut barrier integrity marker. Lowered on follow-up.
Pseudomonas spp.	1.11 x 10 ⁴	2.69 x 10 ⁴	<1.00 x 10 ⁴ CFU/g	High. Inflammatory dysbiosis, mast-cell pattern. Increased slightly on follow-up.
Staphylococcus aureus	6.60 x 10 ²	<dl	<5.00 x 10 ² CFU/g	High. Digestive insufficiency, food intolerance. Lowered on follow-up.
Candida spp.	5.53 x 10 ²	3.59 x 10 ²	<5.00 x 10 ³ org/g	Higher than ideal. Fungal imbalance. Lowered on follow-up.
Elastase-1	347	458	>200 µg/g	Within range and improved on follow-up – suggests improved digestive capacity.
Enterococcus spp.	1.34 x 10 ⁶	6.65 x 10 ⁸	1.9 x 10 ⁵ -2.0 x 10 ⁸	Gram-positive; potential concern when too high. Considered opportunistic.
Morganella spp.	<dl	3.2 x 10 ⁵	<1.00 x 10 ³ CFU/g	High. May increase LPS. Gram-negative.
Pseudomonas aeruginosa	<dl	1.93 x 10 ³	<5.00 x 10 ² CFU/g	High. May increase LPS. Gram-negative.

Table 4: Neurotransmitter and Hormone Changes.

Analytes Tested	12/28/23	8/23/24	Ideal	%Change	Direction
Tryptophan	42.8	42.8	10.1-74.3	0.0%	No change
5-Hydroxyindoleacetic Acid	6	203.5	<23.3	3,291.7%	Increase (massive)
Kynurenine	3.3	2.8	<11.6	-15.2%	Decrease
KT Ratio	0.077	0.065	<0.313	-15.6%	Decrease
Kynurenic Acid	17.6	30.6	7.8-54.0	73.9%	Increase
Quinolinic Acid	56.3	68.1	29.4-178.5	20.9%	Increase
Tyrosine	77.5	74.2	<99.0	-4.3%	Decrease (minor)
γ-Aminobutyric Acid	<dl	12.5	<9.5	–	
Homovanillic Acid	20.8	45.6	<42.1	119.2%	Increase
Vanillylmandelic Acid	6	10.5	5.3-36.1	75.0%	Increase
Cortisol	13.9	15	<82.0	7.9%	Increase (minor)

Table 5: Cognitive Domain Improvements.

Cognitive Domain	Baseline (3/16/23)	12/22/23	%Change	Direction
Neurocognitive Index	82	98	19.5%	Increase
Composite Memory	84	91	8.3%	Increase
Verbal Memory	79	80	1.3%	Increase
Visual Memory	95	106	11.6%	Increase
Psychomotor Speed	86	117	36.0%	Increase

Cognitive Domain	Baseline (3/16/23)	12/22/23	%Change	Direction
Reaction Time	67	70	4.5%	Increase
Complex Attention	91	112	23.1%	Increase
Cognitive Flexibility	82	99	20.7%	Increase
Processing Speed	84	109	29.8%	Increase
Executive Function	81	98	21.0%	Increase
Social Acuity	–	112	– (no baseline)	–
Reasoning	–	112	– (no baseline)	–
Working Memory	115	–	– (no 12/22 value)	–
Sustained Attention	108	–	– (no 12/22 value)	–
Simple Attention	94	107	13.8%	Increase
Motor Speed	94	118	25.5%	Increase

Table 6: Fit3D Body Composition Outcomes.

Body Comp. Metric	Baseline (7/24/24)	3/11/25	%Change	Direction
Waist Circumference	51.6"	43.9"	-14.9%	Decrease
Waist-to-Hip Ratio	1.01	1.01	0%	No Change
Trunk-to-Leg Vol. Ratio	1.6	1.4	-12.5%	Decrease
Body Fat %	32.14%	25.09%	-22.0%	Decrease
Weight	296.1 lb	248.8 lb	-15.9%	Decrease
Fat Mass	95.2 lb	62.4 lb	-34.4%	Decrease
Lean Mass	200.9 lb	186.4 lb	-7.2%	Decrease
Basal Metabolic Rate	2320 kcal	2101 kcal	-9.4%	Decrease
Body Shape Rating	14	25	78.6%	Increase

Table 7.

Date	Clinical Event
2023-03-09	Patient arrived after a fall that resulted in mild head trauma, neck pain, and toe injuries. IV vitamin C 12,500 mg and glutathione 1,000 mg were administered
2023-03-16	Initial Central Nervous System (CNS) Vital Signs Report
2023-03-23	Initiated treatment with acupuncture, chiropractic manipulation, physical rehabilitation, and regenerative injections (continued through May)
2023-04-03	Began regenerative tissue injection series (x5 injections of HylaPure and Vitti Pure – monthly through Aug 10)
2023-04-10	Began BPC-157 and GHK-Cu peptide therapy; began NAD+ 500 mg IV series (x4 through Apr 26)
2023-04-25	Second CNS Vital Signs Report
2023-05-01	Initiated treatment of IV antioxidants (continued through September)
2023-05-05	Initiated treatment with hyperbaric oxygen dives 4 days/week, 1- hour sessions, for 10 weeks (continued through July)
2023-05-25	Third CNS Vital Signs Report
2023-06-20	Fourth CNS Vital Signs Report
2023-08-15	Fifth CNS Vital Signs Report
2023-10-02	Initiated Total Gut Restoration Kit (12 weeks) and IV vitamin C (continued through December)
2023-12-22	Sixth CNS Vital Signs Report
2023-12-28	Metabolomics Spotlight OAT Assessment test and DFH Spotlight Functional Stool Test Assessment
2024-01-03	Dysbiosis protocol (6 weeks) included nutraceutical regimen of Berberine Synergy, IgGI Shield, GI Revive, GI Microb-X, Allicilin, and Tri-Butyrin, along with 9 day Rhythm Reset (continued through February)
2024-02-14	Second Total Gut Restoration Kit Protocol (12 weeks), consisting of Microbiome Labs MegaSporeBiotic, MegaPre, and MegaMucosa. Along with Vascanox HP, Adrenotone, and Whole Body Collagen
2024-03-01	qEEG recording – baseline

Date	Clinical Event
2024-04-18	Genetic-guided supplementation (B vitamin status and detoxification) Homocysteine Supreme, Hepatone Plus, Libidostim-M, OmegaVail Hi-Po (Continued through December)
2024-05-27	Seventh CNS Vital Signs Report
2024-07-24	Initial Fit3D Body Scan Report
2024-08-23	Follow-up DFH Metabolomics Spotlight OAT Assessment and GI Spotlight Functional Stool Test Assessment
2024-10-02	qEEG recording – interim
2025-01-08	Second Fit3D Body Scan Report
2025-01-08	Initiated second series of IV NAD+ 500 mg infusions (x6 through Feb 12)
2025-02-12	Third Fit3D Body Scan Report; Final qEEG recording
2025-02-20	Fourth Fit3D Body Scan Report
2025-03-11	Final Fit3D Body Scan Report

CNS Vital Signs Graphical Report		Generated By Justin Dearing on August 26, 2024															
		Above Average		Average		Low Average		Low		Very Low							
Test Time		Neurocognitive Index	Composite Memory	Verbal Memory	Visual Memory	Psychomotor Speed	Reaction Time*	Complex Attention*	Cognitive Flexibility	Processing Speed	Executive Function	Social Acuity	Reasoning	Working Memory	Sustained Attention	Simple Attention	Motor Speed
◆ BaSt01	2023-03-16 06:10:05	82	84	79	95	86	67	91	82	84	81			115	108	94	94
◆ BaSt01	2023-04-25 06:45:38	90	94	82	107	101	58	105	91	91	89					106	109
◆ BaSt01	2023-05-25 06:01:56	89	82	89	83	102	53	111	98	93	98		118	84	93	106	109
◆ BaSt01	2023-06-20 08:47:25	91	53	82	43	108	76	114	106	100	106			121	121	106	110
◆ BaSt01	2023-08-15 07:30:03	75	55	66	61	102	38	97	83	97	82			109	110	106	107
◆ BaSt01	2023-12-22 05:49:40	98	91	80	106	117	70	112	99	109	98	112	112			107	118

Figure 1: Longitudinal CNS Vital Signs cognitive-domain report. Six serial CNS Vital Signs assessments, March–December 2023. Rows are testing dates, columns cognitive domains; cells are color-coded by performance band (above average to very low) from age-adjusted standard scores (mean 100, SD 15). Scores progressed from low-average toward average/above-average; higher scores indicate better performance.





Figure 2: Cognitive performance trajectories (CNS Vital Signs). Age-adjusted standard scores (mean 100, SD 15) for all CNS Vital Signs (CNS VS) domains across six serial assessments, March–December 2023; each line is one domain. Trajectories are broadly upward, with a transient August 2023 decline; reflects the variability typical of serial neurocognitive testing.



Discussion

In this section, the scientific relevance of the constituents, supplement ingredients, and other therapeutics that were used in the 18-month program will be outlined. This will help elucidate some of the potential effects, benefits, and hypotheses underlying the positive outcomes.

Interventions were selected based on clinical outcomes and documented mechanisms in neuro-metabolic recovery:

1. Dysbiosis and Gut Restoration Protocols

Foundation for inflammation control.

2. Total Gut Restoration Kit (12-week Microbiome Labs program)

Barrier repair and SCFA support.

3. Hyperbaric Oxygen Therapy

Mitochondrial oxygenation and angiogenesis.

4. IV Antioxidant Therapy

Redox modulation and microcirculation.

5. NAD and High-Dose Nutrient IVs

Mitochondrial and DNA-repair support.

6. Stem Cell and Exosome Biologics

Regenerative signaling.

7. Rhythm Reset Nutrition Program

Metabolic switching and autophagy activation.

The sequence followed a reduce, repair, restore, and retrain framework.

Dysbiosis Protocol (6 Weeks of Berberine Synergy, IgGI Shield, GI Revive, GI Microb-X, Allicillin, and Tri-Butyrin)

The dysbiosis protocol coincided with sustained improvement of the patient, as he was still adapting his diet following the transition from an NFL-level caloric intake of 6,000–12,000 calories per day to a standard maintenance range of 2,000–3,000 calories. Without this intervention, persistent issues with kynurenine metabolism and neurotransmitter function would likely have remained unresolved. Impaired digestive capacity for proteins, fats, and carbohydrates had contributed to gram-negative bacterial dysbiosis and *Candida* overgrowth, resulting in excessive production of lipopolysaccharides (LPS), oxalates, and sugar alcohols. In combination with an elevated kynurenine-to-tryptophan ratio, these imbalances promoted ongoing formation of neurotoxic metabolites originating in the gut. This intestinal leakiness, inflammation, and endotoxin burden would have impeded continued neurological recovery beyond the period of intensive clinical care.

Berberine Synergy

Berberine Synergy was recommended based on the patient's state of obesity and metabolic syndrome. This supplement provided 400 mg of berberine HCl from *Berberis aristata* with 50 mg of alpha lipoic acid per serving. This supplement was utilized to support metabolic health, as this combination is regarded for its efficacy in supporting blood glucose regulation and healthy insulin metabolism [34]. Berberine exhibits the ability to be supportive in lowering numerous markers of metabolic dysfunction, including fasting blood glucose, hemoglobin A1C, low-density lipoprotein cholesterol (LDL-C), triglycerides, and fasting insulin [35,36]. Berberine may play a role in healthy metabolic status and healthy mitochondria through its ability to support the production of GLP-1 by stimulating intestinal L-cells and mitigating oxidative stress [37,38]. Alpha lipoic acid is a molecule that acts as a cofactor for enzymatic reactions related to blood sugar metabolism and supports mitochondrial energy production by acting directly as a cofactor for pyruvate dehydrogenase and α -ketoglutarate dehydrogenase complexes [39].

IgGI Shield

In order to give more support to the gastrointestinal tract, the patient supplemented with 2.5 g serum-derived bovine immunoglobulin concentrate (ImmunoLin[®]), 1.1 g of immunoglobulin G, and 1 g of N-acetyl-D-glucosamine daily for one month. This combination helps to improve mucosal barrier function, facilitates a balanced and healthy inflammatory response, and improves ability to bind and clear toxins and lipopolysaccharide (LPS) [40,41]. This was deemed necessary based on the high levels of Secretory IgA, an immunoglobulin that binds to toxins and pathogens in the mucosa-associated lymphoid tissues (MALT) and is associated with food intolerances [41]. Further, bovine immunoglobulins have shown an ability to increase the production of short-chain fatty acids (SCFAs) in humans [40].

GI Revive

Improving the integrity of the patient's gut membrane was critical due to the elevated levels of Zonulin, secretory IgA, *Pseudomonas* spp. — a histamine-producing bacterium with an LPS outer wall — and *Staphylococcus aureus*, an opportunistic pathogenic bacterium associated with food intolerances [26,41-45]. GI Revive was utilized due to its multi-ingredient formulation of 15 gut-supportive compounds, including zinc carnosine, L-glutamine, N-acetyl-D-Glucosamine, citrus pectin cellulose, DGL, aloe vera extract, slippery elm extract, mucin, marshmallow root, chamomile extract, okra extract, cat's claw, MSM, quercetin, and prune. Zinc carnosine is a chelated compound of L-carnosine and zinc that helps stabilize the gut mucosa and cell membrane lining, supporting gastric tissue integrity and promoting a healthy inflammatory status within the gut [46–48]. The patient took one scoop daily for one month. Quantitative changes (Table 3) include an 86.7% decrease in Zonulin levels, a 58.2% decrease in secretory IgA, and a decrease of *Staphylococcus aureus* to below detectable limits over a 7-month period. *Pseudomonas* spp. levels remained elevated on follow-up testing.

GI Microb-X

The patient showed an imbalance in *Candida* spp. on his GI test (553 org/g). GI Microb-X is a targeted blend of nutrients and botanicals with a long history of use for supporting a healthy microbial balance within the gastrointestinal tract. This proprietary blend of botanicals includes Tribulus extract, berberine, black walnut powder, barberry extract, artemisinin, along with magnesium and caprylic acid from magnesium caprylate. Research shows that the bioactive constituents in these botanicals possess properties that may help promote a healthy balance of normal gut flora [49-51]. Tribulus contains many bioactive compounds, including saponins and alkaloids, which may have health-promoting properties, such as the support of GI microbial balance [50]. It exhibits properties that are anti-fungal and can bring levels of *Candida* spp. Down [50]. Follow-up testing demonstrated a 35% reduction in *Candida* spp. during a 7-month period (553 to 359 org/g).

Allicillin

To address the dysbiosis that was shown on the patient's GI test, a recommendation of Allicillin was made. This product offers garlic and parsley oils to help support a healthy microbial ecosystem, primarily by inhibiting the growth of unfavorable microbes, such as *Pseudomonas* spp. and *Staphylococcus aureus*. Garlic contains diverse, bioactive compounds that have been studied for a broad range of health benefits. The most well-known compound is allicin. However, the instability of allicin forces it to convert into the following more stable compounds: ajoene, diallyl sulfide (DAS), diallyl disulfide (DADS), and diallyl trisulfide (DATS). Ajoene has been proposed to be more bioactive than allicin by reacting with cysteine residues on proteins such as glutathione reductase, trypanothione reductase, and gastric lipase [52,53]. Ajoene's reaction with the thiol residues on these molecules

is largely responsible for the strong antimicrobial properties garlic offers against several gram-positive and gram-negative bacteria [54]. In addition to gastrointestinal support, the garlic sulfides DAS, DADS, and DATS may also help in supporting metabolic health. A meta-analysis (n = 768) of randomized controlled trials concluded that 0.05 g to 1.5 g of garlic supplementation per day for 12 weeks played a positive and sustained role in normalizing blood glucose, total cholesterol, high-density lipoprotein cholesterol, and low-density lipoprotein cholesterol in the management of type 2 diabetes [55].

Tri-Butyrin Supreme

This supplement features CoreBiome™, a patent-pending form of tributyrin, consisting of three molecules of butyric acid (butyrate) bound to glycerol, delivering butyrate to the colon where it can be fermented into SCFAs. Butyrate supplementation is considered a postbiotic—metabolites of bacteria that benefit the human GI tract. Low butyric acid level was observed on the OAT, suggesting reduced production of short-chain fatty acids and possible dysbiosis affecting the colonic environment. Butyrate is critically important for maintaining gut barrier integrity, modulating inflammation, and providing energy to colonocytes [56].

Total Gut Restoration Kit (TGRK)

Major findings in the GI Spotlight Functional Stool Test were high normal Zonulin, high normal Firmicute, indicating some level of disruption to the gut barrier integrity, along with high *Pseudomonas* spp., a pathogenic gram-negative bacterium, enveloped in lipopolysaccharide (LPS) [26]. The three supplements in the TGRK (MegaSporeBiotic™, MegaPre™, and MegaMucosa™) may have contributed to increasing SCFA production, supporting intestinal barrier function, and regulating a healthy immune response [57,58]. MegaSporeBiotic is the first phase, also called the recondition phase, where 5 *Bacillus* spore-based probiotics (*Bacillus Licheniformis*, *Bacillus Indicus* HU36, *Bacillus Subtilis* HU58, *Bacillus Clausii*, *Bacillus Coagulans*) are introduced. These are meant to survive digestion, colonize, and promote microbial diversity, supporting gut barrier integrity and immune balance. In phase two—the reinforce phase—MegaPre supplies select prebiotic substrates (oligosaccharides) that feed beneficial bacteria while not feeding undesirable ones, helping to reinforce the shifts introduced by the previous phase. In the third phase—the rebuild phase—MegaMucosa helps repair the gut barrier and supports gastrointestinal immune function through a combination of amino acids, immunoglobulins, and bioflavonoids. From 2023 to 2024, immune markers normalized: neutrophils increased by 26.5%, from borderline-low to optimal innate immune activity, lymphocytes decreased by 20.2%, indicating less viral activation, monocytes decreased by 30.4%, showing resolution of chronic inflammatory load, while basophils remained stable, and eosinophils increased slightly, suggesting mild residual stress tone (see Table 2).

Hyperbaric Oxygen Therapy

HBOT was incorporated to support tissue repair, neuroprotection, and overall recovery. By increasing the partial pressure of oxygen in the blood, HBOT supports tissue oxygenation, promoting angiogenesis, collagen synthesis, and cellular metabolism [59]. In the context of post-concussive syndrome and cervical musculoskeletal injury, improved oxygen delivery can reduce hypoxia-related inflammation, support mitochondrial function, and facilitate the repair of damaged neural and connective tissues [59,60]. The patient showed impressive improvements in cognitive performance from the March 2023 baseline CNS vital signs testing to the December 2023 testing, which is discussed further in the following section titled “Neurocognitive and qEEG Findings” as well as in the cognitive performance domain reports (see Figures 1 & 2).

IV Antioxidant Therapy (Vitamin C and Glutathione)

IV administration of vitamin C (12,500 mg) and glutathione (1,000 mg) was utilized to support antioxidant capacity, attenuate inflammation, and enhance tissue repair. Delivering these nutrients intravenously bypasses intestinal absorption limits, allowing for significantly higher plasma concentrations than those achievable through oral intake. Vitamin C transporters in the gastrointestinal tract become saturated at doses between 200–400 mg, limiting systemic bioavailability; however, IV delivery circumvents this threshold and enables therapeutic levels to be reached rapidly [61]. Vitamin C serves as a critical cofactor for prolyl and lysyl hydroxylase—enzymes required for hydroxylation of proline and lysine residues during collagen synthesis—thereby directly contributing to connective tissue remodeling and wound healing [62]. Beyond its structural role, vitamin C neutralizes reactive oxygen species generated during tissue injury, modulates inflammatory pathways, and supports cellular recovery [62–64]. Although human research on IV glutathione remains more limited, emerging data suggest beneficial effects on vascular and metabolic function. In one clinical study, individuals with peripheral arterial disease receiving IV glutathione twice daily for five days demonstrated improved pain-free walking distance and enhanced peripheral circulation compared with placebo [65]. Collectively, this IV antioxidant protocol was intended to reduce oxidative burden, support tissue regeneration, and complement the broader regenerative and mitochondrial restoration strategies implemented in this case.

IV NAD+ Therapy

Nicotinamide adenine dinucleotide (NAD⁺) was incorporated into the patient’s treatment protocol for its central role in mitochondrial function, cellular energy metabolism, and neuroprotection [66–68]. Emerging evidence links suboptimal NAD⁺ levels to neurodegenerative processes, including Alzheimer’s disease and other forms of cognitive decline, where mitochondrial dysfunction and oxidative stress are key pathophysiologic features [67,68]. As an essential coenzyme

in oxidative phosphorylation, NAD⁺ facilitates electron transfer within the mitochondrial respiratory chain, thereby sustaining neuronal energy production and redox balance [69]. In addition to its neuroprotective effects, NAD⁺ plays a vital role in cardiovascular and endothelial health by regulating sirtuin activity, promoting DNA repair, and modulating inflammatory signaling pathways [70]. Through these mechanisms, NAD⁺ repletion supports both cerebral and systemic metabolic recovery, aligning with the comprehensive integrative approach utilized in this case to address mitochondrial dysfunction, cognitive impairment, and vascular dysregulation.

IV Regenerative Therapy (HylaPure and Vitti Pure)

HylaPure Micronized Wharton's Jelly is a specialized formulation derived from Wharton's jelly, the gelatinous connective tissue within the umbilical cord. This tissue is rich in mesenchymal stem cells (MSCs), hyaluronic acid, collagen, and growth factors, making it highly valuable in regenerative applications. HylaPure is classified as a 361-HCT/P product—specifically, a micronized umbilical/placental tissue matrix that is minimally manipulated and cryogenically preserved. It retains the extracellular matrix without the use of DMSO, and is intended for regenerative and reparative purposes. Vitti Pure is a complementary regenerative preparation produced by Vitti-Labs as an alternative to platelet-rich plasma (PRP). It is a growth factor-dense, non-cellular plasma derivative formulated to mimic the regenerative stimulus of PRP but derived from placental and umbilical tissue rather than autologous blood. This stem cell-based injectable offers a cost-effective option, as it does not contain Wharton's jelly or live cells. Each 4.5 cc vial can be administered across multiple joints, tendons, or ligaments. The product delivers non-cellular growth factors from the same immune-privileged birth tissue source as PRP.

The Rhythm Reset 9-Day Program

The Rhythm Reset Program was incorporated as a foundational intervention to re-establish physiologic coherence between the patient's metabolic, neurologic, and circadian systems—areas commonly disrupted in chronic TBI [71-73]. Repetitive head and neck trauma, as seen in many professional athletes, leads to dysregulation of the autonomic nervous system, mitochondrial fatigue, altered sleep-wake cycles, and chronic neuroinflammation. The Rhythm Reset's structured nine-day protocol integrates targeted nutritional support, breath-work, circadian entrainment practices, and lymphatic activation to restore systemic balance. By supporting gut-brain axis repair, reducing neuroinflammatory signaling, and improving oxygen utilization at the cellular level, the program helps reestablish the "rhythmic" synchronization necessary for optimal cognitive recovery and metabolic resilience. This approach aligns with emerging evidence linking circadian rhythm restoration and autonomic recalibration to improved neuronal repair, sleep quality, and neuroendocrine balance in post-concussive individuals. The program's emphasis on nutrient-dense and anti-inflammatory foods, targeted supplementation, guided movement, and structured rest supported mitochondrial efficiency, lymphatic drainage, and gut integrity—three pillars critical

for neurological healing. When combined with IV nutrition, HBOT, chiropractic rehabilitation, and precision nutraceuticals, the Rhythm Reset provided the physiologic groundwork for improved neuronal communication and decreased systemic inflammation.

Peptide Therapy (BPC 157 Oral and GHK-Cu Injections)

The patient supplemented with BPC 157 500mcg 1 capsule daily for 2 months. The inclusion of oral BPC-157 in this patient's comprehensive protocol was based on its multifaceted regenerative and neuroprotective properties relevant to post-concussive and systemic dysfunction [74,75]. BPC-157 has been shown in preclinical studies to promote neuronal repair through upregulation of growth factors such as brain-derived neurotrophic factor (BDNF) and vascular endothelial growth factor (VEGF), while attenuating neuroinflammation via suppression of proinflammatory cytokines and NF- κ B signaling [74-76]. These mechanisms directly address the chronic neuroinflammation, oxidative stress, and neuronal degeneration associated with traumatic brain injury and early cognitive decline that the patient was experiencing. Additionally, oral administration supports gastrointestinal mucosal healing, enhances gut barrier integrity, and modulates the gut-brain axis—key considerations given the patient's significant gastrointestinal dysfunction and its contribution to systemic inflammation [74,77]. BPC-157 provided an integrative therapeutic bridge between the neurologic, metabolic, and structural components of this patient's complex clinical picture, complementing the effects of regenerative IV therapies, HBOT, and targeted nutraceuticals. Another peptide, GHK-Cu (glycyl-L-histidyl-L-lysine-copper), was introduced as part of this patient's regenerative treatment protocol due to its broad reparative and neurorestorative properties [78-80].

This tripeptide-copper complex has been shown to modulate thousands of human genes involved in tissue remodeling, antioxidant defense, and nervous system development [76]. Through suppression of pro-inflammatory mediators such as NF- κ B, TNF- α , and IL-6, and concurrent activation of antioxidant pathways including superoxide dismutase (SOD) and metallothionein, GHK-Cu exerts potent anti-inflammatory and oxidative stress-reducing effects [77,78]. Preclinical studies further demonstrate that GHK-Cu enhances collagen synthesis, angiogenesis, and fibroblast proliferation, key processes for musculoskeletal and connective tissue repair, which are particularly relevant to this patient's cervical enthesopathy and chronic pain [81,82]. Additionally, its ability to improve mitochondrial respiration and neuronal growth supports cognitive and metabolic recovery following traumatic brain injury. When integrated alongside BPC-157, IV regenerative therapy, and HBOT, GHK-Cu serves to promote cellular repair, vascular health, and combat neurodegeneration.

Acupuncture

Acupuncture was incorporated into the patient's integrative protocol to address chronic pain and support tissue healing [83,84]. Needle stimulation promotes pain reduction through the release of endogenous opioids and modulation of central pain pathways, which

is particularly relevant for the patient's cervical enthesopathy and widespread myalgia [85]. In addition, acupuncture enhances local microcirculation, improving the delivery of oxygen and nutrients to injured tissues and supporting connective tissue repair [86,87].

Chiropractic

Chiropractic manipulation therapy was included in the patient's integrative protocol to address musculoskeletal dysfunction and support functional recovery. Spinal adjustments can restore joint alignment and mobility and reduce mechanical stress on muscles and ligaments, which was critical for treatment of the patient's neck pain and chronic myalgia [85]. In addition, chiropractic care may enhance neuromuscular coordination and proprioceptive feedback, supporting postural stability and improving movement patterns following repetitive head and neck trauma [86]. By optimizing joint and soft tissue function, chiropractic interventions complement physical therapy, regenerative peptide therapy, and nutraceutical support, facilitating pain reduction, tissue repair, and overall rehabilitation.

CNS Vital Signs

CNS Vital Signs was chosen as the primary cognitive assessment tool because it provides a validated, quantitative profile of multiple brain functions that are often disrupted in individuals with repetitive sub-concussive exposure and metabolic dysfunction. The platform measures domains such as memory, processing speed, executive function, and attention with high test-retest reliability, allowing objective tracking of progress over time. Unlike standard symptom questionnaires, CNS Vital Signs captures subtle neurocognitive recovery that parallels physiologic changes seen on qEEG and metabolic testing. In this case, it documented the transition from slowed processing and cognitive fatigue to normalized or above-average performance, confirming restoration of network efficiency and functional brain resilience.

Neurocognitive and qEEG Findings

Six serial assessments from March 2023 through December 2023 documented steady cognitive improvement (Figures 1 & 2 and Table 5). Neurocognitive Index (+19.5), Executive Function (+21%), Psychomotor speed (+36%), Processing Speed (+29.8%), Composite Memory (+8.3%), and Cognitive flexibility (+20.7%) progressed from low-average to expected or above-average norms, coinciding with the phases of gut restoration and metabolic therapy. When the patient first arrived, his brain maps and cognitive tests showed a system running in "low gear." He could think and remember, but mental tasks required extra effort. The qEEG confirmed slower healing rhythms (theta and delta) and weak communication between key control networks, patterns that reflect inflammation and metabolic fatigue, much like his lab markers of anemia and immune under-drive.

The biggest change came once his metabolic and gut-repair measures had been in place long enough to lower inflammation and improve cellular energy. That shift, together with oxygen therapies,

NAD+ support, and targeted brain training, allowed the neural networks to resynchronize. In short, the "hardware" of his brain finally had the fuel and stability it needed to run at full speed again. As gut and metabolic therapies normalized nutrition status, oxygenation, and inflammatory control, those brain-wave patterns normalized. CNS Vital Signs testing confirmed faster processing, sharper memory, and sustained attention returning to normal or better.

By early 2025, the Default Mode and Executive Networks were back in sync: slow, drowsy waves quieted, while faster beta activity strengthened and became more coordinated (Figure 3). The frontal and cingulate regions, centers of attention, self-control, and consciousness, reconnected fully, explaining his renewed clarity, energy, and cognitive resilience. Today, his test scores and brain activity patterns resemble those of a healthy, well-regulated nervous system. He can sustain focus, process information rapidly, and recover from stress or fatigue without setbacks. The data show not just symptom relief but restoration of brain efficiency—the ability to switch smoothly between rest, focus, and recovery, which is the foundation of lasting cognitive and emotional resilience.

Fit3D Body Scans

Fit3D scanning began in July 2024, one year into treatment. The first scan reported some body mass improvement, at 296.1 lbs. (-7.5%) relative to the original 320 lb. baseline we measured on the date of his first visit. Across four scans (July 2024 → March 2025), weight decreased into the low-220-lb range (-15.9%), BMI shifted from 37.9 to 32, and body-fat percentage improved (-22%) with some lean-mass preservation (-7.2%), and an overall significant improvement in Body Shape Rating (+78.6%). Basal Metabolic Rate adjusted from 2320 kcal to 2102 kcal (-9.4%) (see Table 6).

Conclusion

This case demonstrates how a phased, multiomics-guided rehabilitation program can improve neurologic and metabolic resilience in a retired professional athlete with a history of repetitive sub-concussive injury. Integration of gut restoration, oxygen-based therapies, metabolic repair, and regenerative biologic interventions resulted in measurable improvement across cognitive, electrophysiologic, hematologic, and body-composition domains. Quantitative EEG provided one of the most striking indicators of recovery, showing a 61-76% reduction in network abnormalities across the Default Mode, Central Executive, and Salience Networks (Figure 3). These objective brain improvements closely paralleled the second series of NAD+ infusions, which enhanced mitochondrial redox signaling and promoted neuronal and neurovascular repair. Together, these interventions reestablished cortical efficiency, coherence, and functional connectivity that had been disrupted by years of microtrauma and metabolic strain. Serial CNS Vital Signs assessments confirmed the clinical significance of these findings, documenting recovery of processing speed, sustained attention, memory, and executive function from below-average to ex-

pected or above-average levels. Metabolic testing and Fit3D analyses supported these outcomes, showing improved mitochondrial performance, immune regulation, and over 100 pounds of healthy weight reduction with preservation of lean mass.

The convergence of these outcomes demonstrates that clinically meaningful neuroregeneration can occur when oxygen dynamics, mitochondrial metabolism, and systemic inflammation are simultaneously corrected. This case supports the concept that integrating multiomics assessment with targeted oxygen and regenerative therapies can achieve durable neurologic and metabolic resilience, offering a reproducible model for addressing post-concussive syndrome and chronic TBI.

Patient Consent

Written informed consent was obtained from the patient for publication of this case report. Identifying information has been removed or minimized to protect patient confidentiality.

Ethics Approval

This case report describes retrospective clinical observations from routine patient care. Institutional review board approval was not required for this single-patient case report, however, the patient provided written informed consent for publication of this case report.

Author Contributions

Justin Dearing contributed to patient care, clinical data collection, case interpretation, and manuscript review. Melanie Luther contributed to literature review, clinical writing, and manuscript preparation. Seamus Kelly added further input on scientific deep dives and clinical writing. Oscar Coetzee contributed to clinical interpretation, manuscript development, revision, and correspondence with the journal. All authors reviewed and approved the final manuscript.

Funding

The authors received no financial support for the research, authorship, or publication of this case report. All treatments were paid for by patient.

Acknowledgement

The authors acknowledge the healthcare professionals involved in the patient's care and collection of clinical records used in this case report. The authors are especially grateful to the team at The Dearing Clinic (Brentwood, Tennessee), including Justin Dearing, DC, DACNB, FIAMA, FAARM; Lauren Dearing, ACNP; Catherine Jones, FNP; and Charles Bell, DO, for their clinical expertise, patient care, and assistance with clinical documentation. The authors also acknowledge Richard J. McAlister, DC, BCN, qEEG-D, for his expertise in quantitative electroencephalography (qEEG) acquisition and analysis. The authors thank Seamus Kelly, MS candidate, for assistance with manuscript preparation, formatting, editorial review, and table standardization.

References

- Zimmerman KA, Kim J, Karton C, Lochhead, D J Sharp, et al. (2021) Player position in American football influences the magnitude of mechanical strains produced in the location of chronic traumatic encephalopathy pathology: A computational modelling study. *J Biomech* 118: 110256.
- George AK, Behera J, Homme RP, Tyagi N, Tyagi SC, et al. (2021) Rebuilding microbiome for mitigating traumatic brain injury: Importance of restructuring the gut-microbiome-brain axis. *Mol Neurobiol* 58(8): 3614-3627.
- Heston MB, Hanslik KL, Zarbock KR, Sandra J Harding, Nancy J Davenport-Sis, et al. (2023) Gut inflammation associated with age and Alzheimer's disease pathology: A human cohort study. *Sci Rep* 13(1):18924.
- Lin Y, Hou C, Wang C, Rui Chen, Yunzhong Zhu, et al. (2024) Research progress on digestive disorders following traumatic brain injury. *Front Immunol* 15:1524495.
- Clarke GJB, Skandsen T, Zetterberg H, Turid Follestad, Cathrine Elisabeth Einarsen, et al. (2024) Longitudinal associations between persistent post-concussion symptoms and blood biomarkers of inflammation and CNS injury after mild traumatic brain injury. *J Neurotrauma* 41(7-8): 862-878.
- Gard A, Vedung F, Piehl F, Mohsen Khademi, Maria Portonova Wernersson, et al. (2023) Cerebrospinal fluid levels of neuroinflammatory biomarkers are increased in athletes with persistent post-concussive symptoms following sports-related concussion. *J Neuroinflammation* 20(1): 189.
- Ryan T, Nagle S, Daly E, Pearce AJ, Ryan L, et al. (2023) A potential role exists for nutritional interventions in the chronic phase of mild traumatic brain injury, concussion and sports-related concussion: A systematic review. *Nutrients* 15(17): 3726.
- Finnegan E, Daly E, Pearce AJ, Ryan L (2022) Nutritional interventions to support acute mTBI recovery. *Front Nutr* 9: 977728.
- Haider MN, Bezherano I, Wertheimer A, Akas H Siddiqui, Emily C Horn, et al. (2021) Exercise for sport-related concussion and persistent post-concussive symptoms. *Sports Health* 13(2): 154-160.
- Art K, Ridenour C, Durbin S, Bauer M, Hassen-Miller A, et al. (2023) The effectiveness of physical therapy interventions for athletes post-concussion: A systematic review. *Int J Sports Phys Ther* 18(1): 26-38.
- Rosales C (2018) Neutrophil: A cell with many roles in inflammation or several cell types?. *Front Physiol* 9: 113.
- Zhang F, Xia Y, Su J, Fushi Quan, Hengzong Zhou, et al. (2024) Neutrophil diversity and function in health and disease. *Signal Transduct Target Ther* 9(1): 343.
- Chi H, Pepper M, Thomas PG (2024) Principles and therapeutic applications of adaptive immunity. *Cell* 187(9): 2052-2078.
- LaRosa DF, Orange JS (2008) Lymphocytes. *J Allergy Clin Immunol* 121(2 Suppl): S364-S369.
- Weinstock A, Fisher EA (2019) Methods to study monocyte and macrophage trafficking in atherosclerosis progression and resolution. *Methods Mol Biol* 1951: 153-165.
- Yang J, Zhang L, Yu C, Yang XF, Wang H, et al. (2014) Monocyte and macrophage differentiation: Circulation inflammatory monocyte as biomarker for inflammatory diseases. *Biomark Res* 2(1): 1.
- Buonacera A, Stancanelli B, Colaci M, Malatino L (2022) Neutrophil-to-lymphocyte ratio: An emerging marker of the relationships between the immune system and diseases. *Int J Mol Sci* 23(7): 3636.
- Li Y, Xue Z, Dong X, Qian Liu, Zhe Liu, et al. (2020) Mitochondrial dysfunction and oxidative stress in bone marrow stromal cells induced by dauno-

- rubicin leads to DNA damage in hematopoietic cells. *Free Radic Biol Med* 146: 211-221.
19. Hampton K, Polski-Delva A, Hellmich C, Rushworth SA (2025) Linking mitochondria, fatty acids, and hematopoietic stem cell expansion during infection: Implications for aging and metabolic diseases. *Stem Cells* 43(11): sxaf053.
 20. Wang J, Shen J, Chen D, Binyou Liao, Xi Chen, et al. (2024) Secretory IgA reduced the ergosterol contents of *Candida albicans* to repress its hyphal growth and virulence. *Appl Microbiol Biotechnol* 108(1): 244.
 21. Moreno-Sabater A, Sterlin D, Imamovic L, et al. (2023) Intestinal *Candida albicans* overgrowth in IgA deficiency. *J Allergy Clin Immunol* 152(3): 748-759.e3.
 22. Zhao J, Bi W, Xiao S, Xiaofeng Cheng, Jiawei Zhang, et al. (2019) Neuroinflammation induced by lipopolysaccharide causes cognitive impairment in mice. *Sci Rep* 9(1): 5790.
 23. Nieminen MT, Salaspuro M (2018) Local acetaldehyde—an essential role in alcohol-related upper gastrointestinal tract carcinogenesis. *Cancers (Basel)* 10(1): 11.
 24. Wan Z, Zheng J, Zhu Z, Lan Sang, Jinwei Zhu, et al. (2022) Intermediate role of gut microbiota in vitamin B nutrition and its influences on human health. *Front Nutr* 9: 1031502.
 25. Meir Z, Osherov N (2018) Vitamin biosynthesis as an antifungal target. *J Fungi (Basel)* 4(2): 72.
 26. Huszczyński SM, Lam JS, Khursigara CM (2019) The role of *Pseudomonas aeruginosa* lipopolysaccharide in bacterial pathogenesis and physiology. *Pathogens* 9(1): 6.
 27. Skrzypczak-Wiercioch A, Sałat K (2022) Lipopolysaccharide-induced model of neuroinflammation: Mechanisms of action, research application and future directions for its use. *Molecules* 27(17): 5481.
 28. Sannasiddappa TH, Lund PA, Clarke SR (2017) *In vitro* antibacterial activity of unconjugated and conjugated bile salts on *Staphylococcus aureus*. *Front Microbiol* 8: 1581.
 29. Liu A, Garrett S, Hong W, Zhang J (2024) *Staphylococcus aureus* infections and human intestinal microbiota. *Pathogens* 13(4): 276.
 30. Ríos-Covián D, Ruas-Madiedo P, Margolles A, Gueimonde M, de Los Reyes-Gavilán CG, et al. (2016) Intestinal short chain fatty acids and their link with diet and human health. *Front Microbiol* 7: 185.
 31. Hodgkinson K, El Abbar F, Dobranowski P, Juliana Manoogian, James Butcher, et al. (2023) Butyrate's role in human health and the current progress towards its clinical application to treat gastrointestinal disease. *Clin Nutr* 42(2): 61-75.
 32. Corder EH, Saunders AM, Strittmatter WJ, D E Schmechel, P C Gaskell, et al. (1993) Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261(5123): 921-923.
 33. Reiman EM, Webster JA, Myers AJ, John Hardy, Travis Dunckley, et al. (2007) GAB2 alleles modify Alzheimer's risk in APOE ϵ 4 carriers. *Neuron* 54(5): 713-720.
 34. Liu D, Zhao H, Zhang Y, Hu J, Xu H, et al. (2025) Efficacy and safety of berberine on the components of metabolic syndrome: A systematic review and meta-analysis of randomized placebo-controlled trials. *Front Pharmacol* 16:1572197.
 35. Yang L, Zhu W, Zhang X, Zhou X, Wu W, et al. (2023) Efficacy and safety of berberine for several cardiovascular diseases: A systematic review and meta-analysis of randomized controlled trials. *Phytomedicine* 112: 154716.
 36. Wang H, Zhu C, Ying Y, Luo L, Huang D, et al. (2018) Metformin and berberine, two versatile drugs in treatment of common metabolic diseases. *Oncotarget* 9(11): 10135-10146.
 37. Araj-Khodaei M, Ayati MH, Azizi Zeinalhajlou A, Tannaz Novinbahador, Mehdi Yousefi, et al. (2024) Berberine-induced glucagon-like peptide-1 and its mechanism for controlling type 2 diabetes mellitus: A comprehensive pathway review. *Arch Physiol Biochem* 130(6): 678-685.
 38. Yang WL, Zhang CY, Ji WY, Fang-Yuan Yang, Lin Zhang, et al. (2024) Berberine metabolites stimulate GLP-1 secretion by alleviating oxidative stress and mitochondrial dysfunction. *Am J Chin Med* 52(1): 253-274.
 39. Salehi B, Yilmaz BY, Antika G, Tugba Boyunegmez Tumer, Mohamad Fawzi Mahomoodally, et al. (2019) Insights on the use of α -lipoic acid for therapeutic purposes. *Biomolecules* 9(8): 356.
 40. Van den Abbeele P, Kunkler CN, Poppe J, Rose A, van Hengel IAJ, et al. (2024) Serum-derived bovine immunoglobulin promotes barrier integrity and lowers inflammation for 24 human adults *ex vivo*. *Nutrients* 16(11): 1585.
 41. Li Y, Jin L, Chen T (2020) The effects of secretory IgA in the mucosal immune system. *Biomed Res Int* 2020: 2032057.
 42. Dib K (2025) Histamine promotes *Pseudomonas aeruginosa* biofilm formation and renders *P. aeruginosa* biofilms more resistant to gentamicin and azithromycin. *Med Princ Pract* 34(3): 238-249.
 43. Touaitia R, Mairi A, Ibrahim NA, Basher NS, Idres T, et al. (2025) *Staphylococcus aureus*: A review of the pathogenesis and virulence mechanisms. *Antibiotics (Basel)* 14(5): 470.
 44. Tsilochristou O, du Toit G, Sayre PH, Graham Roberts, Kaitie Lawson, et al. (2019) Association of *Staphylococcus aureus* colonization with food allergy occurs independently of eczema severity. *J Allergy Clin Immunol* 144(2): 494-503.
 45. Fasano A (2020) All disease begins in the (leaky) gut: Role of zonulin-mediated gut permeability in the pathogenesis of some chronic inflammatory diseases. *F1000Res* 9: F1000 Faculty Rev-69.
 46. Mahmood A, FitzGerald AJ, Marchbank T, D Murray, S Ghosh, et al. (2007) Zinc carnosine, a health food supplement that stabilises small bowel integrity and stimulates gut repair processes. *Gut* 56(2): 168-175.
 47. Hewlings S, Kalman D (2020) A review of zinc-L-carnosine and its positive effects on oral mucositis, taste disorders, and gastrointestinal disorders. *Nutrients* 12(3): 665.
 48. Efthymakis K, Neri M (2022) The role of zinc L-carnosine in the prevention and treatment of gastrointestinal mucosal disease in humans: A review. *Clin Res Hepatol Gastroenterol* 46(7): 101954.
 49. Ho KV, Lei Z, Sumner LW, Mark V Coggeshall, Hsin-Yeh Hsieh, et al. (2018) Identifying antibacterial compounds in black walnuts (*Juglans nigra*) using a metabolomics approach. *Metabolites* 8(4): 58.
 50. Al-Bayati FA, Al-Mola HF (2008) Antibacterial and antifungal activities of different parts of *Tribulus terrestris* L. growing in Iraq. *J Zhejiang Univ Sci B* 9(2): 154-159.
 51. Abiri R, Silva ALM, de Mesquita LSS, José Wilson Carvalho de Mesquita, Narges Atabaki, et al. (2018) Towards a better understanding of *Artemisia vulgaris*: Botany, phytochemistry, pharmacological and biotechnological potential. *Food Res Int* 109: 403-415.
 52. Bouyahya A, Kamari F, El Omari N, et al. (2025) Unlocking the natural chemical sources, nutritional properties, biological effects, and molecular actions of ajoene. *J Funct Foods* 127: 106714.
 53. Rose P, Moore PK, Whiteman M, Zhu YZ (2019) An appraisal of develop-

- ments in *Allium sulfur* chemistry: Expanding the pharmacopeia of garlic. *Molecules* 24(21): 4006.
54. Bhatwalkar SB, Mondal R, Krishna SBN, Adam JK, Govender P, et al. (2021) Antibacterial properties of organosulfur compounds of garlic (*Allium sativum*). *Front Microbiol* 12: 613077.
 55. Wang J, Zhang X, Lan H, Wang W (2017) Effect of garlic supplement in the management of type 2 diabetes mellitus (T2DM): A meta-analysis of randomized controlled trials. *Food Nutr Res* 61(1): 1377571.
 56. Canani RB, Costanzo MD, Leone L, Pedata M, Meli R, et al. (2011) Potential beneficial effects of butyrate in intestinal and extraintestinal diseases. *World J Gastroenterol* 17(12): 1519-1528.
 57. Marzorati M, Bubeck S, Bayne T, Krishnan K, Giusto M, et al. (2022) Effects of combined prebiotic, probiotic, IgG and amino acid supplementation on the gut microbiome of patients with inflammatory bowel disease. *Future Microbiol* 17: 1307-1324.
 58. Catanean A, Neag MA, Krishnan K, Corina Ioana Bocsan, Raluca Maria Pop, et al. (2020) Probiotic *Bacillus* spores together with amino acids and immunoglobulins exert protective effects on a rat model of ulcerative colitis. *Nutrients* 12(12): 3607.
 59. Schottlender N, Gottfried I, Ashery U (2021) Hyperbaric oxygen treatment: Effects on mitochondrial function and oxidative stress. *Biomolecules* 11(12): 1827.
 60. Shabi Shlifer A, Suzin G, Shorer R, Erez Lang, Shachar Finci, et al. (2025) Hyperbaric oxygen therapy improves post-concussion symptoms in adults with childhood traumatic brain injury: A retrospective cohort study. *Front Neurol* 16: 1641033.
 61. Lykkesfeldt J, Carr AC, Tveden-Nyborg P (2025) The pharmacology of vitamin C. *Pharmacol Rev* 77(2): 100043.
 62. DePhillipo NN, Aman ZS, Kennedy MI, Begley JP, Moatshe G, et al. (2018) Efficacy of vitamin C supplementation on collagen synthesis and oxidative stress after musculoskeletal injuries: A systematic review. *Orthop J Sports Med* 6(10): 2325967118804544.
 63. Bechara N, Flood VM, Gunton JE (2022) A systematic review on the role of vitamin C in tissue healing. *Antioxidants (Basel)* 11(8): 1605.
 64. Xu C, Yi T, Tan S, Yu Hu, Junpeng Ma, et al. (2023) Association of oral or intravenous vitamin C supplementation with mortality: A systematic review and meta-analysis. *Nutrients* 15(8): 1848.
 65. Arosio E, De Marchi S, Zannoni M, Prior M, Lechi A, et al. (2002) Effect of glutathione infusion on leg arterial circulation, cutaneous microcirculation, and pain-free walking distance in patients with peripheral obstructive arterial disease: A randomized, double-blind, placebo-controlled trial. *Mayo Clin Proc* 77(8): 754-759.
 66. Lautrup S, Sinclair DA, Mattson MP, Fang EF (2019) NAD⁺ in brain aging and neurodegenerative disorders. *Cell Metab* 30(4): 630-655.
 67. Clement J, Wong M, Poljak A, Sachdev P, Braid N (2019) The plasma NAD⁺ metabolome is dysregulated in "normal" aging. *Rejuvenation Res* 22(2): 121-130.
 68. Grant R, Berg J, Mestayer R, Nady Braid, James Bennett, et al. (2019) A pilot study investigating changes in the human plasma and urine NAD⁺ metabolome during a 6-hour intravenous infusion of NAD. *Front Aging Neurosci* 11: 257.
 69. Cantó C, Menzies KJ, Auwerx J (2015) NAD⁺ metabolism and the control of energy homeostasis: A balancing act between mitochondria and the nucleus. *Cell Metab* 22(1): 31-53.
 70. Csiszar A, Tarantini S, Yabluchanskiy A, Tamas Kiss, Eszter Farkas, et al. (2019) Role of endothelial NAD⁺ deficiency in age-related vascular dysfunction. *Am J Physiol Heart Circ Physiol* 316(6): H1253-H1266.
 71. Kuo LT, Lu HY, Chen YH (2024) Traumatic brain injury-induced disruption of the circadian clock. *J Mol Med (Berl)* 102(3): 403-414.
 72. Oft HC, Simon DW, Sun D (2024) New insights into metabolism dysregulation after TBI. *J Neuroinflammation* 21(1): 184.
 73. Haria JM, Singh NK, Kumar J, Jain SK, Pamidimarri D, et al. (2025) Metabolic and endocrine dysfunctions in traumatic brain injury: Implications for cognitive recovery and therapeutic strategies. *Behav Brain Res* 493: 115697.
 74. Vukojevic J, Milavić M, Perović D, Spomenko Ilić, Andrea Zemba Čilić, et al. (2022) Pentadecapeptide BPC 157 and the central nervous system. *Neural Regen Res* 17(3): 482-487.
 75. Klicek R, Kolenc D, Suran J, D Drmic, L Brcic, et al. (2013) Stable gastric pentadecapeptide BPC 157 heals cysteamine-colitis and colon-colon-anastomosis and counteracts cuprizone brain injuries and motor disability. *J Physiol Pharmacol* 64(5): 597-612.
 76. Perovic D, Kolenc D, Bilic V, Domagoj Drmic, Esmat Elabjer, et al. (2019) Stable gastric pentadecapeptide BPC 157 can improve the healing course of spinal cord injury and lead to functional recovery in rats. *J Orthop Surg Res* 14(1): 199.
 77. Tohyama Y, Sikirić P, Diksic M (2004) Effects of pentadecapeptide BPC157 on regional serotonin synthesis in the rat brain: Alpha-methyl-L-tryptophan autoradiographic measurements. *Life Sci* 76(3): 345-357.
 78. Pickart L, Vasquez-Soltero JM, Margolina A (2015) GHK peptide as a natural modulator of multiple cellular pathways in skin regeneration. *Biomed Res Int* 2015: 648108.
 79. Pickart L, Vasquez-Soltero JM, Margolina A (2017) The effect of the human peptide GHK on gene expression relevant to nervous system function and cognitive decline. *Brain Sci* 7(2): 20.
 80. Siméon A, Emonard H, Hornebeck W, Maquart FX (2000) The tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ stimulates matrix metalloproteinase-2 expression by fibroblast cultures. *Life Sci* 67(18): 2257-2265.
 81. Jiang F, Wu Y, Liu Z, Hong M, Huang Y, et al. (2023) Synergy of GHK-Cu and hyaluronic acid on collagen IV upregulation via fibroblast and ex vivo skin tests. *J Cosmet Dermatol* 22(9): 2598-2604.
 82. Vickers AJ, Cronin AM, Maschino AC, George Lewith, Hugh MacPherson, et al. (2012) Acupuncture for chronic pain: Individual patient data meta-analysis. *Arch Intern Med* 172(19): 1444-1453.
 83. Xue L, Yan Y, Chen Z, Hong Y (2025) Acupuncture and massage therapy enhance postoperative recovery and quality of life in orthopedic patients. *Medicine (Baltimore)* 104(37): e44314.
 84. Gao P, Gao XI, Fu T, Xu D, Wen Q, et al. (2015) Acupuncture: Emerging evidence for its use as an analgesic (review). *Exp Ther Med* 9(5): 1577-1581.
 85. Valentini J, Sigl M, Dunckel C, Krisam J, Amendt K, et al. (2024) Can acupuncture increase microcirculation in peripheral artery disease and diabetic foot syndrome? A pilot study. *Front Med (Lausanne)* 11: 1371056.
 86. Komori M, Takada K, Tomizawa Y, Keiko Nishiyama, Izumi Kondo, et al. (2009) Microcirculatory responses to acupuncture stimulation and phototherapy. *Anesth Analg* 108(2): 635-640.
 87. Haneline MT, Cooperstein R (2009) Chiropractic care for patients with acute neck pain: Results of a pragmatic practice-based feasibility study. *J Chiropr Med* 8(4): 143-155.
 88. Holt KR, Haavik H, Lee ACL, Murphy B, Elley CR, et al. (2016) Effectiveness of chiropractic care to improve sensorimotor function associated with falls risk in older people: A randomized controlled trial. *J Manipulative Physiol Ther* 39(4): 267-278.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2026.66.010291](https://doi.org/10.26717/BJSTR.2026.66.010291)

Justin Dearing . Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>