

Use of Primer Design for the Detection of *Ehrlichia Canis* by Conventional Polymerase Chain Reaction

Echeverría-Barría G and Navarro-Venegas C*

Veterinary and Animal Sciences Faculty. FAVET. University of Chile

*Corresponding author: Navarro Venegas C, email ID: carnaven444@gmail.com Veterinary and Animal Sciences Faculty. FAVET. University of Chile

ARTICLE INFO

Received:  July 11, 2026

Published:  July 22, 2026

Citation: Echeverría-Barría G and Navarro-Venegas C. Use of Primer Design for the Detection of *Ehrlichia Canis* by Conventional Polymerase Chain Reaction. Biomed J Sci & Tech Res 66(2)-2026. BJSTR. MS.ID.010315.

ABSTRACT

Canine Ehrlichiosis is an infectious disease with high morbidity and mortality among domestic dogs and other members of the *Canidae* family. It is found in countries throughout the world. It is transmitted by ticks and is caused by *Ehrlichia* spp., an obligate intracellular gram-negative bacterium. It has tropism for lymphocytes, monocytes, and granulocytes. It is endemic in tropical and subtropical regions and is increasingly spreading to areas with temperate climates. This may be due to both climatic and environmental changes that affect the distribution of ticks and travel made with pets, allowing the arrival of this disease to non-endemic areas. However, an increase in diagnosis is also contemplated due to an improvement in diagnostic techniques. Depending on the phase of the disease (acute, subclinical or chronic), it can produce signs such as high fever, anorexia, corneal opacity, weight loss, epistaxis, vomiting, diarrhea, anemia, thrombocytopenia. Diagnostic techniques include hematology, serology, bacterial isolation, and molecular techniques. Among these, the polymerase chain reaction (PCR) method offers a definitive diagnosis due to its greater sensitivity and specificity, as it uses genus-specific and species-specific primers. Diagnosis is often difficult because coinfection with other tick-borne pathogens is common, complicating pathogenesis, clinical manifestations, diagnosis, and treatment. Therefore, it is important to choose a diagnostic technique with greater specificity and sensitivity. This disease is a problem in veterinary medicine for the reasons mentioned above, as well as for the zoonotic potential of these agents and their potential impact on human health. Thus, this work proposes a molecular protocol with *in silico*-designed primers for the *trp36* gene of *Ehrlichia canis* as a diagnostic method.

Abbreviations: PCR: Polymerase Chain Reaction; LPS: Lipopolysaccharides; ABS: Anti-Ehrlichia Antibodies; ELISA: Enzyme-Linked Immunosorbent Assay; IIF: Indirect Immunofluorescence

Background

Pathogenesis and Pathology

Ehrlichia spp. are obligate intracellular Gram-negative bacteria that infect monocytes, granulocytes, and platelets [1]. They do not synthesize lipopolysaccharides (LPS) and produce little peptidoglycan, which diminishes the host's immune response and provides flexibility and plasticity, facilitating intravascular circulation in infected leukocytes [2]. In its infective form, it enters the cell by endocytosis, where it matures and multiplies, forming morulae with 20–40 infective cells. These morulae are released after a few days, initiating a new infectious cycle [2,3]. It spreads via the blood or lymphatic system within infected cells, reaching organs [4], such as the liver, spleen, bone marrow and lymph nodes where they multiply (incubation period of 8-20 days) [3,5].

Transmission

Rhipicephalus sanguineus ticks (*R. sanguineus*) are the main vector of *E. canis* transmission (hematophagous arthropods of the *Ixodidae* family, *Arachnida* class). *R. sanguineus* is the most widely distributed tick species in both tropical and subtropical areas [1], which are most susceptible to *E. canis* infestation [6,7]. Its prevalence is high year-round, while in temperate climates it only increases in summer [8]. There is no assessment of infestation by age, sex, or breed [1]. The tick becomes infected from larvae to nymphs and from nymphs to adults, or during the infection, after ingesting blood from an infected dog. Once infected, they can inoculate the bacteria into a new host [3]. *E. canis* DNA has been detected in fox fleas (*Vulpes vulpes*) and in small mammals that do not belong to the *Canidae* family, suggesting another transmission cycle that may include other arthropods [3], blood transfusions and accidents with objects contaminated with infected blood as serious potential forms of transmission [3,4,9].

Clinical Signs

They vary depending on the phase: acute, subclinical, or chronic [2,10], the level of virulence of the strain, the pathogen dose, the host immune system, and breed (German shepherds have higher morbidity and mortality) [3,4,9], and coinfection with other pathogens [3]. Common symptoms at different stages are nonspecific, the most common being fever, anorexia, corneal opacity, weight loss, epistaxis, vomiting, diarrhea, anemia, and thrombocytopenia (occurring in more than 90% of infected individuals). In severe cases, hepatomegaly, splenomegaly, nervous system damage (meningitis, ataxia, and seizure symptoms), hemorrhages, and significant weight loss may occur [2,3,9]. Respiratory signs such as dyspnea, seropurulent secretions in the nasal passages and conjunctival sacs, and even interstitial pneumonia are also present. In addition to lymphadenomegaly, there is a tendency to bleed (dermal petechiae and/or ecchymosis) due to thrombocytopenia, which is attributed to increased platelet consumption due to vasculitis, increased splenic sequestration and immunological destruction or decreased platelet half-life [3,9]. In the acute phase, unilateral or bilateral epistaxis, extravasculations at injection sites and in the anterior chamber of the eyes, blood in urine and feces have been described. These signs remit in 1-4 weeks, but if not treated, they can enter a subclinical phase, for months [10] and even up to 10 years without eliminating it and without clinical signs (asymptomatic) with mild thrombocytopenia. If the infection passes to the subclinical phase, it becomes chronic [3,9], a phase that can present in a mild or severe form, where we find pancytopenia, hemorrhagic diathesis, bone marrow aplasia (myelosuppression) [3], marked splenomegaly, glomerulonephritis, kidney failure, interstitial pneumonitis, anterior uveitis and meningitis [9]. Death is usually due to extensive hemorrhages or secondary infections [3]. Other alterations are: hypoalbuminemia, hyperglobulinemia and hypergammaglobulinemia, increased alkaline phosphatase, increased alanine aminotransferase and increased urea and creatinine concentrations, non-regenerative normocytic anemia and a level of leukopenia with monocytosis [3].

Diagnosis

Microscopic Visualization

Giemsa staining of peripheral blood smears can visualize morulae in monocytes using microscopy during the acute phase. It is inexpensive and rapid, but has low sensitivity [2]. White-cap smears, made from concentrated leukocytes, are also available to detect morulae, which has greater sensitivity. False negatives can occur, resulting in a diagnostic disadvantage and requiring highly trained personnel [2,3].

Serology

Serology involves the search for anti-*Ehrlichia* antibodies (ABs) using methods such as enzyme-linked immunosorbent assay (ELISA) and indirect immunofluorescence (IIF). The latter is considered the gold standard, but the ABs detected are not specific and cross-react

with other *Ehrlichia* spp. agents [2,3,10]. Anti-*E. canis* IgG antibody detection tests are available (SNAP 3Dx® and SNAP 4Dx® from IDEXX laboratories) [2,3] with a sensitivity and specificity of 97.8% and 92.3%, respectively [2]. The IgG antibody response can persist for weeks or even months after infection, so we can have false positives [9].

Molecular Diagnosis

Polymerase Chain Reaction (PCR) has proven to be an effective tool for the molecular detection of *E. canis* [1]. It is genus and species-specific and allows for diagnosis in acute and chronic phases [2], confirming active *E. canis* infection before seroconversion by AC occurs [3]. There are methods based on the amplification of coding genes with highly conserved regions, achieving 100% specificity [2]. For *E. canis*, the 16S rRNA gene and the genes encoding the p28 and p30 proteins have been used as target genes, the latter being the most sensitive [2, 3]. PCR samples can be blood, aspirates, or biopsies from organs such as lymph nodes, liver, bone marrow, or spleen [9]. The latter is more sensitive to assessing *E. canis* clearance. The disadvantage of PCR is that it is easily contaminated, so the use of controls at each step is recommended [3].

Treatment

Doxycycline is the antibiotic of choice and should be administered at a dose of 10 mg/kg every 24 hours for 3-4 weeks [3,9,11], as it eliminated *E. canis* infection in most dogs in studies [12]. In chronic cases, supportive therapy may be required [9]. The phase of the disease during which treatment is initiated influences the final results; patients with acute or subclinical phases maintained negative PCR results when clinical parameters improved, but those treated in the chronic phase remained intermittently positive for *E. canis* [3].

Prevention

Prevention of ehrlichiosis and other tick-borne diseases is achieved by eliminating ticks from the environment and preventing re-infestation. This includes prophylactic management of the pet's environment [3,4]. Chemical control and infestation prevention with acaricides, collars, sprays, oral products, injectables, and chewable tablets are available [13]. If you live in areas with a high tick population, pets should avoid contact with wildlife. Environmental care should be provided in areas where pets are kept, such as areas surrounding the home, generally through the application of tick-repellent solutions [3]. Maintaining a balanced diet produces a better immune response. It does not prevent *E. canis* infection, but it does prevent complications [4]. Progress has been made in developing a vaccine based on an attenuated strain of *E. canis*, but it has not yet been successful [14].

Materials and Methods

This project can be carried out in any laboratory in the Third World, thanks to Kary Mullis's brilliant idea [15] and the availability

of various bio-tools that are still free on the Internet. For primer design, the Genbank database will be used to search for the trp36 gene sequence, and the Invitrogen Oligoperfect Design program will be used to select the primer.

Design Of Primers from the Ehrlichia Canis Trp36 Gene

From the Genbank® database, we will access the *E. canis* trp36 gene sequences, with a total of 65 sequences. From these, we will select those containing at least 700 base pairs. Within these, we will search for a consensus sequence using the Oligoperfect Design® program, where a ranking of optimal matches will be provided.

For primer selection, the following criteria will be used:

- Primer length: between 18-25 nucleotides
- GC content (%GC): 50%
- Melting temperature (Tm): Between 50-60°
- Absence of secondary structures: the primers must not form Dimers (self- or heterodimers) or hairpin structures.
- Specificity: region of the trp36 gene

Propose a Protocol for Conventional PCR Detection of E. Canis

The proposal for specific conditions for molecular detection considers the use of the assays designed in the previous point, in conjunction with the generic conditions for any PCR reaction, such as:

- PCR reaction mix.** The mixture to be formed in each DNA-like PCR tube (from a suspect sample, the vaccine, a positive control, or a negative control), the designed primers, and a Master Mix solution (containing Taq Polymerase, nucleotides ([A, T, C, G]), and Mg²⁺). The PCR tube, referred to as the reagent control, will not contain DNA but nuclease-free water and all the other reagents already mentioned.
- PCR reaction.** The PCR reaction uses a protocol compatible with conventional PCR, which generally includes three steps: denaturation at 94°C for 30 seconds; annealing at a temperature defined by the primer design, initially (Tm-5)°C for 30 seconds; and the polymerization process at 72°C for 1 minute. The above is repeated 35 times, followed by a final extension step at 72°C for 10 minutes.
- Visualization of amplified products.** Once the PCR reaction is complete, the product must be visualized using 2% agarose gel electrophoresis in TAE (Tris-acetate-EDTA) buffer. This electrophoresis can be performed at 90 volts for 40 minutes, using a molecular size marker. The agarose gel is then incubated in a GelRed solution for 30 minutes at room temperature.

Finally, the gel is observed under an ultraviolet light transilluminator, and a photographic record is obtained. The positivity criterion includes the visualization of an expected DNA fragment (e.g., 400 bp) compared to the migration of the marker used. The resulting nucleotide sequences can be aligned using the Clustal Omega program, yielding a consensus sequence for each fragment. These consensus sequences can then be entered into the BLAST program to determine the percentage of nucleotide identity with respect to the Genbank® database.

- Biosafety measures.** Laboratory procedures will be carried out in accordance with the biosafety standards established for the FAVET Microbiology and Virology laboratories, which include the use of clean materials, proper waste disposal, and the use of a closed apron and gloves during laboratory work. Visualization of the magnified product will involve the use of red gel and a UV light transilluminator. Finally, an acrylic plate and UV-filtering glasses will be used when viewing the gel.

Acknowledgement

To Dr. Aron Mosnaim of the Wolf Foundation, Illinois, USA, for his continued support of our ideas and projects. This research was partially funded by Project JWF-CH-005, which is partly funded by the Wolf Foundation, Illinois, USA.

References

- Vasquez S, Aguilar F, Guerrero A, Mocha B, Baylon M, et al. (2024) Detección Molecular con la Técnica de PCR en un solo Paso para Ehrlichia Canis en Perros del Cantón Ponce Enríquez, Ecuador. Ciencia Latina Revista Científica Multidisciplinar 8(2): 8018-8026.
- Franco Zetina M, Adame Gallegos J, Dzul Rosado K (2019) Efectividad de los métodos diagnósticos para la detección de ehrlichiosis monocítica humana y canina. Revista chilena de infectología 36(5): 650-655.
- Gutierrez C, Perez Yabarra L, Agrela I F (2016) Ehrlichiosis Canina. Saber 28(4): 641-665.
- Huerto Medina E, Damaso Mata B (2015) Factores asociados a la infección por Ehrlichia canis en perros infestados con garrapatas en la ciudad de Huánuco, Perú. Revista Peruana de Medicina Experimental y Salud Publica 32(4): 756-760.
- Aragon Lopez C, Luna Nevarez P, Ortiz Encinas V, Leyva Corona J, Cantu Soto E, et al. (2021) Detección molecular de Ehrlichia canis, Anaplasma platys y Rickettsia rickettsii en caninos domésticos del municipio de Cajeme, Sonora, México. Abanico veterinario 11: e129.
- Cicuttin G, Tarragona E, De Salvo M, Mangold A, Nava S (2015) Infection with Ehrlichia canis and Anaplasma platys (Rickettsiales: Anaplasmataceae) in two lineages of Rhipicephalus sanguineus sensu lato (Acari: Ixodidae) from Argentina. Ticks Tick Borne Dis 6(6): 724-729.
- Moraes Filho J, Krawczak F, Costa F, Soares J, Labruna M (2015) Comparative evaluation of the vector competence of four south american populations of the Rhipicephalus sanguineus group for the bacterium Ehrlichia canis, the agent of canine monocytic ehrlichiosis. PLoS ONE 10(9): e0139386.

8. Ruiz A, Velarde J (2024) Frecuencia y factores asociados al diagnóstico de Ehrlichia canis y Anaplasma spp. en perros. Revista Mexicana de Ciencias Pecuarias 15(3): 749-761.
9. Foley J (2024) Ehrlichiosis, Anaplasmosis, and Related Infections in Animals. MSD Veterinary Manual.
10. Guendulain C, Tamiozzo P, Bessone A, Caffaratti M, Cabello S, et al. (2021) Detección molecular de Ehrlichia canis en perros de diferentes localidades de la provincia de Córdoba, Argentina. Ab Intus 7(4): 43-49.
11. Harrus S (2015) Perspectives on the pathogenesis and treatment of canine monocytic ehrlichiosis (Ehrlichia canis). Vet J 204(3): 239-240.
12. Fourie J, Horak I, Crafford D, Erasmus H, Botha O (2015) The efficacy of a generic doxycycline tablet in the treatment of canine monocytic ehrlichiosis. JSAVA 86(1).
13. Mctier T, Chubb N, Curtis M, Hedges L, Inskeep G, et al. (2016) Discovery of sarolaner: a novel, oral administered, broad-spectrum, isoxazoline ectoparasiticide for dogs. Vet Parasitol 222: 3-11.
14. Rudoler N, Harrus S, Martinez Subiela S, Tvarijonaviciute A, Van Straten M, et al. (2015) Comparison of the acute phase protein and antioxidant responses in dogs vaccinated against canine monocytic ehrlichiosis and naive-challenged dogs. Parasit Vectors 8: 175.
15. Mullis KB, Faloona FA (1987) Specific synthesis of DNA *in vitro* via a polymerase-catalyzed chain reaction. Methods Enzymol 155: 335-350.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2026.66.010315

Navarro Venegas C. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>