

Follow-Up of Asymptomatic Uterine Fibroids After Menopause

Hanaka Fujita, Shigeru Abe, Hiroshi Takagi, Satoshi Ichigo, Kazushige Yamamoto and Kazutoshi Matsunami, Atsushi Imai*

Department of Obstetrics and Gynecology, Matsunami General Hospital, Japan

*Corresponding author: Atsushi Imai, Department of Obstetrics and Gynecology, Matsunami General Hospital, 185-1 Dendai, Kasamatsu, Gifu 501-6062, Japan

ARTICLE INFO

Received:  September 15, 2026

Published:  September 30, 2026

Citation: Hanaka Fujita, Shigeru Abe, Hiroshi Takagi, Satoshi Ichigo, Kazushige Yamamoto and Kazutoshi Matsunami, Atsushi Imai. Follow-Up of Asymptomatic Uterine Fibroids After Menopause. Biomed J Sci & Tech Res 66(5)-2026. BJSTR. MS.ID.010419.

ABSTRACT

The development and growth of uterine fibroids are primarily influenced by ovarian estrogen and progesterone. Following menopause, the decline in ovarian steroid hormone production generally induces spontaneous regression of uterine fibroids and improvement of associated symptoms. However, this regression is not necessarily complete, and some fibroids may persist or demonstrate limited reduction in size. For asymptomatic uterine fibroids or adenomyosis, routine imaging examinations, including magnetic resonance imaging (MRI), are generally unnecessary after menopause. However, if new symptoms such as genital bleeding, lower abdominal pain, or abdominal distension develop, prompt re-evaluation is essential to exclude malignant disease. Uterine fibroids are benign mesenchymal tumours that occur in approximately 30% of women over 30 years of age and in 40–50% of women over 40 [1-3]. Although approximately half of affected women are asymptomatic, many undergo periodic clinical surveillance or receive treatment for fibroids. The development and growth of uterine fibroids are strongly influenced by ovarian estrogen and progesterone. Many clinicians are concerned about the management of uterine fibroids after menopause. This article examines the appropriate management and follow-up of uterine fibroids during the menopausal transition and postmenopausal period.

Mini Review

Following menopause, ovarian secretion of estrogen and progesterone declines substantially. Consequently, uterine fibroids generally undergo spontaneous regression, accompanied by improvement in associated symptoms, although the lesions themselves do not necessarily disappear completely. In a small proportion of women, however, fibroids may fail to demonstrate substantial regression after menopause. Possible explanations include peripheral estrogen production by adipose tissue, as well as the local production of growth factors, inflammatory mediators, or estrogen within the fibroid tissue itself [1-3]. This phenomenon may be particularly evident in women with obesity and a substantial amount of adipose tissue. Furthermore, calcification and degenerative changes within fibroids may limit or prevent appreciable reductions in lesion size. Women receiving estrogen-containing preparations [4-6] or supplements containing phytoestrogens [7-9], such as soy isoflavones (Table 1) -which are com-

monly used to alleviate menopausal symptoms or reduce the risk of osteoporosis-may likewise experience limited regression of uterine fibroids after menopause. Information regarding dietary compounds that may influence uterine fibroids is available at: <https://www.mdpi.com/article/10.3390/nu14040734/s1>.

Table 1: The active compounds from plants influencing uterine fibroids.

Active Compounds	Plants	References
Isoflavone	Soybean	[13]
Polyphenols	Green tea	[14]
Curcumin	Turmeric plants	[15]
Quercetin	Cruciferous vegetables	[16]
Indole 3-carbinol	Onion	[16]

Adenomyosis, a condition sharing certain clinical and pathological characteristics with uterine fibroids [10,11], is similarly influenced by estrogen with respect to lesion growth and symptom progression. Accordingly, both lesion size and associated symptoms generally decrease after menopause. Although adenomyosis often produces more severe symptoms than uterine fibroids, it typically exhibits a more pronounced response to the hormonal changes associated with menopause. An extremely rare condition, occurring in less than 1% of cases, is endometrial carcinoma arising within adenomyosis [12]. This malignancy is thought to result from the malignant transformation of ectopic endometrial tissue within an adenomyotic lesion and is referred to as endometrial carcinoma arising in adenomyosis (EC-AIA). Although EC-AIA is considered to occur predominantly after menopause, no specific diagnostic imaging findings have yet been established. In many cases, uterine fibroids identified after menopause do not require routine surveillance with periodic imaging examinations. However, the development of new symptoms after menopause—such as genital bleeding, lower abdominal pain, or abdominal distension, including a sensation of increased abdominal fullness or tightness of clothing or undergarments—warrants further evaluation to exclude the possibility of malignancy.

In addition, very large fibroids may be associated with age-related complications, including an increased risk of deep vein thrombosis or renal impairment. Surgical intervention may also be considered when fibroid-related mass effects and compressive symptoms, such as urinary frequency or constipation, become clinically significant. For asymptomatic uterine fibroids or adenomyosis, routine imaging examinations, including MRI, are generally unnecessary after menopause. However, the emergence of new symptoms, such as genital bleeding, lower abdominal pain, or abdominal distension, warrants prompt clinical re-evaluation to exclude malignant disease [13-16]. The menopausal transition is a period during which a variety of physical symptoms and health concerns may arise as a consequence of hormonal fluctuations, particularly alterations in estrogen levels. Careful attention to changes in one's physical condition and timely consultation with a healthcare professional when concerns arise are therefore important. Maintaining regular gynaecological examinations remains an essential component of preventive healthcare and contributes to the preservation of health and well-being throughout later life.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Yang Q, Ciebiera M, Bariani MV, Ali M, Elkafas H, et al. (2022) Comprehensive review of uterine fibroids: developmental origin, pathogenesis, and treatment. *Endocr Rev* 43(4): 678-719.
2. De La Cruz MS, Buchanan EM (2017) Uterine fibroids: diagnosis and treatment. *Am Fam Physician* 95(2): 100-107.
3. Ulin M, Ali M, Chaudhry ZT, Al HENDY A, Yang Q (2020) Uterine fibroids in menopause and perimenopause. *Menopause* 27(2): 238-242.
4. Sener AB, Seçkin NC, Özmen S, Gökmen O, Doğu N, et al. (1996) The effects of hormone replacement therapy on uterine fibroids in postmenopausal women. *Fertil Steril* 65(2): 354-357.
5. Yang CH, Lee JN, Hsu SC, Kuo CH, Tsai EM (2002) Effect of hormone replacement therapy on uterine fibroids in postmenopausal women: a 3-year study. *Maturitas* 43(1): 35-39.
6. Moro E, Degli Esposti E, Borghese G, Manzara F, Zanello M, et al. (2019) The impact of hormonal replacement treatment in postmenopausal women with uterine fibroids: a state-of-the-art review of the literature. *Medicina (Kaunas)* 55(9): 549.
7. Szydlowska I, Nawrocka Rutkowska J, Brodowska A, Marciniak A, Starczewski A, et al. (2022) Dietary natural compounds and vitamins as potential cofactors in uterine fibroids Growth and development. *Nutrients* 14(4): 734.
8. Qin H, Lin Z, Vásquez E, Luan X, Guo F, et al. (2019) High soy isoflavone or soy-based food intake during infancy and in adulthood is associated with an increased risk of uterine fibroids in premenopausal women: a meta-analysis. *Nutr Res* 71: 30-42.
9. Atkinson C, Lampe JW, Scholes D, Chen C, Wähälä K, et al. (2006) Lignan and isoflavone excretion in relation to uterine fibroids: a case-control study of young to middle-aged women in the United States. *Am J Clin Nutr* 84(3): 587-593.
10. Bulun SE, Yildiz S, Adli M, Wei JJ (2021) Adenomyosis pathogenesis: insights from next-generation sequencing. *Hum Reprod Update* 27(6): 1086-1097.
11. García Solares J, Donnez J, Donnez O, Dolmans MM (2018) Pathogenesis of uterine adenomyosis: invagination or metaplasia? *Fertil Steril* 109(3): 371-379.
12. Raffone A, Raimondo D, Maletta M, Travaglino A, Renzulli F, et al. (2023) Endometrial cancer arising in adenomyosis (EC-AIA): a systematic review. *Cancers (Basel)* 15(4): 1142.
13. Nagata C, Nakamura K, Oba S, Hayashi M, Takeda N, et al. (2009) Association of intakes of fat, dietary fibre, soya isoflavones and alcohol with uterine fibroids in Japanese women. *Br J Nutr* 101(10): 1427-1431.
14. Grandi G, Del Savio MC, Melotti C, Feliciello L, Facchinetti F (2021) Vitamin D and green tea extracts for the treatment of uterine fibroids in late reproductive life: A pilot, prospective, daily-diary based study. *Gynecol Endocrinol* 38(1): 63-67.
15. Malik M, Mendoza M, Payson M, Catherino WH (2009) Curcumin, a nutritional supplement with antineoplastic activity, enhances leiomyoma cell apoptosis and decreases fibronectin expression. *Fertil Steril* 91(5 Suppl): 2177-2184.
16. Greco S, Islam S, Zannotti A, Carpinì GD, Giannubilo SR, et al. (2020) Quercetin and indole-3-carbinol inhibit extracellular matrix expression in human primary uterine leiomyoma cells. *Reprod Biomed* 40(4): 593-602.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2026.66.010419](https://doi.org/10.26717/BJSTR.2026.66.010419)

Atsushi Imai . Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>