

Antifungal and Antibacterial Activity of *Leptodermis virgata* Ethanolic Extract Against *Candida* Species and Pathogenic Bacteria: Applications in Pharmaceutical Development, Herbal Drug Formulation, Antifungal Therapeutics and Antibacterial Agents

Aaqib Hussain Shah^{1*} and Zia Ul Hassan²

¹Department of Chemistry, Government Post Graduate College Mansehra, Pakistan

²Department of Chemistry, Government Degree College Balakot Mansehra, Pakistan

*Corresponding author: Aaqib Hussain Shah, Department of Chemistry, Government Post Graduate College Mansehra, Pakistan.

ARTICLE INFO

Received: 📅 July 17, 2026

Published: 📅 October 05, 2026

Citation: Aaqib Hussain Shah and Zia Ul Hassan. Antifungal and Antibacterial Activity of *Leptodermis virgata* Ethanolic Extract Against *Candida* Species and Pathogenic Bacteria: Applications in Pharmaceutical Development, Herbal Drug Formulation, Antifungal Therapeutics and Antibacterial Agents. Biomed J Sci & Tech Res 67(1)-2026. BJSTR. MS.ID.010428.

ABSTRACT

Fungal infections, especially those which are caused by *Candida* species, are an increasing global health concern due to the rising resistance of pathogenic microorganisms to conventional antifungal and antibacterial drugs. In this work, a systematic investigation of *Leptodermis virgata*, which is traditionally used in Himalayan shrub that belongs to the family Rubiaceae, was carried out to analyze its antifungal activity, antibacterial activity, overall antimicrobial potential, and phytochemical composition. The ethanolic extract of *L. virgata* was prepared using the cold maceration technique, and its antimicrobial efficiency was evaluated using the agar well diffusion method against *Candida albicans*, *Candida parapsilosis*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. The extract exhibited strong antifungal activity, with zones of inhibition of 13 mm against *C. albicans* and 19 mm against *C. parapsilosis*, along with notable antibacterial activity against *E. coli*, that produce an inhibition zone of 20 mm at a concentration of 25 µL. Phytochemical screening confirmed that the presence of major bioactive constituents, that includes flavonoids, tannins, terpenoids, saponins, glycosides, steroids, and anthraquinones. The results of this study is shown that *L. virgata* possesses significant antifungal, antibacterial, and antimicrobial properties, that highlights its potential as a promising natural source for the development of novel therapeutic agents.

Keywords: Phytochemistry; Antifungal Activity; Antibacterial Activity; Ethanolic Extract; *Candida* Species; Pathogenic Bacteria

Introduction

Microbiology is the branch of biological sciences that investigates the microorganisms, that includes the archaea, viruses, protozoa, bacteria, and fungi, encompassing their structure, physiology, biochemistry, ecology, and clinical significance [1,2]. The foundations of microbiology were established in the seventeenth century by Antonie van Leeuwenhoek, whose pioneering microscopic observations first revealed the previous unseen microbial world [3]. The special study of fungi, which is known as mycology, and it focuses on a diverse group of eukaryotic organisms, and whose range is from unicellular

yeasts to complex multicellular forms, that exhibits the sizes from a few micrometers to several centimeters [4]. Fungi play the vital roles in ecosystem functioning, specially in the nutrient cycling, organic matter decomposition, and the formation of mutualistic mycorrhizal associations, which support the growth and survival of over 90% of terrestrial plant species [5]. Although their ecological and economical importance, fungal biodiversity remains poorly characterized; it is estimated that only about 7-10% of fungal species have been described recently, with global diversity projected to range between 3.2 and 3.8 million species [6,7].

Fungi are classified as eukaryotic organisms, which is characterized by the presence of a true nucleus and exist as unicellular as well as multicellular forms. Based on morphological characteristics and modes of growth, fungi are categorized into yeasts, yeast-like fungi, molds, and dimorphic fungi [8,9]. Yeasts are unicellular, oval or spherical organisms that reproduce sexually through budding, with *Cryptococcus neoformans* serves as a clinically significant example [10]. Yeast-like fungi, such as *Candida albicans*, shows pseudohyphal growth, which shows the elongated cells that resemble true hyphae while retaining yeast-like characteristics [11]. In opposite, molds are multicellular fungi, that are composed of filamentous hyphae with diameters and it usually ranging from 2 to 10 μm , observed in genera such as *Aspergillus* and *Penicillium* [12]. Dimorphic fungi shows a medically important group which is capable of switching between yeast and mold forms in response of environmental conditions, with notable pathogens that includes the *Blastomyces* and *Histoplasma species* [13]. This morphological diversity underpins fungal adaptability and pathogenicity, emphasizes the clinical and biological importance of fungal classification systems.

Fungi play a very vital role and multifaceted role in human welfare, environmental sustainability, and industrial biotechnology. It is a very rich source of bioactive compounds, in which we have includes some of the important therapeutic agents that are used in modern medicine. The discovery of penicillin from *Penicillium* species is the initialization of the antibiotic era and revolutionized the treatment of bacterial infections [14,15]. Additionally, in the antibiotics, fungi make a wide range of secondary metabolites with anticancer, immunosuppressive, and cholesterol-lowering properties [16]. Many entomopathogenic fungi are used as biological control agents against insect pests, and antagonistic fungi such as *Trichoderma species*, to suppress plant pathogenic fungi, which show their importance in sustainable agriculture [17]. Additionally, fungi play a very important role in food and beverage production, with yeasts being a central component in baking, brewing, wine fermentation, and cheese manufacturing [18].

The Fungi are indispensable components of natural ecosystems due to their primary role as decomposers. It contribute to the degradation of complex organic materials, which includes the lignin, cellulose, pesticides, herbicides, and other xenobiotic compounds, and it's facilitates the nutrient recycling and environmental detoxification [19]. The enzymatic activity, fungi mineralize is dead organic matter and releases the carbon dioxide and nitrogenous compounds that sustain primary method and maintain ecological balance [20]. Unlike their numerous benefits, fungi are also responsible for significant plant, animal, and human diseases, which is ranges from superficial infections to severe systemic mycoses such as candidiasis, cryptococcal meningitis, and plant blights [21]. The incidence of fungal infections and increasing resistance rises to exists antifungal drugs underscore the urgent need for novel antifungal agents, specially those derived from medicinal plants, which served as valuable sources of

therapeutic compounds [22]. The use of plants as medicines to pre-historic times, although the exact origins of medicinal plant use remain unknown.

Archaeological evidence indicates that plants are used therapeutically as early as 60,000 years ago, which is shown by plant remains discovered in the grave of a Neanderthal in present-day Iraq, as well as medicinal plants that are found among the personal belongings of Ötzi the Iceman [23,24]. The earliest written medical records, dating back approximately 4,000–5,000 years, originate from ancient civilizations such as Mesopotamia, where Sumerian clay tablets documented herbal drug preparations is derived from over 250 plant species [25]. Traditional systems of medicine, specially Ayurveda which is often regarded as the oldest organized healthcare system and emphasizes the therapeutic value of medicinal plants [26]. Similarly, ancient Chinese pharmacopoeias, notably the *Pen T'Sao* attributed to Emperor Shen Nung (circa 2500 BC), described hundreds of medicinal plants, many of which remain in use today [27].

Leptodermis virgata belongs to the family Rubiaceae, one of the large angiosperm families, that comprises approximately 13,500 species distributed among about 620-630 genera with a cosmopolitan presence, specially concentrated in tropical and subtropical regions [28,29]. Members of Rubiaceae are characterized by opposite or verticillate leaves with interpetiolar stipules, actinomorphic sympetalous flowers, and predominantly woody growth forms, that includes the shrubs and trees [30]. The genus *Leptodermis* Wall. is mainly distributed across the Himalayan region, China, and Japan, and derives its name from the Greek words *leptos* (thin) and *derma* (skin), that referes to its thin, membranous bractlets [31]. Species within this genus are shrubby, emit a characteristic odor when bruised, and possess opposite leaves with persistent stipules, although the genus remains taxonomically complex [32]. *Leptodermis virgata* is an erect, much-branched shrub typically 30–70 cm in height, bearing linear leaves, purplish flowers, and cylindrical capsules with fibrous seeds, features that distinguish it from related species [33]. Given the medicinal importance of several Rubiaceae members, including *Coffea* and *Cinchona*, the biological and pharmacological potential of *L. virgata* warrants systematic scientific investigation. *Leptodermis virgata* (Rubiaceae) is a shrubby species distributed predominantly in the Western Himalayas, that includes the regions of Pakistan, where it inhabits dry, stony soils up to elevations of approximately 2300 m. The family of Rubiaceae, one of the largest angiosperm families, is taxonomically and economically significant, encompasses in medicinal, ornamental, and industrially very important plants such as *Coffea* and *Cinchona*. The members of this family are characterized by opposite leaves with interpetiolar stipules, sympetalous flowers, and predominantly woody growth forms, features that are well expressed in the genus *Leptodermis*. Now beyond its taxonomic relevance, it increases the scientific interest which has focused on the pharmacological potential of Rubiaceae taxa, specially their antifungal proper-

ties and rich phytochemical profiles. Antifungal agents, historically rooted in the concept of chemotherapy which is introduced by Paul Ehrlich, and it play a very critical role in combating fungal infections through selective inhibition of fungal cell wall or membrane synthesis. Concurrently, plant-derived phytochemicals such as flavonoids, alkaloids, terpenoids, tannins, and phenolic compounds have been broadly recognized for their anti-microbial and therapeutic efficacy. Standardized bioassays, that includes the agar diffusion (Kirby-Bauer) method, and it provides the reliable means to evaluate antifungal

activity. In this method, the present study shows the botanical classification, antifungal relevance, and phytochemical significance of *L. virgata*, that supports its potential as a valuable source of bioactive compounds for pharmaceutical applications. Figure 1 shows the natural growth of *Lip-todermis virgata*, presenting its fine branching structure and dense green foliage. The image shows the characteristic leaf arrangement and morphology, which support botanical identification and ecological assessment.



Figure 1: Liptodermis virgata.

Main Contributions

The main contributions of this study are summarized as follows:

- a. A comprehensive investigation of *Leptodermis virgata*, is used in Himalayan shrub of the family Rubiaceae, is conducted to evaluate its antimicrobial potential.
- b. The study systematically analyzed the antifungal activity, antibacterial activity, overall antimicrobial efficacy, and phytochemical composition of *L. virgata* in laboratory.
- c. Ethanolic extracts of *L. virgata* are prepared using the cold maceration technique, and their efficacy is tested against multiple pathogenic microorganisms, that includes the *Candida albicans*, *Candida parapsilosis*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*.
- d. The extract shows the strong antifungal activity with inhibition zones of 13 mm and 19 mm against *C. albicans* and *C. parapsilosis*, respectively, and notable antibacterial activity against *E. coli* with a 20 mm inhibition zone at 25 μ L.
- e. Phytochemical screening reveals the presence of key bioactive compounds, that includes the flavonoids, tannins, terpenoids,

saponins, glycosides, steroids, and anthraquinones, which contribute to the observed antimicrobial effects.

- f. This research highlights the potential of *L. virgata* as a natural source for the development of novel antifungal and antibacterial agents.

Background Study

The use of medicinal plants is a cornerstone of traditional medicine, with knowledge passed through generations for the treatment of human and animal diseases. Ethnic groups worldwide have relied on plants as sources of natural remedies, primarily due to their production of antimicrobial secondary metabolites, and it offers a relatively non-toxic and cost-effective method to synthetic drugs. Medicinal plants are evaluated based on their therapeutic, aromatic, or salty qualities, and are collected particularly for medicinal purposes and are recognized as true medicinal plants. The antimicrobial activity of these plants is linked to the presence of volatile oils and terpenes, and abundant in essential oils which is used in pharmaceuticals, cooking, and perfumery [34,35]. Plant extracts & essential oils are evaluated for their antifungal and antibacterial properties, and it shows their importance as natural bioactive compounds.

Many studies show the potent antimicrobial activity of plant extracts against both bacteria and fungi. Like, eleven ethanolic extracts from spices such as *Melissa officinalis*, *Mentha piperita*, *Laurus nobilis*, and *Juniperus oxycedrus*, and assayed against Gram-positive (*Bacillus subtilis*, *Staphylococcus aureus*, *S. epidermidis*) and Gram-negative (*Escherichia coli*, *Pseudomonas aeruginosa*) bacteria, as well as *Candida albicans* and *Aspergillus niger*. The minimum inhibitory concentration (MIC) of *M. piperita*, *L. nobilis*, and *J. oxycedrus* extracts is 5 mg/mL for all tested microorganisms, with *P. aeruginosa* being specially sensitive to *P. nigrum* and *E. arborea* extracts. The extracts of *L. no-*

bilis, *D. coryophyllum*, *J. oxycedrus*, and *C. arborescens* demonstrated higher antifungal activity than the standard drug nystatin [36]. Figure 2 shows the molecular structure of sulfanilamide, presenting a benzene ring which is substituted with amino and sulfonamide functional groups. The arrangement of these groups gives information about its chemical reactivity and underlies its antibacterial properties in pharmaceutical applications. Figure 3 shows the different regions on a petri dish, which includes the fungal development around the periphery and a distinct clear zone. The inhibition zone near the center shows the suppressed microbial growth due to the applied agent.




Figure 2: Structure of Sulfanilamide.




Figure 3: Different regions on the petri dish.

Similarly, six ethanolic extracts from spices including ginger (*Zingiber officinale*), cinnamon (*Cinnamomum verum*), black cumin (*Nigella sativa*), clove (*Syzygium aromaticum*), black pepper (*Piper nigrum*), and chamomile (*Anthemis nobilis*) were tested for antimicrobial activity against standard bacterial and fungal strains. Clove extract exhibited strong inhibitory activity against *B. subtilis*, *P. aeruginosa*, and *C. albicans*, whereas cinnamon was effective against *B. subtilis* and *C. albicans* only. Other extracts show either no activity or stimulatory effects. *S. aureus* and *E. coli* are resistant to most extracts, clove and cinnamon shown to potent antimicrobial effects at low MIC values and it ranges from 3.125 to 25 µg/mL [37].

Traditional Chinese medicine and ethnobotanical knowledge historically provided a rich source of empirical information on the therapeutic value of plants. Before the emergence of modern medicine, herbalists accumulated significant knowledge about plant-based remedies through empirical observations of efficacy, failures, and successes. Modern pharmacology has built upon these traditional practices, that isolate active compounds from plant extracts for therapeutic purposes. Today, medicinal plants remain very important in combating infectious diseases; they are rich in secondary metabolites such as alkaloids, flavonoids, tannins, and terpenoids, which exhibit antibacterial, antifungal, and anticancer activities [38]. The study on plant extracts shown their selective antimicrobial and antifungal activities. Among 22 plant extracts evaluated, clove and ginger show the inhibitory activity against all tested fungal isolates, whereas garlic was effective against most fungi except *Penicillium oxalicum*. Extracts of cumin, turmeric, and cinnamon displayed activity against a limited number of fungi. Clove extract shows the highest mycelial growth inhibition (70-100%), which is followed by ginger extract (30-100%), and this suggests their potential as natural antimicrobial additives to improve food shelf-life.

The pharmacological potential of medicinal plants is also reflected in their chemical composition. Essential oils, flavonoids, terpenoids, saponins, steroids, and phenolic compounds contribute to the biological activity of plants. These phytochemicals not only confer the antimicrobial properties and also serve as antioxidants and nutraceuticals. The use of plant compounds like food preservatives is recognized due to consumer preference for natural products [39]. Research in different areas, like the Himalayas and Cameroon, highlights the nutritional and medicinal significance of indigenous plants. *Pimpinella stewartii* is identified as a rich source of phenolic compounds and antioxidants. And Cameroonian medicinal plants are reveals the presence of tannins, flavonoids, saponins, steroids, glycosides, and alkaloids, emphasizing their therapeutic potential. Certain plants, such as *Morinda lucida* and *Spilanthes filicaulis*, show the degrees of

chemical diversity, which directly influence their efficacy in treating diseases [40-43].

Materials and Methods

Materials

Chemicals

a) For antifungal activity: The chemicals which are used for the evaluation of antifungal activity includes ethanol, distilled water, dimethyl sulfoxide (DMSO), and agar.

b) For phytochemical screening: The reagents which is employed for preliminary phytochemical analysis are ferric chloride (FeCl_3), hydrochloric acid (HCl), magnesium ribbon, lead acetate, acetic acid (CH_3COOH), sulphuric acid (H_2SO_4), Fehling's solution, acetic anhydride, chloroform (CHCl_3), and ammonia.

Apparatus

The apparatus are used in the present study comprises mortar and pestle, porcelain cloth, Whatman filter paper, beakers, volumetric flasks, micropipettes, aluminum foil, a magnetic stirrer, Eppendorf tubes, a cork borer, a digital balance, and sterilized Petri dishes.

Methodology

A systematic methodology is adopted to evaluate the antifungal activity of *Leptodermis virgata*. The experimental procedure involves the sequential steps, includes the collection and grinding of the plant material, preparation of crude extracts, preparation of a suitable culture medium, culturing of fungal strains with well formation, and assessment of antifungal activity of the plant extract.

Collection and Grinding of Plant Material

The plant, *Leptodermis virgata*, is collected from multiple locations in the Mansehra district. Initially, the samples are taken from Khait Sarash and Mang, Tehsil Balakot. Additional collections also collect from Tehsil Darband, Village Tarmang, and Tehsil Oghi, Village Chamial, and it is confirmed the presence of *L. virgata* in these regions. The collected plant material is shade dried at room temperature for a period of 65-70 days, in order to preserve its phytochemical constituents. After complete drying of plant, the whole plant, which includes the flowers, leaves, stems, bark, and branches, is finely ground into a uniform powder using a mortar and pestle. The powdered material is stored in clean, airtight containers for further experimental use. Figure 4 shows the powdered form of *Liptodermis virgata* after grinding, which presents a uniform and fibrous texture. This processed form is suitable for extraction and further phytochemical and biological analysis.



Figure 4: Grinded form of plant leptodermis virgata.

Preparation of Crude Extracts

The ethanolic extract of *Leptodermis virgata* is prepared using the maceration technique. An estimated 4 kg of the dried and powdered plant material is soaked in 7.5 L of ethanol at room temperature. The mixture is intermittently stirred two to three times within a 24-hour period using a stainless-steel rod to guarantee the maximum extraction of bioactive constituents. After maceration, the extract is first filtered through a porcelain cloth and subsequently re-filtered using Whatman filter paper to obtain a clear filtrate. This extraction and

filtration procedure is repeated three times to enhance this process. The combined filtrates are then left in the open air to allow complete evaporation of ethanol. This solvent evaporation process requires approximately two months, which show the results in a concentrated crude extract that is stored for further feature experimental use. Figure 5 shows the plant material soaked in ethanol, show the extraction stage where bioactive compounds are dissolved into the solvent. Figure 6 shows the solid plant extract and is obtained after solvent removal, it also represents the concentrated form which is also used for further experimental analysis.



Figure 5: Plant soaked in ethanol.



Figure 6: Solid extract of plant sample.

Preparation of Growth Medium

We evaluate the antifungal activity, and a suitable growth medium is prepared before microbial inoculation. We select nutrient agar as the culture medium due to its ability to support the growth of a wide range of microorganisms, and this includes fungi. The preparation method involves the dissolving of 20 g of nutrient agar powder in 500 mL of distilled water in a conical flask. The mixture is also heated with continuous stirring using a magnetic stirrer until the complete dissolution of the agar components is achieved. The flask is then covered with aluminum foil and autoclaved at 100°C on a hot plate for one hour. After sterilization, the molten agar is allowed to cool without solidification and subsequently poured into sterile Petri dishes. The plates are then placed in a pressure steam sterilizer to guarantee the complete sterilization and left undisturbed until the agar solidifies.

Culturing of Fungi

Many techniques are available for fungal culturing, and the agar plate method is used in the present study. Fungi are ubiquitous organisms that exist as unicellular or complex multicellular forms and it play the essential roles in nutrient cycling within ecosystems. Some fungal species are pathogenic and responsible for diseases in plants, animals, and humans, which leads to significant agricultural and health-related losses. For the assessment of antifungal activity, two yeast species are selected as test organisms.

a) **Candida albicans:** *Candida albicans* is a pathogenic yeast that is commonly found as a commensal organism in the human gastrointestinal tract, but it may also exist outside the human body. It is also responsible for superficial infections such as oral and vaginal candidiasis, as well as it severe systemic infections that can be life-threatening, especially in immuno-compromised individuals.

b) **Candida parapsilosis:** *Candida parapsilosis* is the second or

third most isolated fungal species from clinical specimens. It is a yeast associated with sepsis, wound infections, and tissue infections, especially in immunodeficient patients.

Assessment of Antifungal Activity

Antifungal activity is defined as the ability of a compound to selectively inhibit or kill fungal pathogens without causing damage to host tissues. Recently developed antimicrobial agents are chemically modified derivatives of natural compounds. The foundation of modern chemotherapy can be attributed to Paul Ehrlich (1910), who introduced the concept of chemotherapy using Salvarsan, an arsenic-based compound effective against syphilis. He also emphasized the selective toxicity of chemotherapeutic agents against pathogens. For antifungal evaluation, a solution of the crude plant extract is prepared using dimethyl sulfoxide (DMSO) as the solvent. Approximately 20 mg of crude extract is dissolved in 2 mL of DMSO in an Eppendorf tube and thoroughly shaken to guarantee the complete dissolution. Using a micropipette, 25 µL of the prepared solution is carefully dispensed into wells that is formed in the agar plates inoculated with fungal cultures. Only DMSO is used as a negative control and is added to a separate well. The same method is repeated for all Petri dishes. The inoculated plates are then incubated at appropriate conditions for 24 hours, after which antifungal activity is assessed by observation in a zones of inhibition around the wells.

Phytochemical Analysis

We also performed the phytochemical analysis in order to identify and evaluate the presence of various bioactive compounds in *Lepidodermis virgata*. This screening is necessary for understanding the therapeutic potential of the plant and for correlating observed biological activities with specific classes of secondary metabolites. The phytochemical tests conducted in this study are follows:

- **Test for Alkaloids:** We have detected the nitrogen-containing basic compounds, which are known for pharmacological activities.
- **Test for Flavonoids:** We identified the polyphenolic compounds with antioxidant and antimicrobial properties.
- **Test for Reducing Sugars:** We also determined the presence of monosaccharides or other reducing sugars.
- **Test for Saponins:** We detected the glycosidic compounds that show the antifungal and surface-active properties.
- **Test for Tannins:** We identified phenolic compounds that are capable of precipitating proteins and have contribution to the antimicrobial activity.
- **Test for Steroids:** We also determined the presence of sterol-based compounds with potential anti-inflammatory and antimicrobial effects.
- **Test for Terpenoids:** We have detected the compounds which are known for their diverse pharmacological activities, that includes the antifungal and antibacterial effects.

These qualitative tests provide us with a preliminary insight into the phytochemical composition of *L. virgata*, which contributes to its observed antifungal and antibacterial activities.

Results

Antifungal Activity

We evaluated the antifungal activity of the ethanolic extract of *Leptodermis virgata* using the agar well diffusion method. We also prepared the 25 μ L solution of the extract in an Eppendorf tube and is subsequently added to the wells created using a cork borer on agar plates inoculated with the test fungi. We also determined the effectiveness of the extract which is measured the zones of inhibition around the wells after incubation. These results present that the *Candida albicans* gives us a zone of inhibition of 13 mm, while other *Candida parapsilosis* displays a larger zone of inhibition measuring 19 mm, which shows the stronger antifungal activity against *C. parapsilosis*. (Table 1) Figure 7 shows the antifungal activity of the plant extract against *Candida albicans*, which is evidenced by the visible inhibition of fungal growth. Figure 8 gives us a similar antifungal response against *Candida parapsilosis*, which indicates the broad-spectrum effectiveness of the extract. Figure 9 shows the inhibition zone, which is measured at a sample concentration of 25 microliter for two fungal strains. The extract shows the stronger antifungal activity against *Candida parapsilosis* and is compared to *Candida albicans*, which is indicated by the larger inhibition zone.




Figure 7: Indicates the antifungal activity of Candida Albicans.




Figure 8: Indicates the antifungal activity of Candida Parasilosis.




Figure 9: Graphical representation of inhibition vs concentration of sample.

Table 1: Zone of Inhibition of Fungi by Leptodermis virgata Ethanolic Extract.

Name of Fungi	Concentration of Solution (μL)	Zone of Inhibition (mm)
Candida albicans	25	13
Candida parapsilosis	25	19

Antibacterial Activity

The antibacterial activity of the ethanolic extract of *Leptodermis virgata* is assessed using the agar well diffusion method. A 25 μL solution of the extract is added to wells in nutrient agar plates inoculated with the test bacterial strains. The zones of inhibition are measured

after incubation to evaluate the effectiveness of the extract against each bacterial species. The results show that *E. coli* is the highest susceptibility with a zone of inhibition of 20 mm. Both *Pseudomonas aeruginosa* and *Streptococcus aureus* give us s smaller zones of inhibition, measuring the 6 mm each, suggesting comparatively lower anti-bacterial activity against these species. (Table 2) Figure 10 presents the zone of inhibition observed at a sample concentration of 25 microliter against different bacterial strains. The extract shows strong antibacterial activity against *Escherichia coli*, while moderate inhibition is observed for *Pseudomonas aeruginosa* and *Staphylococcus aureus*. Figures 11–13 shows the antibacterial activity of the plant which is extract against three clinically relevant bacterial strains.

Table 2: Zone of Inhibition of Bacteria by *Leptodermis virgata* Ethanolic Extract.

Name of Bacteria	Concentration of Solution (µL)	Zone of Inhibition (mm)
<i>Pseudomonas aeruginosa</i>	25	6
<i>Streptococcus aureus</i>	25	6
<i>E. coli</i>	25	20

Figure 11 gives the information about the response of *Pseudomonas aeruginosa*, where a distinct inhibition zone confirms the sensi-

tivity of this strain to the extract. Figure 12 demonstrates the antibacterial effect against *Staphylococcus aureus*, which indicates moderate growth suppression around the applied sample. Figure 13 shows the antibacterial activity against *Escherichia coli*, where a comparatively larger inhibition zone reflects stronger susceptibility. The variation in inhibition patterns shows the strain-dependent antibacterial potential of the plant extract. These observations suggest differences in cell wall structure and metabolic response among Gram positive and Gram-negative bacteria. The experimental results support the broad-spectrum antibacterial capability of the extracted compounds.




Figure 11: Indicates the antibacterial activity of *Pseudomonas Aeregenosa*.




Figure 12: Indicates the antibacterial activity of *Streptococcus Aureus*.




Figure 13: Indicates the antibacterial activity of E Coli.

Phytochemical Analysis

Qualitative phytochemical analysis is performed to detect and localize bioactive compounds present in the ethanolic extract of *Lep-todermis virgata*. Different standard tests are used to identify the various classes of phytochemicals, that are:

- a) **Flavonoids:** The alkaline reagent test is used for flavonoids. Two milliliters of 2% NaOH solution are added to the crude extract. The appearance of a yellow color indicates the presence of flavonoids.
- b) **Saponins:** The extract is mixed with 5 mL of distilled water in a test tube and warmed. The absence of significant change indicates the presence of saponins in the extract.
- c) **Alkaloids:** Salkowski's test is performed by mixing 2 mL of chloroform with the crude extract, which is followed by 2 mL of concentrated sulfuric acid. The development of a green color confirmed the presence of alkaloids.
- d) **Anthraquinones:** We extract the two milliliters of plant and then mixed with 1 mL of 10% ammonia solution and shaken for 4–5 minutes. The formation of a deep red color shows the presence of anthraquinones.

- e) **Tannins:** We also conduct the Ferric chloride test by stirring 2 mL of plant extract with distilled water and adding a few drops of FeCl_3 . The formation of a green precipitate shows the presence of condensed tannins.
- f) **Steroids:** We also added the two milliliters of chloroform into 2 mL of plant extract, which is followed by 1 mL of concentrated sulfuric acid. The appearance of a blue-green color indicates the presence of steroids.
- g) **Terpenoids:** We dissolve two milliliters of plant extract in 2 mL of chloroform, and 2 mL of concentrated sulfuric acid. The mixture is heated for 2 minutes, and the appearance of a greyish color confirmed the presence of terpenoids. Figure 14 shows the phytochemical screening of the plant extract using standard qualitative tests. Distinct color changes and precipitate formations show the presence of bioactive constituents such as alkaloids, flavonoids, tannins, and phenolic compounds. These phytochemicals are likely responsible for the observed antibacterial effects, and they validate the medicinal relevance of the plant extract.



Figure 14: Indicates the phytochemical test of plant extract.

Results of Phytochemical Tests

The qualitative phytochemical analysis of the ethanolic extract of *Leptodermis virgata* shows the presence of various bioactive compounds. The results are summarized in Table 3.

Table 3: Results of phytochemical tests for *Leptodermis virgata* ethanolic extract.

S.No	Test	Result
1	Flavonoids	+
2	Saponins	+
3	Glycosides	+
4	Anthraquinones	+
5	Tannins	+
6	Steroids	+
7	Terpenoids	+

Discussion

The active components of the plant *Leptodermis virgata* are extracted using ethanol and evaluate for their antibacterial and antifungal activities against some selected pathogenic microorganisms. The ethanolic extract also shows the activity against *Pseudomonas aeruginosa*, *Streptococcus aureus*, and *E. coli* for antibacterial testing, against *Candida albicans* and *Candida parapsilosis* for antifungal testing. We also observed the antimicrobial activity at room temperature and measured it through the zones of inhibition on agar plates. The results show the extract contains bioactive compounds such as flavonoids, terpenoids, glycosides, tannins, saponins, steroids, and anthraquinones, which contribute to the observed antimicrobial effects. These results give us the guarantee that *L. virgata* can be utilized for the treatment of gastroenteritis, urethritis, wound infections, and fungal diseases like sepsis.

Conclusion

The results of this study reveal that the ethanolic extract of *Leptodermis virgata* possesses antifungal, antibacterial, and overall antimicrobial activities. The extract shows the stronger antibacterial activity against *Escherichia coli* and as compared to *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Moreover, both tested fungal strains, i.e. *Candida albicans* and *Candida parapsilosis*, show the marked susceptibility to the plant extract, show the antifungal activity of *L. virgata* is more pronounced and useful than its antibacterial activity. Phytochemical screening confirmed the presence of many major bioactive constituents, which includes flavonoids, tannins, terpenoids, saponins, glycosides, steroids, and anthraquinones, which are collectively contribute to the observed antimicrobial effects. These findings support the traditional use of *L. virgata* in the treatment of various infectious diseases, especially in fungal infections. Further studies will be used to investigate its efficacy against a broader spectrum of bacterial and fungal pathogens, as well as to isolate and characterize the active compounds responsible for its antimicrobial activity, that develop novel plant-based therapeutic agents.

Data Availability

No datasets were used or analyzed in this study.

Acknowledgements

The author is deeply grateful to his parents and uncle, especially Nazakat Hussain Shah, for their unwavering support and encouragement throughout his education and this research. The author also extends sincere thanks to the Department of Chemistry, Government Degree College Balakot, Mansehra, for their valuable support.

Conflict of Interest

There is no conflict of interest among authors, and all authors did equal work.

References

- Wiarat C (2006) *Etnopharmacology of medicinal plants*. Humana Press, New Jersey.
- Tucakov J (1971) *Healing with plants – phytotherapy*. Culture, Beograd.
- Glesinger L (1954) *Medicine through centuries*. Zora, Zagreb.
- Tucakov J (1964) *Pharmacognosy*. Institute for textbook issuing in SR SR-bije, Beograd.
- Dimitrova Z (1999) *The history of pharmacy*. St Clement of Ohrid, Sofija.
- Toplak Galle (2005) *Domestic medicinal plants*. Mozaic book, Zagreb.
- Bojadzievski P (1992) *The health services in Bitola through the centuries*. Society of science and art, Bitola.
- Bazala V (1943) *The historical development of medicine in the Croatian lands*. Croatian publishing bibliographic institute, Zagreb.
- Nikolovski B (1995) *Essays on the history of health culture in Macedonia*. Macedonian pharmaceutical association, Skopje.
- Thorwald J (1991) *Power and knowledge of ancient physicians*. August Cesarec, Zagreb.
- Katic R (1967) *The Serbian medicine from 9th to 19th centuries*. Scientific work, Beograd.
- Stojcevska VA (1982) *Clement and naum of ohrid in folk tradition*. Our book, Skopje.
- Katic R (1958) *La medicine en serbie au moyen age*. Scientific work, Beograd.
- Igersheim A, C Puff, P Leins, C Erbar (1994) Gynoecial development of gaertnera lam. and of presumably allied taxa of the psychotrieae (rubiaceae): Secondary 'superior' vs. inferior ovaries. *Botanische Jahrbuchher fu'r Systematik* 116: 401-414.
- Robbrecht E (1988) Tropical woody rubiaceae. *Opera Botanica Belgica* 1: 1-271.
- Anderson WR (1973) A morphological hypothesis for the origin of heterostyly in the rubiaceae. *Taxon* 22(5/6): 537-542.
- Roskov Y, T Kunze, T Orrell, L Abucay, Paglinawan AL, et al. (2014) *Species 2000 & itis catalogue of life: 2014 annual checklist, 2014*. Edited by V. Didz'ulis.
- Davis AP, R Govaerts, DM Bridson, M Ruhsam, J Moat, et al. (2009) A global assessment of distribution, diversity, endemism, and taxonomic effort in the rubiaceae. *Annals of the Missouri Botanical Garden* 96(1): 68-78.
- Bolhuis MS, PN Panday, AD Pranger, JG Kosterink, JC Alfenaar (2011) Pharmacokinetic drug interactions of antimicrobial drugs: A systematic review on oxazolidinones, rifamycines, macrolides, fluoroquinolones, and beta-lactams. *Pharmaceutics* 3(4): 865-913.
- Mesa Arango AC, L Scorzoni, O Zaragoza (2012) It only takes one to do many jobs: Amphotericin b as antifungal and immunomodulatory drug. *Front Microbiol* 3: 286.
- Swain SS, SK Paidesetty, RN Padhy (2017) Antibacterial, antifungal and antimycobacterial compounds from cyanobacteria. *Biomed Pharmacother* 90: 760-776.
- Nakatani N (1994) Antioxidative and antimicrobial constituents of herbs and spices. In G. Charalambous, 1st editor, *Spices, Herbs, and Edible Fungi*, Elsevier, Amsterdam, Pp 251-272.
- Hamburger M, K Hostettmann (1991) Bioactivity in plants: The link between phytochemistry and medicine. *Phytochemistry*, 30(12): 3864-3874.
- Ahn K (2017) The worldwide trend of using botanical drugs and strategies for developing global drugs. *BMB Rep* 50(3): 111-116.
- Sumner J (2000) *The Natural History of Medicinal Plants*. Timber Press, pp 235.
- Avato P, C Vitali, P Mongelli, A Tava (1997) Antimicrobial activity of polyacetylenes from bellis perennis and synthetic derivatives. *Planta Medica*, 63(6): 503-507.
- Ababutain IM (2011) Antimicrobial activity of ethanolic extracts from some medicinal plants. *Australian Journal of Basic and Applied Sciences* 5(11): 678-683.
- Abalaka ME, A Mann, SO Adeyemo (2011) Studies on in-vitro antioxidant and free radical scavenging potential and phytochemical screening of leaves of ziziphus mauritiana l. and ziziphus spina christi l. compared with ascorbic acid. *Journal of Medical Genetics and Genomics* 3(2): 28-34.
- Agarwal SK, SS Singh, S Verma, S Kumar (2000) Antifungal activity of anthraquinone derivatives from rheum emodi. *J Ethnopharmacology* 72(1-2): 43-46.
- Garbacki N, L Angenot, C Bassleer, J Damas, M Tits (2002) Effects of prodelphinidins isolated from ribes ni-grum on chondrocyte metabolism and cox activity. *Naunyn-Schmiedeberg's Archives of Pharmacology* 365 (6): 434-441.
- Chung IM, M Ali, SJ Hahn, NA Siddiqui, Y Lim, et al. (2005) Chemical constituents from the hulls of oryza sativa with cytotoxic activity. *Chemistry of Natural Compounds* 41: 182-189.
- Fujii Y, M Furukawa, Y Yakawa, K Sugawara, T Shibuya (1991) Survey of japanese medicinal plants for the detection of allelopathic properties. *Weed Research Japan* 36 (1): 36-42.
- Lambert J, J Srivastava, N Vietmeyer (1997) *Medicinal plants. Rescuing a global heritage*. World Bank, Wash-ington DC. World Bank Technical Paper 1: 355.
- Ramu G, GK Mohan (2012) Preliminary phytochemical and antioxidant study of hydroalcoholic extracts from selected genera of indian lamiaceae. *Asian Pacific Journal of Tropical Biomedicine* Pp: 685-688.
- Ozcan M, et al. (2011) Antimicrobial properties of spice extracts. *Food Control*.
- Al Snafi AE (2015) Pharmacological importance of medicinal plants. *International Journal of Pharmacology*.
- Shen X, et al. (2005) Traditional chinese medicine: Pharmacology and clinical applications. *Journal of Ethnopharmacology*.
- Liu X et al. (2010) Ethnobotanical and phytochemical studies of medicinal plants. *Ethnobotany Research and Applications*.
- Burt S (2004) Essential oils: Their antimicrobial properties and applications. *Int J Food Microbio* 94(3): 223-253.
- Boklari FM (2009) Antifungal activity of some medicinal plants used in jeddah, saudi arabia. *Mycopath* 7(1): 51-57.
- Dukic NM, B Bozin, M Sokovic, N Simin (2004) Antimicrobial and antioxidant activity of melissa officinalis l. (lamiaceae) essential oil. *J Agric Food Chemistry* 52(9): 2485-2489.

42. Souza DEL, TLM Stamford, EO Lima, JMB Filho (2005) Antimicrobial effectiveness of spices: An approach for use in food conservation system. Braz Arch Biol Technol 48(4): 549-558.
43. Pundir RK, P Jain (2010) Antifungal activity of twenty two ethanolic plant extracts against food-associated fungi. Journal of Pharmacy Research 3(1): 506-510.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2026.67.010428

Aaqib Hussain Shah . Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>