

# CFTR is a Target of Particulate Matter Exposure Beyond Lung Epithelium

Aztatzi Aguilar OG<sup>1\*</sup>, Debray-García Y<sup>1†</sup>, Guerrero-Cruz M<sup>2†</sup>, Uribe-Ramírez TM<sup>3</sup>, Narváez-Morales J<sup>3</sup> and Barbier O<sup>3</sup>

<sup>1</sup>Instituto Nacional de Enfermedades Respiratorias Ismael Cosío Villegas. Departamento de Investigación en Toxicología y Medicina Ambiental, CDMX, México.

<sup>2</sup>Universidad del Valle de México. Campus Chapultepec, CDMX, México.

<sup>3</sup>Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional. Departamento de Toxicología, CDMX, México.

ORCID of Debray Gracia Y: 0000-0002-3247-2841

ORCID OF Uribe-Ramírez TM: 0000-0003-3282-1287

ORCID of Narváez-Morales J: 0000-0002-6411-036X

ORCID of Barbier O: 0000-0002-0378-3675

ORCID of Aztatzi-Aguilar O.G: 0000-0003-1761-9443

**\*Corresponding author:** Aztatzi Aguilar OG, Instituto Nacional de Enfermedades Respiratorias Ismael Cosío Villegas, Departamento de Investigación en Toxicología y Medicina Ambiental, CDMX, México.

## ARTICLE INFO

**Received:** 📅 August 25, 2026

**Published:** 📅 October 09, 2026

**Citation:** Aztatzi Aguilar OG, Debray-García Y, Guerrero-Cruz M, Uribe-Ramírez TM, Narváez-Morales J and Barbier O. Hypertensive Crisis Associated with Methylphenidate Use During Total Intravenous Anaesthesia for Orthognathic Surgery: A Case Report. Biomed J Sci & Tech Res 67(1)-2026. BJSTR. MS.ID.010432.

## ABSTRACT

The cystic fibrosis transmembrane conductance regulator (CFTR) is implicated in epithelial ion transport and mucus secretion; dysfunction or lower expression is related with physiological dysfunction of organs such as lungs. It has been reported that cigarette smoke and diesel particles affect CFTR expression in lung epithelium models, but not urban particles. The goal of the present brief communication is to present evidence of urban particles affect CFTR in an *in vitro* and *in vivo* experimental model affecting also non-respiratory tissues. In A549 cell line after 24h exposure to PM<sub>10</sub> and PM<sub>2.5</sub> from subway and outdoor environments evaluate the CFTR expression and intracellular chloride concentrations. On the other hand, in male rats evaluated CFTR expression after acute, and sub-chronic exposure to PM<sub>2.5</sub> and filtered air (FA) in lung and kidney cortex tissues; serum- and urine-chloride concentrations were evaluated. *In vitro*, assays demonstrated that PM<sub>10</sub> and PM<sub>2.5</sub> can modulate CFTR expression and increased intracellular chloride concentrations. *In vivo* results showed that PM<sub>2.5</sub> exposure reduced CFTR protein in lungs and kidney cortex, additionally, serum and urine chloride concentrations exhibited time-dependent changes. CFTR is a toxicological target of urban PM, which affects expression, and local and systemic chloride ion concentrations changes.

**Keywords:** Particulate matter, CFTR, lung tissue, non-respiratory tissues, kidney.

## Introduction

Air pollution by particulate matter (PM) is associated with the development of cardio-respiratory diseases [1,2] and there is evidence that air quality also contributes to neurological [3], and renal diseases [4]. The PM mechanism related with its toxicity is unknown and unexplained to the many diseases associated with air pollution by PM. Additionally, chronic diseases worsen under polluted ambient

air conditions, such as metabolic diseases [5]. Some molecular targets have been suggested which mediate PM toxicity, that include Transient Receptor Potential (TRP) family channels [6], which are able to interact with other co-modulators channels such cystic fibrosis transmembrane conductance regulator (CFTR) [7]. The CFTR gene codifies for an integral membrane protein, which is a member of the ABC transporter family and is implicated in chloride and bicarbonate ion

transport in many epithelial secretions. CFTR also mediates anionic glutathione flux, which is the most important extracellular non-enzymatic antioxidant in the lungs [8,9]. The absence or malfunctioning of CFTR is implicated in Cystic fibrosis (CF), a hereditary recessive autosomal disorder caused by a mutation in CFTR gene [10]. The different CFTR mutations in the CF patients are related with diverse effects including airways, pancreas, male genital system, intestine, liver, bone, and kidney. Besides the fat malabsorption, CF causes chronic pulmonary infections that ultimately drives the development of bronchiectasis and progressive lung damage [11]. In CF, lungs become chronically infected by pathogens such *Pseudomonas aeruginosa*, suggested by the altered anionic composition of airway surface fluid of CF patients which affect the activity of innate immune response proteins called defensins. For the above reason CFTR it has been suggested as a pleiotropic protein [12].

PM exposure has been associated with the increment of respiratory infections. Fine particle exposure has been related with initial *Pseudomonas aeruginosa* infection in the human alveolar epithelial cell line A549, and corroborate through epidemiological studies, where it was reported a risk augment (23-26%) in the *P. aeruginosa* acquisition by each 10 µg/m<sup>3</sup> augment in PM<sub>2.5</sub> in children with a median age of 15 months (25th- 75th percentile, 9-25 mo) [13]. Chloride concentration has been used as a biomarker CFTR function in CF patients. It has been reported the CFTR participation in pneumocyte in chloride and fluid transport, it suggests the CFTR function in the pneumocyte layer could participate in the resolution of pulmonary edema [14]. Which is a pathological state, when increase permeability of the alveolar capillary barrier without changes in hydrostatic pressure, recurrent in burning smoke intoxication where many toxic products are formed including hydrogen chloride, chlorine, oxides, aldehydes, and particulate matter [15]. CFTR expression in rat isolated alveolar pneumocyte type-II [16], in alveoli from adult healthy human and alveoli from distal airway disease [17] and basal expression of CFTR in the lung adenocarcinoma cell line A549 (<https://www.proteinatlas.org/ENSG00000001626-CFTR/cell>) have been reported [18]. In addition, extensive analysis of punctual mutation on CFTR gene sequence has allowed importance in tissues and organs dysfunction, however, the gene-environment interaction on CFTR transcript and protein continues being subject of study.

In this sense, CFTR might be a molecular target of air pollution, the changes in CFTR could associate an aggravating factor susceptibility of alveolar epithelium with the further development of pulmonary disease or compromise physio-pathological condition of other illnesses. Nguyen et al., (2021) reports the effect of Diesel exhaust particles and tobacco smoke extract in airway epithelial cells (Calu-3) on CFTR expression and dysfunction, but not by the exposure to urban particulate matter [19]. The present study evaluates whether urban PM exposure can modify CFTR mRNA and protein in the *in vitro*, A549 cell line, and in the lungs and non-respiratory tissues, such kidneys, of rats exposed to concentrated PM.

## Materials and Methods

We investigated the effect of PM<sub>10</sub> and PM<sub>2.5</sub> collected in subway microenvironment and outdoor from Mexico City to explore whether CFTR is a molecular target of PM in the cell line A549. To corroborate the observations *in vitro*, we analyzed changes in CFTR mRNA and protein levels in lungs and in non-respiratory target of PM, kidney tissues, of rats exposed to whole body-chamber through a particulate concentrator.

### *In vitro* Exposure

Particles were provided by Dr. Andrea de Vizcaya CINVESTAV-IPN. Particles were collected in September-October 2009 from subway microenvironment and outdoor of Azcapotzalco subway station. PM<sub>10</sub> and PM<sub>2.5</sub> were obtained with a HiVol sampler, nitrocellulose filters were scraped off in order to obtain the particles [20].

Human lung adenocarcinoma cells, A549, were obtained from ATCC. The A549 cell line was subcultured in F-12/Ham culture media (Pyruvate 0.05 mg/mL, 2 mM L-glutamine, penicillin/streptomycin 50 IU/mL and /50g/mL, from Sigma Aldrich, USA) with 10% inactivated FBS (GIBCO, Thermo Fisher Scientific, USA). Cells were maintained at 37°C in a 5% CO<sub>2</sub> incubator. Cells were then plated at a density of 45 x 10<sup>3</sup> cells/cm<sup>2</sup> 24 h before PM exposure. Exposure media consisted of F-12 Ham supplemented with 0.5% FBS and PM solutions. The A549 cell line was exposed 20 and 40 µg/cm<sup>2</sup> of PM<sub>10</sub> and PM<sub>2.5</sub> for 24 h, which are amount equal or below annual average established by the official Mexican standard [21]. LDH assay was performed to corroborate cytotoxicity (Sigma Aldrich, USA). All assays were performed in triplicate with at least three independent experiments.

### *In vivo* Exposure

Sprague Dawley male rats (250 ± 40 g) were acutely (5 hours per day for 3 days) and subchronically (5 hours per day, 5 days per week, for 8 weeks) exposed to PM<sub>2.5</sub> using a particulate concentrator at CINVESTAV-IPN in the North of Mexico City during months of April-July 2013. Animal care and use were carried out according to the "Principles of Laboratory Animal Care" [22] and Mexican guidelines (NOM-062-ZOO-1999) [23]. All procedures were approved by the institutional ethics committee for the care and use of animals. Concentration in chambers by particle concentrator was 445 and 375 µg/m<sup>3</sup> in acute and subchronic exposure, respectively. After each exposure, urine from exposed animals was harvested 12 hours after being collected from a metabolic cage, followed by 15 min centrifugation at 1,000 rpm, and preserved at -80°C for analysis. Twenty-four hours after the last exposure, anesthetized animals (40 mg/kg sodic pentobarbital) were sacrificed by exsanguination. Blood was obtained by abdominal aorta puncture and collected with a gel activator tube, lungs and kidney cortex tissue were dissected. Serum was separated after centrifugation at 1,500 rpm for 15 min and then aliquoted and preserved at -80°C until their analysis.

## Chloride Determination

Chloride concentrations in cell lysates obtained by osmotic shock with ultrapure MilliQ-water, serum, and urine were analyzed spectrophotometrically at 500 nm by the Mercury (II)-Thiocyanate method using a commercial kit (CL1645, Randox, UK) and read in a TECAN Infinite 200 PRO reader.

## CFTR mRNA Expression

Total RNA was isolated by TRIzol reagent (Invitrogen, Thermo Fisher Scientific, USA) from cell cultures and lung and kidney cortex tissues. We then subjected 10 µg of total RNA to DNase treatment (Ambion, Turbo DNase-free™, Life Technologies, USA). cDNA synthesis was performed with 3 µg of total-RNA using Super Script II (Invitrogen, Thermo Fisher Scientific, USA). End point PCR was performed with the following CFTR oligonucleotides: forward 5'-TTCTGCCTCTTACCATATTTGACT-3' and reverse 5'-GCCTTCAGAT-TCCAGTTGTTTG-3', and we used actin and GADPH as housekeeping genes; GADPH forward 5'-ACCACAGTCCATGCCATCAC-3' and GADPH reverse 5'-TGCCAGTGAGCTTCCCGTT-3'; and Actin forward 5'-AC-TATCGGCAATGAGCGGTTCC-3' and Actin reverse 5'-AGCACTGTGTTG-GCATAGAGGTC-3'.

## CFTR Immunodetection

Total protein from A549 cells exposed to 0 and 20 µg/cm<sup>2</sup> of the different PMs was obtained with Nonidet-P40 buffer (150 mM NaCl, 1 % NP-40, 50 mM Tris-HCl pH 8.0 and protease inhibitors) and lysed using an insulin syringe with an ultrafine 6 mm needle, centrifuged at 10,000 rpm and 4 °C, and the supernatant was collected and stored at -70 °C until use. Lung and kidney cortex tissues were homogenized with Nonidet-P40 buffer and centrifuged at 8,000 rpm and 4 °C. Protein concentrations were determined by Bradford protein assay. The total protein (20 µg) per sample was loaded in SDS-polyacrylamide gels and transferred in wet transfer units to a PVDF membrane at a constant voltage (25 V) and kept overnight (12 hours). Membranes were then blocked for 1 h with 5% non-fat milk and incubated overnight with the primary antibodies Anti-CFTR (1:400, ProSci, inc. 7717) and GADPH (1:1000, Santa Cruz Biotechnology SC-32233). HRP secondary antibodies Goat Anti-Rabbit (BioRad, 1706515) and

Goat Anti-Mouse (BioRad, 1706516) were incubated for 1 h at a dilution of 1:10,000 to detect CFTR and GADPH, respectively. Immunoreactivity was detected using Luminata forte western blotting detection reagent (Millipore, WBLUF0500, Germany). Bands were visualized with a UVP EC3 imaging system (UVP Inc, USA). Finally, a semiquantitative analysis was performed with ImageJ free online software and expressed as arbitrary units.


## Statistical Analysis

The normality of all data was assessed with a Kolmogorov-Smirnov test. We then performed one-way analyses of variance (ANOVA) or Kruskal-Wallis tests followed by Holm-Sidak or Dunn's multiple comparisons post-hoc tests, respectively, to compare treatment groups with respect to the control group. All statistical analyses were carried out in GraphPad Prism version 6.00 for Windows (GraphPad Software, La Jolla California USA).

## Results


### Effect of Ambient Particles on CFTR Expression in vitro

After 24 hours of exposure to subway microenvironment and outdoor PM<sub>2.5</sub> and PM<sub>10</sub> particles, we observed cytotoxicity below 10% with respect to the control group (Figure 1a). Subway microenvironment PM<sub>2.5</sub> at 20 and 40 µg/cm<sup>2</sup> decreased CFTR expression, and a similar but less pronounced effect was observed with outdoor PM<sub>2.5</sub> (Figure 1b). Subway PM<sub>10</sub> at 40 µg/cm<sup>2</sup> induce an increment statistically significant on CFTR mRNA, however, outdoor PM<sub>10</sub> showed a statistical significant decrease at 20 µg/cm<sup>2</sup>, but not at 40 µg/cm<sup>2</sup> (Figure 1b). To corroborate the effect of the different particles on CFTR expression, we immunodetected CFTR on the total protein of A549 cells. The results show that CFTR protein levels decreased with respect to the control group 24 hours after exposure to 20 µg/cm<sup>2</sup>, independently of the origin and size of the particle (Figure 2a). One proper function of CFTR is maintaining the balance between the intra and extracellular chloride ions concentration. Thus, we evaluated intracellular chloride ion concentrations in cell lysates obtained by osmotic shock. Chloride concentration increased statistically significantly into the cell lysates of cultures exposed to PM at 20 µg/cm<sup>2</sup> in all particles tested, respect to the control cell group (Figure 2b).



**Figure 1:** CFTR mRNA expression is origin-dependent of the particle: Particulate concentrations were evaluated previously to CFTR mRNA expression; LDH assay demonstrates that PM<sub>10</sub> from different environments are the most toxic a). Subway microenvironment and outdoor PM<sub>2.5</sub> induce a down-regulation of CFTR mRNA. However, PM<sub>10</sub> from subway air induces the CFTR mRNA expression and outdoor PM<sub>10</sub> decreases CFTR expression.

b). Particle concentrations are expressed as micrograms per square centimeter (µg/cm²). Bar graphs show average ± standard deviation. One way-ANOVA test follows Holm-Sidak multiple comparisons test. \*Indicate significant statistical differences respect control group, zero particle concentration.




**Figure 2:** Subway microenvironment and outdoor particles induce CFTR protein down-regulation and intracellular chloride accumulation. A549 cell cultures lysates of exposed twenty-four hours to subway microenvironment and outdoor particles were analyzed by SDS-PAGE to immunodetect CFTR protein; data showing average ± standard deviation, One-way ANOVA with post hoc Fisher's LSD test. All particles at 20 µg/cm², independently of source, induce a decrement of CFTR protein levels a). Twenty-four hours after exposure to subway microenvironment or outdoor PM<sub>2.5</sub> and PM<sub>10</sub> at 20 µg/cm², cells were lysate by osmotic shock with MilliQ water. Data show as Box plot graph, Kruskal Wallis test follows Dunn's multiple comparisons test. Intracellular accumulation of chloride ion after PM exposure was observed b). \*Indicate significant statistical differences respect control group, zero particle concentration (p < 0.05).


### Effect of PM<sub>2.5</sub> Exposure on CFTR Expression in vivo

To provide evidence that CFTR is a molecular target of particle toxicity, we evaluated CFTR mRNA levels in lung tissue after acute (3 days, 5 hour/day) and subchronic (8 weeks, 5 days/week, 5 hour/day) exposure. We observed a time-dependent response of CFTR mRNA levels (Figure 3). Acute exposure resulted in a statistically significant two-fold increment in CFTR mRNA ( $p < 0.05$ ; Figure 3a); however, subchronic exposure to PM<sub>2.5</sub> reduced statistically significant CFTR mRNA to half with respect to the control group ( $p < 0.05$ ; Figure 3b). CFTR also has a function in other types of tissues, such as the pancreas and kidneys [24]. Therefore, we evaluated CFTR mRNA in kidney cortex tissue only in subchronic exposure to PM<sub>2.5</sub> and observed a statistically significant reduction in CFTR mRNA in the particle-ex-

posed group ( $p < 0.05$ , Figure 3c), suggesting that a repeated exposure to PM<sub>2.5</sub> during long periods can affect CFTR expression in non-respiratory tissues such as the kidneys. However, we also observed a statistically significant reduction ( $p < 0.05$ ) in CFTR protein expression in lung tissue after acute and subchronic exposure to PM<sub>2.5</sub> (Figure 3d, 3e, respectively), as well as in kidney tissue (Figure 3f). CFTR protein levels were modified by PM<sub>2.5</sub> exposure, suggesting changes in ionic transport such chloride anion. We observed significant statistical differences ( $p < 0.05$ ) in serum and urine chloride concentrations in both exposure periods (Figure 4). Our results show an increase in chloride ion concentrations in serum (Figure 4a) and urine (Figure 4c) after acute exposure to PM<sub>2.5</sub>; however, subchronic exposure decreased chloride ion concentrations in both fluids (Figure 4b and 4d).



**Figure 3:** Exposure to PM<sub>2.5</sub> *in vivo* modulates CFTR mRNA and protein levels in lung and non-respiratory tissues, kidney. After acute (3 days, 5 hours/ day) a) and d), and subchronic (8 weeks, 4 days/week, 5 hours/ day) b) and e) exposure to PM<sub>2.5</sub>, CFTR mRNA expression in lung tissue was evaluated by semiquantitative PCR and Western blot, respectively. In rat kidney tissue, CFTR mRNA and protein expression were evaluated by semiquantitative PCR and Western blot, respectively, only after subchronic exposure, c) and f). Graphs show the mean ± SD, n= 6 per group. \*Indicates statistical differences respect control group, filtered air (FA), Student's t-test ( $p < 0.05$ ).



**Figure 4:** Acute and subchronic exposure to PM<sub>2.5</sub> induce changes in serum and urine chloride concentration. Male rats were exposed to PM<sub>2.5</sub> by particulate matter concentrator. After acute (3 days, 5 hours/ day) and subchronic (8 weeks, 4 days/week, 5 hours/ day) exposure, animals were placed in metabolic cages to collect the 12 h-urine. After the last exposure, 24 h, the animals were sacrificed, and blood was collected to obtain serum. Acute exposure to PM<sub>2.5</sub> shows an increment statistical significance in serum and urine, a) and c), respectively. However, subchronic exposure to PM<sub>2.5</sub> induces a minor concentration of serum b) and urine chloride d). Boxplot of six animals per group, U-Mann Whitney test. \*Indicates statistical differences respect control group, filtered air (FA), (p<0.05).

## Discussion

CFTR has been studied at different molecular levels, such as genome, transcript, proteomic, and cellular function. However, the effect of ambient factors, such as atmospheric PM, on CFTR expression was recently reported with effects on CFTR expression and function in Calu-3 cell line exposed to tobacco smoke extract and diesel exhaust particles. On the other hand, our study exposed A549 cell line and a rodent model we complement the findings reported by Nguyen, et al. [19], we observed that urban particles can impact CFTR expression on lungs and in extrapulmonary tissues in time-dependent course. The in vitro results provide evidence that CFTR mRNA levels respond to PM<sub>10</sub> and PM<sub>2.5</sub> from subway and outdoor environments. Particle size is related to the source, and chemical composition; however, independently of that factors marginal or statistical significant differences were observed in the cells exposed to PM<sub>10</sub> and PM<sub>2.5</sub> on

CFTR mRNA. Bargon, et al. [25] has been reported that divalent cations as calcium and magnesium can down regulate CFTR mRNA in human bronchial cells [25]. Differences between PM<sub>10</sub> and PM<sub>2.5</sub> on CFTR mRNA could be influenced by the variances in the presence and concentrations of anions and cations by type and the particle origin. Regardless of the effect of particles on CFTR mRNA, all particles reduce statistical significant CFTR protein expression, concomitant with the retention of intracellular chloride ions under non-cytotoxic concentrations, cell viability minor to 10% through LDH assay. Chloride transport has been demonstrated in human and rat pneumocyte type-II alveolar epithelial. Apical pharmacological treatment of pneumocyte type-II monolayers with forskolin induce Cl<sup>-</sup> influx, which is inhibited by the CFTR<sub>inh</sub>-172, when high apical and low basal chloride concentrations are disposed in transwell assays [14]. Besides, it has been demonstrated that CFTR in pneumocyte type-II, through patch-clamp studies, can regulate epithelial Na<sup>+</sup> channel (ENaC), even when

CFTR protein were minimal [26], as well other chloride channels of SLC26 family [27]. The pneumocyte exposition to PM affect CFTR expression contributing to change the chloride and fluid transport, because the presence of ions (e.g. chloride and sodium) as part of chemical PM composition are dissolved in the pulmonary lining fluid modulated apically to CFTR. This effect could be contributed to chloride influx or water resorption across the alveolar epithelium modified transcellular transport.

The CFTR expression reduction has been reported by cigarette smoke (CS) exposure, and the reduced chloride secretion by CS was reported since 80's, before the description of CFTR gene, later was demonstrated the CS inhibit CFTR and induce its internalization, producing dehydration of liquid surface of airspace, a propound effect to development COPD [28]. Furthermore, the present results imply that CFTR is a molecular target of airborne PM and suggest that this effect occurs at different levels, such as mRNA, protein, and ion transport. CFTR regulation appears to be independent of particle size and chemical composition. The effect of PM<sub>2.5</sub> on CFTR was confirmed in a rodent model. It is important to note that the effect was time-dependent, where repeated exposure chronically induced a down-regulation of CFTR mRNA. Additionally, we also observed a reduction on CFTR mRNA in non-respiratory tissues, the kidneys, which are affected in CF, and their function decline after PM<sub>2.5</sub> exposure (4). Serum chloride concentrations corresponded to CFTR mRNA levels but not to CFTR protein expression, which suggests a difference in the temporality of the transcription, protein synthesis, and possibly channel activity.

Xu et al. [29] recently reported that the exposure to PM<sub>2.5</sub> is a potential inductor of pulmonary fibrosis in *in vitro* and *in vivo* models by activating epithelial mesenchymal transition (EMT), and TGFβ1/SMAD3 pathway, and increasing α-smooth muscle actin and Collagen type I [29]. Adding to this evidence, our results suggest that PM from different sources in non-cytotoxic concentrations can deregulate CFTR at different molecular levels, promoting an alteration in fluid secretion, ion transport, and it also likely contributes to EMT. It is significant to remark that the effect of PM on CFTR could contribute to the loss of glutathione transport, which is the most abundant antioxidant in lung epithelium lining fluid, and responsible for decreasing and attenuating oxidative stress and inflammation against toxic inhalants, such CS, [30] and which is reduced in CF patients [31]. One of our most relevant results is the CFTR down-regulation in kidney cortex tissue after subchronic exposure to PM<sub>2.5</sub>, which suggests that PM has an impact on other types of tissues and physiological functions of CFTR. In this sense, the effect of PM on CFTR expression in non-pulmonary tissues might contribute to the development of other diseases. Metabolic pathologies such as Diabetes, the most common extrapulmonary complications of CF, since epidemiological studies have found an association between an increase in PM particles and the number of cases of DM [32,33].

Is well clinically documented Cystic Fibrosis-Related Diabetes (CFRD) as a frequent comorbidity of CF by CFTR impairment [34,35]. Further studies are needed to verify the effect of subchronic or chronic effect of PM on CFTR on pancreatic function. The impairment, severity, and lack of control of diseases caused by the exposure to air pollution has been reported in other pulmonary genetic diseases, such α1-antitrypsin deficiency, which predisposes to the development of chronic obstructive pulmonary disease (COPD); PM<sub>10</sub> is associated with a decline in FEV1 and DLCO tests [36, 37]. Further epidemiological studies are necessary to demonstrate how the exposure to PM on CFTR can contribute to the impairment, increase in severity, and diseases control. The present experimental evidence demonstrates that CFTR is a toxicological target of PM *in vitro* and *vivo* models, affecting CFTR mRNA and protein expression, which could be related to chloride ion transport and probably with other physiological functions. These results imply a risk of developing diseases because of CFTR down-regulation by PM exposure.

## Acknowledgements

The authors acknowledge to Toxicology Department from CINVESTAV-IPN for the equipment and infrastructure to carry out the present study.

## Author Contributions

Aztatzí-Aguilar OG performed the *in vivo* and *in vitro* exposure, analyzed the data and wrote the draft manuscript; Cruz-Guerrero M carried out the PCR analysis; Debray-García Y performed Western Blot and wrote the draft manuscript; Uribe-Ramírez TM participated in the *in vivo* and *in vitro* exposure; Narváez-Morales J participated in the metabolic cage and chloride quantification; Barbier O provided the resources to analyze samples by Western Blot and PCR.

## Funding and/or Conflicts of Interests/Competing Interests.

The authors declare that no funds, grants, or other support were received during the preparation of this manuscript. The authors have no relevant financial or non-financial interests to disclose.

## References

1. Mills NL, Ken Donaldson, Paddy WH, Nicholas AB, William M, et al. (2009) Adverse cardiovascular effects of air pollution. *Nature Reviews Cardiology* 6(1): 36-44.
2. Kelly FJ, Fussell JC (2011) Air pollution and airway disease. *Clinical & Experimental Allergy* 41(8): 1059-1071.
3. Calderon GL, William R, Robert RM, Carlos HR, Ricardo DC, et al. (2004) Brain inflammation and Alzheimer's-like pathology in individuals exposed to severe air pollution. *Toxicologic pathology*. 32(6): 650-658.
4. Mehta AJ, Antonella Z, Marie ACB, Itai Kloog, Petros K, et al. (2016) Long-term exposure to ambient fine particulate matter and renal function in older men: The Veterans Administration Normative Aging Study. *Environmental health perspectives* 124(9): 1353-1360.

5. Rajagopalan S, RD Brook (2012) Air pollution and type 2 diabetes: mechanistic insights. *Diabetes* 61(12): 3037-3045.
6. Milici A, K Talavera (2021) TRP Channels as Cellular Targets of Particulate Matter. *International Journal of Molecular Sciences* 22(5): 2783.
7. Grebert C, F Becq, C Vandebrouck (2019) Focus on TRP channels in cystic fibrosis. *Cell calcium* 81: 29-37.
8. Kogan I, Mohabir R, Canhui Li, Jackie FK, Yanchun W, et al. (2003) CFTR directly mediates nucleotide-regulated glutathione flux. *The EMBO journal* 22(9): p. 1981-1989.
9. Linsdell P, JW, Hanrahan (1998) Glutathione permeability of CFTR. *American Journal of Physiology-Cell Physiology* 275(1): C323-C326.
10. Collins FS (1992) Cystic fibrosis: molecular biology and therapeutic implications. *Science* 256(5058): 774-779.
11. Castellani C, BM Assael (2017) Cystic fibrosis: a clinical view. *Cellular and Molecular Life Sciences* 74(1): 129-140.
12. Vankeerberghen A, H Cuppens, JJ Cassiman (2002) The cystic fibrosis transmembrane conductance regulator: an intriguing protein with pleiotropic functions. *Journal of Cystic Fibrosis* 1(1): 13-29.
13. Psoter KJ, Anneclaire JDR, Jonathan DM, Joel DK, Jon Wakefield, et al. (2015) Fine particulate matter exposure and initial *Pseudomonas aeruginosa* acquisition in cystic fibrosis. *Annals of the American Thoracic Society* 12(3): 385-391.
14. Fang X, Yuanlin S, Jan Hirsch, Luis JVG, Nicoletta P, et al. (2006) Contribution of CFTR to apical-basolateral fluid transport in cultured human alveolar epithelial type II cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 290(2): L242-L249.
15. Witten ML, Quan SF, Sobonya RE, Lemen RJ (1988) New developments in the pathogenesis of smoke inhalation-induced pulmonary edema. *Western Journal of Medicine* 148(1): 33-36.
16. Brochiero E, André Dagenais, Anik P, Yves B, Ryszard G (2004) Evidence of a functional CFTR Cl<sup>-</sup> channel in adult alveolar epithelial cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 287(2): L382-L392.
17. Engelhardt JF, M Zepeda, JA Cohn, JR Yankaskas, JM Wilson (1994) Expression of the cystic fibrosis gene in adult human lung. *The Journal of clinical investigation* 93(2): 737-749.
18. Thul PJ, C Lindskog (2018) The human protein atlas: A spatial map of the human proteome. *Protein Science* 27(1): 233-244.
19. Nguyen JP, Ryan D Huff, Quynh T Cao, Nicholas Tiessen, Christopher Carlsen, et al. (2021) Effects of environmental air pollutants on CFTR expression and function in human airway epithelial cells. *Toxicology in Vitro* 77: 105253.
20. Alfaro ME, Victor Torres, Javier Miranda, Leticia Martínez, Claudia GC, et al. (2009) Induction of IL-6 and inhibition of IL-8 secretion in the human airway cell line Calu-3 by urban particulate matter collected with a modified method of PM sampling. *Environmental research* 109(5): 528-535.
21. MEXICANOS P (2014) NORMA Oficial Mexicana NOM-025-SSA1-2014, Salud ambiental. Valores límite permisibles para la concentración de partículas suspendidas PM10 y PM2.5 en el aire ambiente y criterios para su evaluación. Al margen un sello con el Escudo Nacional, que dice: Estados Unidos Mexicanos.-Secretaría de Salud.
22. (1985) Health N.I.O. Guide for the care and use of laboratory animals.: National Academies.
23. (1999) NOM-062-ZOO, N.O.M. and E.T.P. LA PRODUCCION, Norma Oficial Mexicana NOM-062-ZOO-1999: Especificaciones técnicas para la producción, cuidado y uso de los animales de laboratorio, Diario Oficial de la Federación México.
24. Ideozu JE, Xi Zhang, Susanna McColley, Hara Levy (2019) Transcriptome profiling and molecular therapeutic advances in cystic fibrosis: recent insights. *Genes* 10(3): 180.
25. Bargon, J, BC Trapnell, CS Chu, ER Rosenthal, K Yoshimura, et al. (1992) Down-regulation of cystic fibrosis transmembrane conductance regulator gene expression by agents that modulate intracellular divalent cations. *Molecular and cellular biology*. 12(4): 1872-1878.
26. Lazrak A, Asta Jurkuvenaitė, Lan Chen, Kim M Keeling, James F Collawn, et al. (2011) Enhancement of alveolar epithelial sodium channel activity with decreased cystic fibrosis transmembrane conductance regulator expression in mouse lung. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 301(4): L557-67.
27. Alper SL, AK Sharma (2013) The SLC26 gene family of anion transporters and channels. *Molecular aspects of medicine* 34(2-3): 494-515.
28. Saint Ciri V, MA Gray (2017) Role of CFTR in epithelial physiology. *Cellular and Molecular Life Sciences* 74(1): 93-115.
29. Xu Z, Zilin Li, Ziyi Liao, Sumeng Gao, Li Hua, et al. (2019) PM2.5 induced pulmonary fibrosis in vivo and in vitro. *Ecotoxicology and environmental safety* 171: 112-121.
30. Gould NS, Elysia Min, Richard J Martin, Brian J Day (2012) CFTR is the primary known apical glutathione transporter involved in cigarette smoke-induced adaptive responses in the lung. *Free Radical Biology and Medicine* 52(7): 1201-1206.
31. Roum J, R Buhl, NG McElvaney, Z Borok, RG Crystal (1993) Systemic deficiency of glutathione in cystic fibrosis. *Journal of applied physiology* 75(6): 2419-2424.
32. Eze IC, Lars G Hemkens, Heiner C Bucher, Barbara Hoffmann, Christian Schindler, et al. (2015) Association between ambient air pollution and diabetes mellitus in Europe and North America: systematic review and meta-analysis. *Environmental health perspectives* 123(5): 381-389.
33. Brook RD, Michael Jerrett, Jeffrey R Brook, Robert L Bard, Murray M Finkelstein (2008) The relationship between diabetes mellitus and traffic-related air pollution. *Journal of occupational and environmental medicine* 50(1): 32-38.
34. Norris AW (2019) Is Cystic Fibrosis-related Diabetes Reversible? New Data on CFTR Potentiation and Insulin Secretion. *Am J Respir Crit Care Med* 199(3): 261-263.
35. Ntimbane T, Blandine Comte, Geneviève Mailhot, Yves Berthiaume, Vincent Poitout, et al. (2009) Cystic fibrosis-related diabetes: from CFTR dysfunction to oxidative stress. *The Clinical Biochemist Reviews* 30(4): 153-177.
36. Wood AM, RM Harrison, S Semple, JG Ayres, RA Stockley (2009) Outdoor air pollution is associated with disease severity in  $\alpha$ 1-antitrypsin deficiency. *European Respiratory Journal* 34(2): 346-353.
37. Wood AM, Roy M Harrison, Sean Semple, Jon G Ayres, Robert A Stockley (2010) Outdoor air pollution is associated with rapid decline of lung function in  $\alpha$ 1-antitrypsin deficiency. *Occupational and environmental medicine* 67(8): 556-561.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2026.67.010432](https://doi.org/10.26717/BJSTR.2026.67.010432)

Aztatzi Aguilar OG . Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>